

4-2015

The Atiyah Class of a dg-Vector Bundle

Rajan Amit Mehta

Smith College, rmehta@smith.edu

Mathieu Stiénon

The Pennsylvania State University

Ping Xu

The Pennsylvania State University

Follow this and additional works at: https://scholarworks.smith.edu/mth_facpubs

Part of the [Mathematics Commons](#)

Recommended Citation

Mehta, Rajan Amit; Stiénon, Mathieu; and Xu, Ping, "The Atiyah Class of a dg-Vector Bundle" (2015). Mathematics and Statistics: Faculty Publications, Smith College, Northampton, MA.
https://scholarworks.smith.edu/mth_facpubs/16

This Article has been accepted for inclusion in Mathematics and Statistics: Faculty Publications by an authorized administrator of Smith ScholarWorks.
For more information, please contact scholarworks@smith.edu

THE ATIYAH CLASS OF A DG-VECTOR BUNDLE

RAJAN AMIT MEHTA, MATHIEU STIÉNON, AND PING XU

En hommage à Charles-Michel Marle à l'occasion de son quatre-vingtième anniversaire

ABSTRACT. We introduce the notions of Atiyah class and Todd class of a differential graded vector bundle with respect to a differential graded Lie algebroid. We prove that the space of vector fields $\mathfrak{X}(\mathcal{M})$ on a dg-manifold \mathcal{M} with homological vector field Q admits a structure of $L_\infty[1]$ -algebra with the Lie derivative L_Q as unary bracket λ_1 , and the Atiyah cocycle $\text{At}_{\mathcal{M}}$ corresponding to a torsion-free affine connection as binary bracket λ_2 .

1. DG-MANIFOLDS AND DG-VECTOR BUNDLES

A \mathbb{Z} -graded manifold \mathcal{M} with base manifold M is a sheaf of \mathbb{Z} -graded, graded-commutative algebras $\{\mathcal{R}_U | U \subset M \text{ open}\}$ over M , locally isomorphic to $C^\infty(U) \otimes \hat{S}(V^\vee)$, where $U \subset M$ is an open submanifold, V is a \mathbb{Z} -graded vector space, and $\hat{S}(V^\vee)$ denotes the graded algebra of formal polynomials on V . By $C^\infty(\mathcal{M})$, we denote the \mathbb{Z} -graded, graded-commutative algebra of global sections. By a dg-manifold, we mean a \mathbb{Z} -graded manifold endowed with a homological vector field, i.e. a vector field Q of degree $+1$ satisfying $[Q, Q] = 0$.

Example 1.1. Let $A \rightarrow M$ be a Lie algebroid over \mathbb{C} . Then $A[1]$ is a dg-manifold with the Chevalley–Eilenberg differential d_{CE} as homological vector field. In fact, according to Vaintrob [12], there is a bijection between the Lie algebroid structures on the vector bundle $A \rightarrow M$ and the homological vector fields on the \mathbb{Z} -graded manifold $A[1]$.

Example 1.2. Let s be a smooth section of a vector bundle $E \rightarrow M$. Then $E[-1]$ is a dg-manifold with the contraction operator i_s as homological vector field.

Example 1.3. Let $\mathfrak{g} = \sum_{i \in \mathbb{Z}} \mathfrak{g}_i$ be a \mathbb{Z} -graded vector space of finite type, i.e. each \mathfrak{g}_i is a finite-dimensional vector space. Then $\mathfrak{g}[1]$ is a dg-manifold if and only if \mathfrak{g} is an L_∞ -algebra.

A dg-vector bundle is a vector bundle in the category of dg-manifolds. We refer the reader to [10, 4] for details on dg-vector bundles. The following example is essentially due to Kotov–Strobl [4].

Example 1.4. Let $A \rightarrow M$ be a gauge Lie algebroid with anchor ρ . Then $A[1] \rightarrow T[1]M$ is a dg-vector bundle, where the homological vector fields on $A[1]$ and $T[1]M$ are the Chevalley–Eilenberg differentials.

Research partially supported by NSF grants DMS1406668, and NSA grants H98230-06-1-0047 and H98230-14-1-0153.

The example above is a special case of a general fact [10], that LA-vector bundles [6, 7, 8] (also known as VB-algebroids [2]) give rise to dg-vector bundles.

Given a vector bundle $\mathcal{E} \xrightarrow{\pi} \mathcal{M}$ of graded manifolds, its space of sections, denoted $\Gamma(\mathcal{E})$, is defined to be $\bigoplus_{j \in \mathbb{Z}} \Gamma_j(\mathcal{E})$, where $\Gamma_j(\mathcal{E})$ consists of degree preserving maps $s \in \text{Hom}(\mathcal{M}, \mathcal{E}[-j])$ such that $(\pi[-j]) \circ s = \text{id}_{\mathcal{M}}$, where $\pi[-j] : \mathcal{E}[-j] \rightarrow \mathcal{M}$ is the natural map induced from π ; see [10] for more details. When $\mathcal{E} \rightarrow \mathcal{M}$ is a dg-vector bundle, the homological vector fields on \mathcal{E} and \mathcal{M} naturally induce a degree 1 operator \mathcal{Q} on $\Gamma(\mathcal{E})$, making $\Gamma(\mathcal{E})$ a dg-module over $C^\infty(\mathcal{M})$. Since the space $\Gamma(\mathcal{E}^\vee)$ of linear functions on \mathcal{E} generates $C^\infty(\mathcal{E})$, the converse is also true.

Lemma 1.5. *Let $\mathcal{E} \rightarrow \mathcal{M}$ be a vector bundle object in the category of graded manifolds and suppose \mathcal{M} is a dg-manifold. If $\Gamma(\mathcal{E})$ is a dg-module over $C^\infty(\mathcal{M})$, then \mathcal{E} admits a natural dg-manifold structure such that $\mathcal{E} \rightarrow \mathcal{M}$ is a dg-vector bundle. In fact, the categories of dg-vector bundles and of locally free dg-modules are equivalent.*

In this case, the degree +1 operator \mathcal{Q} on $\Gamma(\mathcal{E})$ gives rise to a cochain complex

$$\cdots \rightarrow \Gamma_i(\mathcal{E}) \xrightarrow{\mathcal{Q}} \Gamma_{i+1}(\mathcal{E}) \rightarrow \cdots,$$

whose cohomology group will be denoted by $H^\bullet(\Gamma(\mathcal{E}), \mathcal{Q})$.

In particular, the space $\mathfrak{X}(\mathcal{M})$ of vector fields on a dg-manifold (\mathcal{M}, Q) (i.e. graded derivations of $C^\infty(\mathcal{M})$), which can be regarded as the space of sections $\Gamma(T\mathcal{M})$, is naturally a dg-module over $C^\infty(\mathcal{M})$ with the Lie derivative $L_Q : \mathfrak{X}(\mathcal{M}) \rightarrow \mathfrak{X}(\mathcal{M})$ playing the role of the degree +1 operator \mathcal{Q} .

Thus we have the following

Corollary 1.6. *For every dg-manifold (\mathcal{M}, Q) , the Lie derivative L_Q makes $\Gamma(T\mathcal{M})$ into a dg-module over $C^\infty(\mathcal{M})$ and therefore $T\mathcal{M} \rightarrow \mathcal{M}$ is naturally a dg-vector bundle.*

Following the classical case, the corresponding homological vector field on $T\mathcal{M}$ is called the *tangent lift* of Q .

Differential graded Lie algebroids are another useful notion. Roughly, a dg-Lie algebroid can be thought of as a Lie algebroid object in the category of dg-manifolds. For more details, we refer the reader to [10], where dg-Lie algebroids are called Q -algebroids.

Differential graded foliations constitute an important class of examples of dg-Lie algebroids.

Lemma 1.7. *Let $\mathcal{D} \subset T\mathcal{M}$ be an integrable distribution on a graded manifold \mathcal{M} . Suppose there exists a homological vector field Q on \mathcal{M} such that $\Gamma(\mathcal{D})$ is stable under L_Q . Then $\mathcal{D} \rightarrow \mathcal{M}$ is a dg-Lie algebroid with the inclusion $\rho : \mathcal{D} \rightarrow T\mathcal{M}$ as its anchor map.*

2. ATIYAH CLASS AND TODD CLASS OF A DG-VECTOR BUNDLE

Let $\mathcal{E} \rightarrow \mathcal{M}$ be a dg-vector bundle and let $\mathcal{A} \rightarrow \mathcal{M}$ be a dg-Lie algebroid with anchor $\rho : \mathcal{A} \rightarrow T\mathcal{M}$. An \mathcal{A} -connection on $\mathcal{E} \rightarrow \mathcal{M}$ is a degree 0 map $\nabla : \Gamma(\mathcal{A}) \otimes \Gamma(\mathcal{E}) \rightarrow \Gamma(\mathcal{E})$ such that

$$\nabla_{fX}s = f\nabla_X s$$

and

$$\nabla_X(fs) = \rho(X)(f)s + (-1)^{|X||f|}f\nabla_Xs$$

for all $f \in C^\infty(\mathcal{M})$, $X \in \Gamma(\mathcal{A})$, and $s \in \Gamma(\mathcal{E})$. Here we use the ‘absolute value’ notation to denote the degree of the argument. When we say that ∇ is of degree 0, we actually mean that $|\nabla_Xs| = |X| + |s|$ for every pair of homogeneous elements X and s . Such connections always exist since the standard partition of unity argument holds in the context of graded manifolds. Given a dg-vector bundle $\mathcal{E} \rightarrow \mathcal{M}$ and an \mathcal{A} -connection ∇ on it, we can consider the bundle map $\text{At}_\mathcal{E} : \mathcal{A} \otimes \mathcal{E} \rightarrow \mathcal{E}$ defined by

$$(1) \quad \text{At}_\mathcal{E}(X, s) := \mathcal{Q}(\nabla_Xs) - \nabla_{\mathcal{Q}(X)}s - (-1)^{|X|}\nabla_X(\mathcal{Q}(s)), \quad \forall X \in \Gamma(\mathcal{A}), s \in \Gamma(\mathcal{E}).$$

Proposition 2.1. (1) $\text{At}_\mathcal{E} : \mathcal{A} \otimes \mathcal{E} \rightarrow \mathcal{E}$ is a degree +1 bundle map and therefore can also be regarded as a degree +1 section of $\mathcal{A}^\vee \otimes \text{End } \mathcal{E}$.

(2) $\text{At}_\mathcal{E}$ is a cocycle: $\mathcal{Q}(\text{At}_\mathcal{E}) = 0$.

(3) The cohomology class of $\text{At}_\mathcal{E}$ is independent of the choice of the connection ∇ .

Thus there is a natural cohomology class $\alpha_\mathcal{E} := [\text{At}_\mathcal{E}]$ in $H^1(\Gamma(\mathcal{A}^\vee \otimes \text{End } \mathcal{E}), Q)$. The class $\alpha_\mathcal{E}$ is called the *Atiyah class* of the dg-vector bundle $\mathcal{E} \rightarrow \mathcal{M}$ relative to the dg-Lie algebroid $\mathcal{A} \rightarrow \mathcal{M}$.

The Atiyah class of a dg-manifold, which is the obstruction to the existence of connections compatible with the differential, was first investigated by Shoikhet [11] in relation with Kontsevich’s formality theorem and Duflo formula. More recently, the Atiyah class of a dg-vector bundle appeared in Costello’s work [1].

We define the *Todd class* $\text{Td}_\mathcal{E}$ of a dg-vector bundle $\mathcal{E} \rightarrow \mathcal{M}$ relative to a dg-Lie algebroid $\mathcal{A} \rightarrow \mathcal{M}$ by

$$(2) \quad \text{Td}_\mathcal{E} := \text{Ber} \left(\frac{1 - e^{-\alpha_\mathcal{E}}}{\alpha_\mathcal{E}} \right) \in \prod_{k \geq 0} H^k(\Gamma(\wedge^k \mathcal{A}^\vee), Q),$$

where Ber denotes the Berezinian [9] and $\wedge^k \mathcal{A}^\vee$ denotes the dg vector bundle $S^k(\mathcal{A}^\vee[-1])[k] \rightarrow \mathcal{M}$. One checks that $\text{Td}_\mathcal{E}$ can be expressed in terms of scalar Atiyah classes $c_k = \frac{1}{k!} \left(\frac{i}{2\pi} \right)^k \text{str } \alpha_\mathcal{E}^k \in H^k(\Gamma(\wedge^k \mathcal{A}^\vee), Q)$. Here $\text{str} : \text{End } \mathcal{E} \rightarrow C^\infty(\mathcal{M})$ denotes the supertrace. Note that $\text{str } \alpha_\mathcal{E}^k \in \Gamma(\wedge^k \mathcal{A}^\vee)$ since $\alpha_\mathcal{E}^k \in \Gamma(\wedge^k \mathcal{A}^\vee) \otimes_{C^\infty(\mathcal{M})} \text{End } \mathcal{E}$. If $\mathcal{A} = T\mathcal{M}$, we write $\Omega^k(\mathcal{M})$ instead of $\Gamma(\wedge^k T^\vee \mathcal{M})$.

3. ATIYAH CLASS AND TODD CLASS OF A DG-MANIFOLD

Consider a dg-manifold (\mathcal{M}, Q) . According to Lemma 1.7, its tangent bundle $T\mathcal{M}$ is indeed a dg-Lie algebroid. By the *Atiyah class of a dg-manifold* (\mathcal{M}, Q) , denoted $\alpha_\mathcal{M}$, we mean the Atiyah class of the tangent dg-vector bundle $T\mathcal{M} \rightarrow \mathcal{M}$ with respect to the dg-Lie algebroid $T\mathcal{M}$. Similarly, the Atiyah 1-cocycle of a dg manifold \mathcal{M} corresponding to an affine connection on \mathcal{M} (i.e. a $T\mathcal{M}$ -connection on $T\mathcal{M} \rightarrow \mathcal{M}$) is the 1-cocycle defined as in Eq. (1).

Lemma 3.1. *Suppose V is a vector space. The only connection on the graded manifold $V[1]$ is the trivial connection.*

Proof. Since the graded algebra of functions on $V[1]$ is $\wedge(V^\vee)$, every vector $v \in V$ determines a degree -1 vector field ι_v on $V[1]$, which acts as a contraction operator on $\wedge(V^\vee)$. The $C^\infty(V[1])$ -module of all vector fields on $V[1]$ is generated by its subset $\{\iota_v\}_{v \in V}$. It follows that a connection ∇ on $V[1]$ is completely determined

by the knowledge of $\nabla_{\iota_v \iota_w}$ for all $v, w \in V$. Since the degree of every vector field $\nabla_{\iota_v \iota_w}$ must be -2 and there are no nonzero vector fields of degree -2 , it follows that $\nabla_{\iota_v \iota_w} = 0$. \square

Given a finite-dimensional Lie algebra \mathfrak{g} , consider the dg-manifold (\mathcal{M}, Q) , where $\mathcal{M} = \mathfrak{g}[1]$ and Q is the Chevalley-Eilenberg differential d_{CE} . The following result can be easily verified using the canonical trivialization $T\mathcal{M} \cong \mathfrak{g}[1] \times \mathfrak{g}[1]$.

Lemma 3.2. *Let $(\mathcal{M}, Q) = (\mathfrak{g}[1], d_{\text{CE}})$ be the canonical dg-manifold corresponding to a finite-dimensional Lie algebra \mathfrak{g} . Then,*

$$H^k(\Gamma(T^\vee \mathcal{M} \otimes \text{End } T\mathcal{M}), Q) \cong H_{\text{CE}}^{k-1}(\mathfrak{g}, \mathfrak{g}^\vee \otimes \mathfrak{g}^\vee \otimes \mathfrak{g}),$$

and

$$H^k(\Omega^k(\mathcal{M}), Q) \cong (S^k \mathfrak{g}^\vee)^\mathfrak{g}.$$

Proposition 3.3. *Let $(\mathcal{M}, Q) = (\mathfrak{g}[1], d_{\text{CE}})$ be the canonical dg-manifold corresponding to a finite-dimensional Lie algebra \mathfrak{g} . Then the Atiyah class $\alpha_{\mathfrak{g}[1]}$ is precisely the Lie bracket of \mathfrak{g} regarded as an element of $(\mathfrak{g}^\vee \otimes \mathfrak{g}^\vee \otimes \mathfrak{g})^\mathfrak{g} \cong H^1(\Gamma(T^\vee \mathcal{M} \otimes \text{End } T\mathcal{M}), Q)$. Consequently, the isomorphism*

$$\prod_k H^k(\Omega^k(\mathcal{M}), Q) \xrightarrow{\cong} (\widehat{S}(\mathfrak{g}^\vee))^\mathfrak{g}$$

maps the Todd class $\text{Td}_{\mathfrak{g}[1]}$ onto the Duflo element of \mathfrak{g} .

Example 3.4. Consider a dg-manifold of the form $\mathcal{M} = (\mathbb{R}^{m|n}, Q)$. Let $(x_1, \dots, x_m; x_{m+1}, \dots, x_{m+n})$ be coordinate functions on $\mathbb{R}^{m|n}$, and write $Q = \sum_k Q_k(x) \frac{\partial}{\partial x_k}$. Then the Atiyah 1-cocycle associated to the trivial connection $\nabla_{\frac{\partial}{\partial x_i}} \frac{\partial}{\partial x_j} = 0$ is given by

$$(3) \quad \text{At}_{\mathcal{M}} \left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j} \right) = (-1)^{|x_i|+|x_j|} \sum_k \frac{\partial^2 Q_k}{\partial x_i \partial x_j} \frac{\partial}{\partial x_k}$$

As we can see from (3), the Atiyah 1-cocycle $\text{At}_{\mathcal{M}}$ includes the information about the homological vector field of second-order and higher.

4. ATIYAH CLASS AND HOMOTOPY LIE ALGEBRAS

Let \mathcal{M} be a graded manifold. A $(1, 2)$ -tensor of degree k on \mathcal{M} is a \mathbb{C} -linear map $\alpha : \mathfrak{X}(\mathcal{M}) \otimes_{\mathbb{C}} \mathfrak{X}(\mathcal{M}) \rightarrow \mathfrak{X}(\mathcal{M})$ such that $|\alpha(X, Y)| = |X| + |Y| + k$ and

$$\alpha(fX, Y) = (-1)^{k|f|} f\alpha(X, Y) = (-1)^{|f||X|} \alpha(X, fY).$$

We denote the space of $(1, 2)$ -tensors of degree k by $\mathcal{T}_k^{1,2}(\mathcal{M})$, and the space of all $(1, 2)$ -tensors by $\mathcal{T}^{1,2}(\mathcal{M}) = \bigoplus_k \mathcal{T}_k^{1,2}(\mathcal{M})$.

The torsion of an affine connection ∇ is given by

$$(4) \quad T(X, Y) = \nabla_X Y - (-1)^{|X||Y|} \nabla_Y X - [X, Y].$$

The torsion is an element in $\mathcal{T}_0^{1,2}(\mathcal{M})$. Given any affine connection, one can define its opposite affine connection ∇^{op} , given by

$$(5) \quad \nabla_X^{\text{op}} Y = \nabla_X Y - T(X, Y) = [X, Y] + (-1)^{|X||Y|} \nabla_Y X.$$

The average $\frac{1}{2}(\nabla + \nabla^{\text{op}})$ is a torsion-free affine connection. This shows that torsion-free affine connections always exist on graded manifolds.

In this section, we show that, as in the classical situation considered by Kapranov in [3, 5], the Atiyah 1-cocycle of a dg-manifold gives rise to an interesting homotopy Lie algebra. As in the last section, let (\mathcal{M}, Q) be a dg-manifold and let ∇ be an affine connection on \mathcal{M} . The following can be easily verified by direct computation.

- (1) The anti-symmetrization of the Atiyah 1-cocycle $\text{At}_{\mathcal{M}}$ is equal to $L_Q T$, so $\text{At}_{\mathcal{M}}$ is graded antisymmetric up to an exact term. In particular, if ∇ is torsion-free, we have

$$\text{At}_{\mathcal{M}}(X, Y) = (-1)^{|X||Y|} \text{At}_{\mathcal{M}}(Y, X).$$

- (2) The degree $1 + |X|$ operator $\text{At}_{\mathcal{M}}(X, -)$ need not be a derivation of the degree $+1$ ‘product’ $\mathfrak{X}(\mathcal{M}) \otimes_{\mathbb{C}} \mathfrak{X}(\mathcal{M}) \xrightarrow{\text{At}_{\mathcal{M}}} \mathfrak{X}(\mathcal{M})$. However, the Jacobiator $(X, Y, Z) \mapsto \text{At}_{\mathcal{M}}(X, \text{At}_{\mathcal{M}}(Y, Z)) - \{(-1)^{|X|+1} \text{At}_{\mathcal{M}}(\text{At}_{\mathcal{M}}(X, Y), Z) + (-1)^{(|X|+1)(|Y|+1)} \text{At}_{\mathcal{M}}(Y, \text{At}_{\mathcal{M}}(X, Z))\}$,

of $\text{At}_{\mathcal{M}}$, which vanishes precisely when $\text{At}_{\mathcal{M}}(X, -)$ is a derivation of $\text{At}_{\mathcal{M}}$, is equal to $\pm L_Q(\nabla \text{At}_{\mathcal{M}})$. Hence $\text{At}_{\mathcal{M}}$ satisfies the graded Jacobi identity up to the exact term $L_Q(\nabla \text{At}_{\mathcal{M}})$.

Armed with this motivation, we can now state the main result of this note.

Theorem 4.1. *Let (\mathcal{M}, Q) be a dg-manifold and let ∇ be a torsion-free affine connection on \mathcal{M} . There exists a sequence $(\lambda_k)_{k \geq 2}$ of maps $\lambda_k \in \text{Hom}(S^k(TM), TM[-1])$ starting with $\lambda_2 := \text{At}_{\mathcal{M}} \in \text{Hom}(S^2(TM), TM[-1])$ which, together with $\lambda_1 := L_Q : \mathfrak{X}(\mathcal{M}) \rightarrow \mathfrak{X}(\mathcal{M})$, satisfy the $L_{\infty}[1]$ -algebra axioms. As a consequence, the space of vector fields $\mathfrak{X}(\mathcal{M})$ on a dg-manifold (\mathcal{M}, Q) admits an $L_{\infty}[1]$ -algebra structure with the Lie derivative L_Q as unary bracket λ_1 and the Atiyah cocycle $\text{At}_{\mathcal{M}}$ as binary bracket λ_2 .*

To prove Theorem 4.1, we introduce a Poincaré–Birkhoff–Witt map for graded manifolds.

It was shown in [5] that every torsion-free affine connection ∇ on a smooth manifold M determines an isomorphism of coalgebras (over $C^{\infty}(M)$)

$$(6) \quad \text{pbw}^{\nabla} : \Gamma(S(TM)) \xrightarrow{\cong} D(M),$$

called the Poincaré–Birkhoff–Witt (PBW) map. Here $D(M)$ denotes the space of differential operators on M .

Geometrically, an affine connection ∇ induces an exponential map $TM \rightarrow M \times M$, which is a well-defined diffeomorphism from a neighborhood of the zero section of TM to a neighborhood of the diagonal $\Delta(M)$ of $M \times M$. Sections of $S(TM)$ can be viewed as fiberwise distributions on TM supported on the zero section, while $D(M)$ can be viewed as the space of source-fiberwise distributions on $M \times M$ supported on the diagonal $\Delta(M)$. The map $\text{pbw}^{\nabla} : \Gamma(S(TM)) \rightarrow D(M)$ is simply the push-forward of fiberwise distributions through the exponential map $\exp^{\nabla} : TM \rightarrow M \times M$ and is clearly an isomorphism of coalgebras over $C^{\infty}(M)$.

Even though, for a *graded* manifold \mathcal{M} endowed with a torsion-free affine connection ∇ , we lack an exponential map $\exp^{\nabla} : \mathcal{T}\mathcal{M} \rightarrow \mathcal{M} \times \mathcal{M}$, a PBW map can still be defined purely algebraically thanks to the iteration formula introduced in [5].

Lemma 4.2. *Let \mathcal{M} be a \mathbb{Z} -graded manifold and let ∇ be a torsion-free affine connection on \mathcal{M} . The Poincaré-Birkhoff-Witt map inductively defined by the relations¹*

$$\begin{aligned} \text{pbw}^\nabla(f) &= f, \quad \forall f \in C^\infty(\mathcal{M}); \\ \text{pbw}^\nabla(X) &= X, \quad \forall X \in \mathfrak{X}(\mathcal{M}); \end{aligned}$$

and

$$\begin{aligned} \text{pbw}^\nabla(X_0 \odot \cdots \odot X_n) &= \frac{1}{n+1} \sum_{k=0}^n (-1)^{|X_k|(|X_0| + \cdots + |X_{k-1}|)} \{ X_k \cdot \text{pbw}^\nabla(X_0 \odot \cdots \odot \widehat{X}_k \odot \cdots \odot X_n) \\ &\quad - \text{pbw}^\nabla(\nabla_{X_k}(X_0 \odot \cdots \odot \widehat{X}_k \odot \cdots \odot X_n)) \}, \end{aligned}$$

for all $n \in \mathbb{N}$ and $X_0, \dots, X_n \in \mathfrak{X}(\mathcal{M})$, establishes an isomorphism

$$(7) \quad \text{pbw}^\nabla : \Gamma(S(T\mathcal{M})) \xrightarrow{\cong} D(\mathcal{M}).$$

of coalgebras over $C^\infty(\mathcal{M})$.

Now assume that (\mathcal{M}, Q) is a dg-manifold. The homological vector field Q induces a degree +1 coderivation of $D(\mathcal{M})$ defined by the Lie derivative:

$$(8) \quad L_Q(X_1 \cdots X_n) = \sum_{k=1}^n (-1)^{|X_1| + \cdots + |X_{k-1}|} X_1 \cdots X_{k-1} [Q, X_k] X_{k+1} \cdots X_n.$$

Now using the isomorphism of coalgebras pbw^∇ as in Eq. (7) to transfer L_Q from $D(\mathcal{M})$ to $\Gamma(S(T\mathcal{M}))$, we obtain $\delta := (\text{pbw}^\nabla)^{-1} \circ L_Q \circ \text{pbw}^\nabla$, a degree 1 coderivation of $\Gamma(S(T\mathcal{M}))$. Finally, dualizing δ , we obtain an operator

$$D : \Gamma(\widehat{S}(T^\vee \mathcal{M})) \rightarrow \Gamma(\widehat{S}(T^\vee \mathcal{M}))$$

as

$$\Gamma(\widehat{S}(T^\vee \mathcal{M})) \cong \text{Hom}_{C^\infty(\mathcal{M})}(\Gamma(S(T\mathcal{M})), C^\infty(\mathcal{M})).$$

Theorem 4.3. *Let (\mathcal{M}, Q) be a dg-manifold and let ∇ be a torsion-free affine connection on \mathcal{M} .*

- (1) *The operator D , dual to $(\text{pbw}^\nabla)^{-1} \circ L_Q \circ \text{pbw}^\nabla$, is a degree +1 derivation of the graded algebra $\Gamma(\widehat{S}(T^\vee \mathcal{M}))$ satisfying $D^2 = 0$.*
- (2) *There exists a sequence $\{R_k\}_{k \geq 2}$ of homomorphisms $R_k \in \text{Hom}(S^k T\mathcal{M}, T\mathcal{M}[-1])$, whose first term R_2 is precisely the Atiyah 1-cocycle $\text{At}_{\mathcal{M}}$, such that $D = L_Q + \sum_{k=2}^{\infty} \widetilde{R}_k$, where \widetilde{R}_k denotes the $C^\infty(\mathcal{M})$ -linear operator on $\Gamma(\widehat{S}(T^\vee \mathcal{M}))$ corresponding to R_k .*

Finally we note that Theorem 4.1 is a consequence of Theorem 4.3.

ACKNOWLEDGEMENTS

We would like to thank several institutions for their hospitality while work on this project was being done: Penn State University (Mehta), and Université Paris Diderot (Xu). We also wish to thank Hsuan-Yi Liao, Dmitry Roytenberg and Boris Shoikhet for inspiring discussions.

¹We would like to thank Hsuan-Yi Liao for correcting a sign error in the inductive formula defining the map pbw^∇ .

REFERENCES

1. Kevin Costello, *A geometric construction of the Witten genus, I*, Proceedings of the International Congress of Mathematicians. Volume II, Hindustan Book Agency, New Delhi, 2010, pp. 942–959. MR 2827826
2. Alfonso Gracia-Saz and Rajan Amit Mehta, *Lie algebroid structures on double vector bundles and representation theory of Lie algebroids*, Adv. Math. **223** (2010), no. 4, 1236–1275. MR 2581370 (2011j:53162)
3. Mikhail Kapranov, *Rozansky-Witten invariants via Atiyah classes*, Compositio Math. **115** (1999), no. 1, 71–113. MR 1671737 (2000h:57056)
4. A. Kotov and T. Strobl, *Characteristic classes associated to Q -bundles*, ArXiv e-prints (2007), [arXiv:0711.4106](https://arxiv.org/abs/0711.4106).
5. C. Laurent-Gengoux, M. Stiénon, and P. Xu, *Kapranov dg-manifolds and Poincaré–Birkhoff–Witt isomorphisms*, ArXiv e-prints (2014), [arXiv:1408.2903](https://arxiv.org/abs/1408.2903).
6. K. C. H. Mackenzie, *Double Lie algebroids and the double of a Lie bialgebroid*, ArXiv Mathematics e-prints (1998), [arXiv:math/9808081](https://arxiv.org/abs/math/9808081).
7. K. C. H. Mackenzie, *Drinfel’d doubles and Ehresmann doubles for Lie algebroids and Lie bialgebroids*, Electron. Res. Announc. Amer. Math. Soc. **4** (1998), 74–87 (electronic).
8. Kirill C. H. Mackenzie, *Ehresmann doubles and Drinfel’d doubles for Lie algebroids and Lie bialgebroids*, J. Reine Angew. Math. **658** (2011), 193–245.
9. Yuri I. Manin, *Gauge field theory and complex geometry*, second ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 289, Springer-Verlag, Berlin, 1997, Translated from the 1984 Russian original by N. Koblitz and J. R. King, With an appendix by Sergei Merkulov. MR 1632008 (99e:32001)
10. Rajan Amit Mehta, *Q -algebroids and their cohomology*, J. Symplectic Geom. **7** (2009), no. 3, 263–293. MR 2534186 (2011b:58040)
11. Boris Shoikhet, *On the Duflo formula for L_∞ -algebras and Q -manifolds*, ArXiv Mathematics e-prints (1998), [arXiv:math/9812009](https://arxiv.org/abs/math/9812009).
12. Arkady Yu. Vaintrob, *Lie algebroids and homological vector fields*, Uspekhi Mat. Nauk **52** (1997), no. 2(314), 161–162. MR 1480150

DEPARTMENT OF MATHEMATICS & STATISTICS, SMITH COLLEGE, 44 COLLEGE LANE, NORTHAMPTON, MA 01063

E-mail address: rmehta@smith.edu

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802

E-mail address: stienon@psu.edu

E-mail address: ping@math.psu.edu