

7-16-2010

Poset Pinball, GKM-Compatible Subspaces, and Hessenberg Varieties

Megumi Harada
McMaster University

Julianna Tymoczko
Smith College, jtymoczko@smith.edu

Follow this and additional works at: https://scholarworks.smith.edu/mth_facpubs

Part of the [Mathematics Commons](#)

Recommended Citation

Harada, Megumi and Tymoczko, Julianna, "Poset Pinball, GKM-Compatible Subspaces, and Hessenberg Varieties" (2010).
Mathematics and Statistics: Faculty Publications, Smith College, Northampton, MA.
https://scholarworks.smith.edu/mth_facpubs/33

This Article has been accepted for inclusion in Mathematics and Statistics: Faculty Publications by an authorized administrator of Smith ScholarWorks.
For more information, please contact scholarworks@smith.edu

POSET PINBALL, GKM-COMPATIBLE SUBSPACES, AND HESSENBERG VARIETIES

MEGUMI HARADA AND JULIANNA TYMOCZKO

ABSTRACT. This paper has three main goals. First, we set up a general framework to address the problem of constructing module bases for the equivariant cohomology of certain subspaces of GKM spaces. To this end we introduce the notion of a **GKM-compatible** subspace of an ambient GKM space. We also discuss **poset-upper-triangularity**, a key combinatorial notion in both GKM theory and more generally in localization theory in equivariant cohomology. With a view toward other applications, we present parts of our setup in a general algebraic and combinatorial framework. Second, motivated by our central problem of building module bases, we introduce a combinatorial game which we dub **poset pinball** and illustrate with several examples. Finally, as first applications, we apply the perspective of GKM-compatible subspaces and poset pinball to construct explicit and computationally convenient module bases for the S^1 -equivariant cohomology of all Peterson varieties of classical Lie type, and subregular Springer varieties of Lie type A . In addition, in the Springer case we use our module basis to lift the classical Springer representation on the ordinary cohomology of subregular Springer varieties to S^1 -equivariant cohomology in Lie type A .

CONTENTS

1. Introduction	1
2. Combinatorial and algebraic preliminaries	4
3. Poset pinball: a combinatorial game on directed graphs	8
4. Poset pinball for GKM-compatible subspaces	12
5. Example: Peterson varieties and other regular nilpotent Hessenberg varieties	18
6. Example: Springer varieties in type A	24
References	31

1. INTRODUCTION

This manuscript has three main goals. First, we develop a general framework and perspective to construct computationally convenient module bases for the equivariant cohomology of certain spaces equipped with group actions. In particular, we introduce the notion of a **GKM-compatible subspace** of an ambient GKM space. While not themselves GKM spaces, GKM-compatible subspaces allow us to exploit the combinatorial advantages of GKM theory applied to the ambient GKM space. We primarily use Borel-equivariant cohomology with field coefficients, but expect future applications in other generalized equivariant cohomology theories. For this reason, we present part of this framework in an abstract algebraic setting, formalizing algebraic properties of the equivariant cohomology of GKM spaces in the language of submodules of product modules indexed by a graded partially ordered set. We also discuss the crucial notion of **poset-upper-triangular** subsets of a module, and give one possible answer to a question of Billey's by providing examples of topological spaces with no combinatorially-natural poset-upper-triangular basis.

Second, we introduce a combinatorial game we call **poset pinball**. The game is designed to address some of the difficulties which arise in the analysis of the equivariant cohomology of GKM-compatible subspaces, but the game itself is purely combinatorial and does not depend on the motivating geometry.

Date: May 21, 2018.

2000 Mathematics Subject Classification. Primary: 55N91; Secondary: 22E46, 14L30.

Key words and phrases. equivariant cohomology and localization, Goresky-Kottwitz-MacPherson theory, graded partially ordered sets, nilpotent Hessenberg varieties, Springer theory.

The first author is partially supported by an NSERC Discovery Grant, an NSERC University Faculty Award, and an Ontario Ministry of Research and Innovation Early Researcher Award. The second author is partially supported by NSF grant DMS-0801554, a Sloan Research Fellowship, and an Old Gold Research Fellowship.

Third, as applications, we use the above theory to describe some **nilpotent Hessenberg varieties**, which are a rich class of algebraic varieties arising in geometric representation theory. GKM theory does not directly apply to nilpotent Hessenberg varieties but our methods do. We first prove in Theorem 5.4 that the **nilpotent Hessenberg varieties** $\mathcal{H}(N, H)$ in Lie type A , and in general Lie type given an extra condition on the parameter N , are GKM-compatible subspaces of the flag variety. Then in Theorem 5.9 we use poset pinball to construct explicit module bases for the S^1 -equivariant cohomology rings of **Peterson varieties in all classical Lie types**; this generalizes earlier work in the Lie type A case [16]. Similarly, we use poset pinball in Theorem 6.7 to construct module bases for the S^1 -equivariant cohomology of **subregular Springer varieties in Lie type A** (also studied by Slodowy [27]). We then use this ‘poset pinball basis’ and Kostant-Kumar’s S_n -action on $H_T^*(\mathcal{Flags}(\mathbb{C}^n); \mathbb{C})$ to explicitly construct in Corollary 6.12 a new geometric representation of S_n on $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ and also prove that it lifts the well-known Springer representation on the ordinary cohomology of subregular Springer varieties to S^1 -equivariant cohomology.

Our work develops out of GKM theory, named for the influential manuscript of Goresky-Kottwitz-MacPherson [11]. If X is a suitable G -space, GKM theory gives a combinatorial description of the generalized equivariant cohomology ring $E_G^*(X)$ via restriction to $E_G^*(X^G)$. If the G -space X has isolated fixed points, there often exists a computationally convenient module basis for $E_G^*(X)$; a classical example is the set of (equivariant) Schubert classes $\{\sigma_w\}_{w \in W}$ which form a basis for $H_T^*(G/B)$ where G is a reductive complex algebraic group and B is a Borel subgroup. Current research in equivariant topology (especially Schubert calculus) frequently exploits combinatorial properties of these module bases to obtain topological information, e.g. the product structure of equivariant cohomology rings (see for instance [18, 20, 22, 24, 35]). However the so-called “GKM conditions” which guarantee that GKM theory applies to a G -space X are stringent (see Section 4.1), so the theory is restricted in scope. On the other hand, many topological spaces are subspaces of a G -space X for which the GKM package holds, since for instance we may take $X = \mathbb{P}^n$ with the standard torus action. Definition 4.5 introduces the notion of a **GKM-compatible subspace** Y of the G -space X , equipped with the action of a subgroup $H \subseteq G$. We show that we can use GKM theory on the ambient space X in order to draw conclusions about the H -equivariant topology of Y . For example, when $G = T$ is a torus and $H = S$ is a subtorus, we present two concrete constructions of combinatorial bases for $H_S^*(Y)$ given a suitable basis for $H_T^*(X)$: Proposition 4.14 gives a *poset pinball basis* and Theorem 4.18 gives a *matching basis*.

One of the goals of this manuscript is to formalize some of the features of GKM theory into purely algebraic and combinatorial terms. We believe that the separation of the algebra and combinatorics from the specifics of the geometry serves to clarify some of the issues involved. We start with a poset \mathcal{J} satisfying conditions arising naturally in geometric applications. We then place the GKM description of equivariant cohomology rings in the more general algebraic setting of a submodule M of a product module $\prod_{i \in \mathcal{J}} M_i$ whose factors are indexed by the poset \mathcal{J} . We also discuss one of the core notions of this manuscript, namely **poset-upper-triangularity**, defined precisely in Definition 2.3. Roughly, a subset $\{x_\alpha\}_{\alpha \in A} \subset \prod_{i \in \mathcal{J}} M_i$ is poset-upper-triangular if for each α there exist distinct $i_\alpha \in \mathcal{J}$ such that $x_\alpha(j) = 0$ for all $j \not\geq i_\alpha$ in the poset. In many contexts, geometric classes in equivariant cohomology give rise to poset-upper-triangular subsets, like the equivariant Schubert classes for flag varieties, or more generally cohomology classes obtained from Morse flows with respect to moment maps on a symplectic manifold. We present in Theorem 4.2 more general circumstances under which poset-upper-triangular module bases exist for Borel-equivariant cohomology. In this algebraic formalism, the analogue of a GKM-compatible subspace Y of X is a subset $\mathcal{J} \subseteq \mathcal{J}$ of an ambient poset \mathcal{J} and a homomorphism $\prod_{i \in \mathcal{J}} M_i \rightarrow \prod_{j \in \mathcal{J}} M'_j$ which is zero on the factors $i \in \mathcal{J}$ with $i \notin \mathcal{J}$. Our central problem, recorded in an algebraic context in Question 2.8 and in a geometric context in Question 4.9, is that a poset-upper-triangular subset of $\prod_{i \in \mathcal{J}} M_i$ may not be poset-upper-triangular when restricted to the components indexed by \mathcal{J} . This is precisely the issue which our **poset pinball** and its variations are designed to address. While geometric in inspiration, we emphasize that poset pinball only requires the combinatorial data of a poset $(\mathcal{J}, <)$ and a choice of initial subset $\mathcal{J} \subseteq \mathcal{J}$.

As a consequence of the examples of poset pinball games computed in this manuscript, we also propose a perspective on poset-upper-triangularity which somewhat differs from that which may be most natural from the point of view of combinatorics (see Remark 4.16). Combinatorists view poset-upper-triangularity as a key computational property; indeed, Billey suggests that it is one of the essential features of the Schubert basis and asks for constructions of such poset-upper-triangular bases in equivariant cohomology rings of other G -spaces. On the other hand, in our poset pinball examples it can happen that we obtain subsets

of modules that are *not* poset-upper-triangular with respect to the original partial order $<$ but are nevertheless poset-upper-triangular with respect to a total order \prec compatible with the original partial order. Poset-upper-triangularity with respect to a total order often suffices to guarantee that a subset is linearly independent and hence a module basis, so in some geometric contexts, it may be more natural to require only that module bases be upper-triangular with respect to some choice of total order compatible with the original partial order.

We now present a concrete example of poset pinball in order to convey the flavor of the game. Let $\mathcal{J} = S_4$ denote the permutation group S_4 . Elements of $\mathcal{J} = S_4$ are the vertices in Figure 1.1 and are labelled by a choice of reduced-word decomposition. (We omit some of the elements of S_4 in the figure because they are not relevant in this example.) The set \mathcal{J} is partially ordered by Bruhat order, so we draw an edge between vertices $w, w' \in \mathcal{J}$ if and only if $w < w'$ and there is no $w'' \in \mathcal{J}$ with $w < w'' < w'$. The vertices are drawn so that the poset's minimal element e is at the bottom, and horizontal levels indicate Bruhat length. Let \mathcal{J} be the subset of \mathcal{J} indicated by the circled vertices in Figure 1.1, so there are 6 elements in the subset \mathcal{J} . To play poset pinball, we successively release a circled vertex, starting from the lowest vertex in \mathcal{J} and then moving up; we imagine each circled vertex rolling down along the edges of the poset until it comes to rest in a lowest-possible unoccupied vertex, at which point the circle turns into a square. (See Section 3.1 for precise statements.) This results in a choice of 6 vertices of \mathcal{J} corresponding to the original elements of \mathcal{J} and indicated by the squared vertices in the figure below. Table (1.1) records the exact correspondence between the initial vertices $w \in \mathcal{J}$ and the squared vertices $v \in \mathcal{J}$.




FIGURE 1.1. An instance of poset pinball, for the Springer variety in $\mathcal{F}lags(\mathbb{C}^4)$ specified by a nilpotent operator N corresponding to the partition $(2, 2)$.

pinball step	w_k	v_k
1	$w_1 = e$	$v_1 = e$
2	$w_2 = s_2$	$v_2 = s_2$
3	$w_3 = s_2s_3$	$v_3 = s_3$
4	$w_4 = s_2s_1$	$v_4 = s_1$
5	$w_5 = s_2s_1s_3$	$v_5 = s_1s_3$
6	$w_6 = s_2s_1s_3s_2$	$v_6 = s_1s_2$

In fact, let $\mathcal{S}_N \subseteq \mathcal{F}lags(\mathbb{C}^4) \cong GL(4, \mathbb{C})/B$ denote the Springer variety of type A corresponding to a nilpotent endomorphism N with two Jordan blocks of size 2. Let the maximal torus T act in the standard way on $\mathcal{F}lags(\mathbb{C}^4)$. An S^1 subtorus of T preserves \mathcal{S}_N and it turns out that the S^1 -fixed points $\mathcal{S}_N^{S^1} = \mathcal{J}$ are exactly the subset of $\mathcal{J} = S_4 \cong \mathcal{F}lags(\mathbb{C}^4)^T$ indicated in Figure 1.1. (See Section 6.) Moreover in this case the choices $\{v_i\}_{i=1}^6$ obtained via poset pinball give rise to an $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis for $H_{S^1}^*(\mathcal{S}_N; \mathbb{F})$. Specifically, consider the ring homomorphism

$$(1.2) \quad H_T^*(\mathcal{F}lags(\mathbb{C}^4); \mathbb{F}) \rightarrow H_{S^1}^*(\mathcal{S}_N; \mathbb{F})$$

obtained by composing the map $H_T^*(\mathcal{F}lags(\mathbb{C}^4); \mathbb{F}) \rightarrow H_{S^1}^*(\mathcal{F}lags(\mathbb{C}^4); \mathbb{F})$ with the map $H_{S^1}^*(\mathcal{F}lags(\mathbb{C}^4); \mathbb{F}) \rightarrow H_{S^1}^*(\mathcal{S}_N; \mathbb{F})$ induced by inclusion of groups $S^1 \hookrightarrow T$ and spaces $\mathcal{S}_N \hookrightarrow \mathcal{F}lags(\mathbb{C}^4)$ respectively. Denote the image of an equivariant Schubert class $\sigma_w \in H_T^*(\mathcal{F}lags(\mathbb{C}^4); \mathbb{F})$ under the map (1.2) by p_w . Then the 6 classes p_{v_1}, \dots, p_{v_6} which correspond to the outcome of the previous poset pinball game form an $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis for $H_{S^1}^*(\mathcal{S}_N; \mathbb{F})$. Moreover, the set of images of the p_{v_i} under the natural restriction map

$$(1.3) \quad \iota^* : H_{S^1}^*(\mathcal{S}_N; \mathbb{F}) \hookrightarrow H_{S^1}^*(\mathcal{S}_N^{S^1}; \mathbb{F}) \cong \bigoplus_{i=1}^6 H_{S^1}^*(\text{pt}; \mathbb{F}) \cong \bigoplus_{i=1}^6 \mathbb{F}[t]$$

is poset-upper-triangular with respect to the partial order on the fixed points $\mathcal{S}_N^{S^1}$ induced from Bruhat order. In other words, for each $p \in H_{S^1}^*(\mathcal{S}_N; \mathbb{F})$ let $p(w_i)$ denote the component of ι^*p in the w_i -th summand of the right side of Equation (1.3). Then for each $i = 1, 2, \dots, 6$ we have

$$(1.4) \quad p_{v_i}(w_j) = 0 \text{ for } w_j \not\geq w_i$$

where the inequality indicates the partial order on $\mathcal{S}_N^{S^1} \subseteq S_4$ induced from Bruhat order on S_4 . These vanishing properties allow us to do explicit computations in $H_{S^1}^*(\mathcal{S}_N; \mathbb{F})$ (see Section 6 for applications).

We now briefly outline the contents of the paper. In Section 2 we present the combinatorial and algebraic preliminaries for the pinball game. Poset pinball itself is described in detail in Section 3. We give two concrete examples of poset pinball in Section 3.2 and make initial observations concerning the role played by principal order ideals in pinball theory in Section 3.3. We then explain the geometric motivation and context in Section 4. We begin with a brief review of relevant GKM theory in Section 4.1, and then in Section 4.2 define GKM-compatible subspaces of GKM spaces. With a view toward future work, we keep the discussion in Sections 2, 4.1, and 4.2 as general as possible. Section 4.3 discusses the case of Borel-equivariant cohomology, which is the main focus of this manuscript. The construction of poset pinball bases for the S^1 -equivariant cohomology of Peterson varieties in classical Lie type occupies Section 5. In Section 6, we construct a pinball basis for the equivariant cohomology $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ of the subregular Springer variety of type A , and lift the usual Springer action on $H^*(\mathcal{S}_N; \mathbb{C})$ to $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$.

Open questions and avenues for future work are mentioned throughout the manuscript.

Acknowledgements. We thank David Anderson, Darius Bayegan, Barry Dewitt, Rebecca Goldin, and Bridget Tenner for helpful conversations. Both authors were supported in part by the NSF-funded Midwest Topology Network's travel research grant. Moreover, some of this work was conducted while the first author was a Research Member at the Mathematical Sciences Research Institute as part of the Symplectic and Contact Geometry and Topology program in spring 2010. Both authors also benefited from the American Institute of Mathematics workshop "Localization techniques in equivariant cohomology" held in March 2010. We gratefully acknowledge the support and generosity of the Midwest Topology Network, MSRI, and AIM.

2. COMBINATORIAL AND ALGEBRAIC PRELIMINARIES

The main goal of this section is to present, in purely algebraic and combinatorial terms, the problem that is the primary focus of this manuscript. The setup is as follows. Let R be a commutative ring and let $(\mathcal{J}, <)$ be a partially ordered set. Let M be an R -module that can be realized as a submodule of a product $\prod_{i \in \mathcal{J}} M_i$ of R -modules M_i for each $i \in \mathcal{J}$. Our motivating geometric examples arise from equivariant topology, where the ring $R = E_G^*(\text{pt})$ is the equivariant cohomology ring of a point, the module $M = E_G^*(X)$ is that of a G -space X , and M injects as a submodule into the product $\prod_{i \in \mathcal{J}} M_i = \prod_{i \in \mathcal{J}} E_G^*(\text{pt})$. (See Section 4.1 for geometric details.) However, in order to highlight the algebraic issues, we have chosen to keep the geometry out of the discussion until Section 4.

With this in mind, our main problem can be stated as follows: given two modules $M \subseteq \prod_{i \in \mathcal{J}} M_i$ and $M' \subseteq \prod_{j \in \mathcal{J}} M'_j$ for $\mathcal{J} \subseteq \mathcal{J}$, together with a homomorphism $M \rightarrow M'$ and a *poset-upper-triangular basis* as defined below, we wish to construct computationally convenient module bases for M' . The next two sections provide the necessary background for this discussion. Section 2.1 establishes notation and terminology for posets and product modules indexed by posets. In particular, we give a precise definition of a *poset-upper-triangular* subset of such a module. We make general observations related to the construction of poset-upper-triangular module bases in Section 2.2. Section 2.3 describes our main problem precisely.

2.1. Combinatorial preliminaries. Let $(\mathcal{J}, <)$ be a partially ordered set. Recall that for $i, j \in \mathcal{J}$, we say that i **covers** j if $j < i$ and, in addition, there is no $i' \in \mathcal{J}$ with $j < i' < i$. A **rank function** $\rho : \mathcal{J} \rightarrow \mathbb{N}$ is an \mathbb{N} -valued function on the poset such that if i covers j then $\rho(i) = \rho(j) + 1$. In the case where \mathcal{J} is infinite, we also require $\rho(i) > \rho(j)$. For $i \in \mathcal{J}$, we call $\rho(i)$ the **rank** of i . A poset $(\mathcal{J}, <, \rho)$ equipped with a rank function is called a **graded poset**.

The partially ordered sets \mathcal{J} in this manuscript will always be assumed to satisfy the following conditions:

- \mathcal{J} is countable,
- \mathcal{J} is graded, and
- for any $d \in \mathbb{N}$ the set $\{i \in \mathcal{J} : \rho(i) \leq d\}$ is finite.

We also recall the following [30, Chapter 3].

Definition 2.1. Given a poset $(\mathcal{J}, <)$ and an element $i \in \mathcal{J}$, the **principal order ideal** $\mathcal{L}_{\mathcal{J}}(i)$ of i is the subset of elements $i' \in \mathcal{J}$ less than or equal to i with respect to $<$. In other words

$$\mathcal{L}_{\mathcal{J}}(i) := \{i' \in \mathcal{J} \mid i' \leq i\}.$$

Similarly, the **principal order filter** $\mathcal{U}_{\mathcal{J}}(i)$ of i is the subset of elements $i' \in \mathcal{J}$ greater than or equal to i with respect to $<$. In other words

$$\mathcal{U}_{\mathcal{J}}(i) := \{i' \in \mathcal{J} \mid i \leq i'\}.$$

The posets appearing in this manuscript arise as indexing sets for products of modules, so we introduce some terminology for this situation. Let R be a commutative ring. Suppose \mathcal{J} is a poset as above and M_i an R -module for each $i \in \mathcal{J}$. Suppose M is a submodule of the product module $\prod_{i \in \mathcal{J}} M_i$. For $x \in M \subseteq \prod_{i \in \mathcal{J}} M_i$ we denote by $x(i) \in M_i$ the component of x in the i -th factor of the direct product. For $x \in M$, let

$$\text{supp}(x) := \{i \in \mathcal{J} : x(i) \neq 0 \in M_i\}$$

denote the **support** of x , i.e. the components $i \in \mathcal{J}$ on which x does not vanish.

Definition 2.2. Suppose $M \subseteq \prod_{i \in \mathcal{J}} M_i$ is a module as above. An element $x \in M$ is a **poset-flow-up (with respect to $<$)** if the support of x contains a minimal element i , the principal order filter of which contains $\text{supp}(x)$; in other words $i \in \text{supp}(x) \subseteq \mathcal{U}_{\mathcal{J}}(i)$. We denote the (unique) element i by $\min(x)$ and call it the **minimum nonzero coordinate of (the poset-flow-up) x** .

Note that if $x \in M$ is a poset-flow-up, then

$$(2.1) \quad x(j) = 0 \text{ for all } j \not\geq \min(x).$$

Throughout the manuscript, we consider collections of elements in a module $M \subseteq \prod_{i \in \mathcal{J}} M_i$ with vanishing properties similar to those in (2.1). We have the following.

Definition 2.3. Suppose $M \subseteq \prod_{i \in \mathcal{J}} M_i$ is a module as above. Suppose $\mathcal{B} = \{x_\alpha\} \subseteq M$. The set \mathcal{B} is **poset-upper-triangular** if

- each x_α is a poset-flow-up, and
- for $\alpha \neq \beta$, we have $\min(x_\alpha) \neq \min(x_\beta)$.

2.2. Upper-triangularity for module generators and bases. In this section, we construct poset-upper-triangular module generators and bases of R -modules $M \subseteq \prod_{i \in \mathcal{J}} M_i$ from a purely algebraic viewpoint. An R -module basis must both generate the module and be R -linearly independent; we address the two conditions separately. We keep the assumptions of Section 2.1. Throughout, we think of M as a topological R -module, considered as the (inverse) limit of the submodules $M \cap \prod_{j \leq i} M_j$. In particular, the terms ‘generator’ and ‘basis’ are understood in the topological sense. (If the poset \mathcal{J} is finite, then this agrees with the usual notions.)

Suppose \prec is a total ordering on \mathcal{J} compatible with the given partial order $<$ on \mathcal{J} . We begin by inductively constructing a set of generators of $M \subseteq \prod_{i \in \mathcal{J}} M_i$ that consists of poset-flow-ups with respect to \prec . Poset-upper-triangularity with respect to the total order \prec is a weaker condition than that with respect to the original partial order $<$ but we will see later in the manuscript that upper-triangularity with respect to \prec suffices for many computational purposes.

Proposition 2.4. *Let $(\mathcal{J}, <)$ be a countable graded partially ordered set with a finite number of elements of each rank. Let R be a commutative ring, M_i an R -module for each $i \in \mathcal{J}$, and M an R -submodule $M \subseteq \prod_{i \in \mathcal{J}} M_i$. Suppose \prec is a total ordering compatible with the partial order $<$ on \mathcal{J} . For each $i \in \mathcal{J}$, define*

$$(2.2) \quad \mathcal{V}^i := \{x \in M \mid x(j) = 0 \text{ for all } j \in \mathcal{J} \text{ with } j \prec i\}$$

and denote by $\mathcal{V}^i(i)$ the image of \mathcal{V}^i in M_i under the natural projection $M \rightarrow M_i$. Then for each $i \in \mathcal{J}$ there exist sets \mathcal{K}_i and nonzero elements $\{x_{i,k}\}_{k \in \mathcal{K}_i} \subseteq M$ such that

- (i) $x_{i,k}(j) = 0$ for all $k \in \mathcal{K}_i$ and $j \in \mathcal{J}$ with $j \prec i$,
- (ii) the set $\{x_{i,k}(i)\}_{k \in \mathcal{K}_i} \subseteq M_i$ generates $\mathcal{V}^i(i)$ as an R -module.

Moreover, the union $\mathcal{B} := \bigcup_{i \in \mathcal{J}} \{x_{i,k}\}_{k \in \mathcal{K}_i}$ is a set of R -module generators of M .

Proof. Note first that \mathcal{V}^i and $\mathcal{V}^i(i)$ are R -submodules of M and M_i respectively. For each $i \in \mathcal{J}$ choose a set of nonzero generators $\{y_{i,k}\}_{k \in \mathcal{K}_i}$ of $\mathcal{V}^i(i)$ indexed by a set \mathcal{K}_i . For each $i \in \mathcal{J}$ and $k \in \mathcal{K}_i$ choose an element $x_{i,k} \in \mathcal{V}^i$ that projects to $y_{i,k}$ under the natural map, namely $x_{i,k}(i) = y_{i,k}$. By construction, the sets \mathcal{K}_i and elements $x_{i,k}$ for $i \in \mathcal{J}, k \in \mathcal{K}_i$ satisfy conditions (i) and (ii) of the proposition.

It remains to show that the union $\mathcal{B} := \bigcup_{i \in \mathcal{J}} \{x_{i,k}\}_{k \in \mathcal{K}_i}$ forms a set of R -module generators for M . We first claim that for any $i \in \mathcal{J}$, the module M is generated as an R -module by the elements $\mathcal{B}_i := \bigcup_{j \prec i} \bigcup_{k \in \mathcal{K}_j} \{x_{j,k}\}$,

together with the submodule \mathcal{V}^i . We proceed by induction. For the base case, let $i_0 \in \mathcal{J}$ be the (unique) minimal element in \mathcal{J} with respect to the total order \prec . In this case $\{j \in \mathcal{J} : j \prec i_0\}$ is empty, so $\mathcal{V}^{i_0} = M$ and the claim holds. Now let $i \in \mathcal{J}$ and suppose by induction that the claim holds for all $j \prec i$. By our assumptions on \mathcal{J} in Section 2.1, the set $\{j \in \mathcal{J} : j \prec i\}$ is finite for any total ordering \prec compatible with the partial order $<$. Hence there exists $i' \in \mathcal{J}$ which is maximal with respect to \prec in $\{j \in \mathcal{J} : j \prec i\}$. Now let $x \in M$. By the inductive hypothesis there exist $x'' \in \mathcal{V}^{i'}$ and $a_{j,k} \in R$ for $j \prec i', k \in \mathcal{K}_j$ such that

$$(2.3) \quad x = x'' + \sum_{j \prec i'} \sum_{k \in \mathcal{K}_j} a_{j,k} x_{j,k}.$$

Since $\{x_{i',k}(i')\}_{k \in \mathcal{K}_{i'}}$ generate $\mathcal{V}^{i'}(i')$ there exist $a_{i',k} \in R$ for $k \in \mathcal{K}_{i'}$ such that

$$(2.4) \quad x''(i') = \sum_{k \in \mathcal{K}_{i'}} a_{i',k} y_{i',k}.$$

Now define

$$(2.5) \quad x' := x'' - \sum_{k \in \mathcal{K}_{i'}} a_{i',k} x_{i',k}.$$

Then by construction

$$(2.6) \quad x = x' + \sum_{j \preceq i'} \sum_{k \in \mathcal{K}_j} a_{j,k} x_{j,k} = x'' + \sum_{j \prec i} \sum_{k \in \mathcal{K}_j} a_{j,k} x_{j,k}.$$

To prove the claim, we show that $x' \in \mathcal{V}^i$. Suppose $j \prec i$. Then either $j \prec i'$ or $j = i'$. First suppose $j \prec i'$. Since $x'' \in \mathcal{V}^{i'}$ we have $x''(j) = 0$. Similarly, each $x_{i',k} \in \mathcal{V}^{i'}$ so $x_{i',k}(j) = 0$. Projection to M_j is R -linear, so

$$x'(j) = x''(j) - \sum_{k \in \mathcal{K}_{i'}} a_{i',k} x_{i',k}(j) = 0$$

as desired. Now suppose $j = i'$. Then by definition of $x_{i',k}$ and by Equation (2.4) we see

$$(2.7) \quad x'(i') = x''(i') - \sum_{k \in \mathcal{K}_{i'}} a_{i',k} x_{i',k}(i') = x''(i') - \sum_{k \in \mathcal{K}_{i'}} a_{i',k} y_{i',k} = 0.$$

Together these mean $x' \in \mathcal{V}^i$. Hence M is generated by the elements in \mathcal{B}_i together with \mathcal{V}^i . The result follows from $M = \varprojlim M \cap \left(\prod_{j \leq i} M_j \right)$. \square

- Remark 2.5.** (1) *It is sometimes impossible to find a set of generators of a module M which is poset-upper-triangular with respect to the original partial order $<$. For example, suppose M is the image of the standard diagonal embedding $R \hookrightarrow \prod_{i \in \mathcal{J}} R$ with $1 < |\mathcal{J}| < \infty$, and take the trivial partial order on \mathcal{J} in which all pairs of elements are incomparable and the trivial rank function $\rho \equiv 0$. Hence the choice of a total order \prec is a crucial step in Proposition 2.4.*
- (2) *On the other hand, Proposition 2.4 extends straightforwardly to partially ordered sets without a choice of total order if we assume that the vanishing submodules \mathcal{V}^i together generate all of M .*

The previous proposition constructed module generators. Our next task is to deal with R -linear independence. For the remainder of the manuscript, we assume that

- R is a domain, and
- for all $i \in \mathcal{J}$ the module M_i is R -torsion-free.

These assumptions imply that the submodule $M \subseteq \prod_{i \in \mathcal{J}} M_i$ is also R -torsion-free. (In fact, in our geometric applications, it is usually the case that $M_i \cong R$ for all i and that M is a free R -module.) We begin by giving conditions under which the construction in Proposition 2.4 in fact yields an R -module basis.

Proposition 2.6. *Let R, \mathcal{J}, M_i, M satisfy the conditions of Proposition 2.4. Assume that R is a domain and that the R -module M_i is R -torsion-free for all $i \in \mathcal{J}$.*

- (1) *If $\{x_\alpha\}$ is poset-upper-triangular, then $\{x_\alpha\}$ is R -linearly independent.*
- (2) *Suppose that for all $i \in \mathcal{J}$, the sets $\{x_{i,k}\}_{k \in \mathcal{K}_i}$ constructed in Proposition 2.4 may be chosen to be R -linearly independent. Then the union $\mathcal{B} = \bigcup_{i \in \mathcal{J}} \{x_{i,k}\}_{k \in \mathcal{K}_i}$ is an R -module basis of M .*
- (3) *Suppose that for all $i \in \mathcal{J}$, the index sets \mathcal{K}_i constructed in Proposition 2.4 may be chosen such that either $|\mathcal{K}_i| = 0$ or $|\mathcal{K}_i| = 1$. Then the union $\mathcal{B} = \bigcup_{i \in \mathcal{J}} \{x_{i,k}\}_{k \in \mathcal{K}_i}$ is an R -module basis of M .*
- (4) *Suppose that R is a principal ideal domain and each M_i is a free R -module of rank 1. Then the index sets \mathcal{K}_i constructed in Proposition 2.4 may be chosen such that either $|\mathcal{K}_i| = 0$ or $|\mathcal{K}_i| = 1$ and the union $\mathcal{B} = \bigcup_{i \in \mathcal{J}} \{x_{i,k}\}_{k \in \mathcal{K}_i}$ is an R -module basis of M .*

Proof. If $\{x_\alpha\}$ is poset-upper-triangular then for each $i \in \mathcal{J}$ there is at most one α with $\min(x_\alpha) = i$. Using this, Part (1) follows by induction. Part (2) is by definition. Part (3) follows from Part (2). Part (4) is a special case of Part (3) since, by definition, an ideal in a PID is generated by a single element. \square

In our geometric applications, we are frequently in the situation of Part (4) of Proposition 2.6. More generally, whenever Part (3) of Proposition 2.6 holds, the module basis elements correspond to elements of the poset, so we may think of the bases \mathcal{B} as being indexed by the poset \mathcal{J} (or possibly a subset of \mathcal{J}). In this way, the combinatorics of the poset \mathcal{J} interacts directly with the algebra of the module M via the basis \mathcal{B} . This links the combinatorial strategies of Section 3 to the algebraic problem of constructing module bases. We close the section with a partial converse to Proposition 2.4.

Lemma 2.7. *Let R, \mathcal{J}, M_i, M satisfy the conditions of Proposition 2.4. Assume that R is a domain and $M_i \cong R$ for each $i \in \mathcal{J}$. Suppose that $\{x_i\}_{i \in \mathcal{J}}$ is poset-upper-triangular with respect to the partial order on \mathcal{J} , that $\min(x_i) = i$ for each i , and that $\{x_i\}_{i \in \mathcal{J}}$ generates M as an R -module. Then the choice of singleton set $\{x_i\}$ for each $i \in \mathcal{J}$ satisfies Conditions (i) and (ii) of Proposition 2.4.*

Proof. The set $\{x_i\}_{i \in \mathcal{J}}$ is poset-upper-triangular with respect to $<$ so it is also poset-upper-triangular with respect to any total ordering \prec compatible with $<$. This is Condition (i) of Proposition 2.4.

To prove Condition (ii), we show that $x_j(j)$ generates $\mathcal{V}^j(j)$ as an R -module for each j . We first claim that each \mathcal{V}^j is generated by $\{x_i\}_{j \prec i \text{ or } j=i}$. We proceed by induction. For the base case, let j_0 be the minimal element of \mathcal{J} with respect to \prec . Then the set $\{i : j_0 \prec i \text{ or } j_0 = i\}$ is all of \mathcal{J} and the claim trivially holds. Now suppose that the claim holds for all i with $i \prec j$ and that i' is maximal in $\{i \prec j\}$. (As in the proof of Proposition 2.4, a maximal i' exists because $\{i : i \prec j\}$ is finite by the hypotheses on \mathcal{J} .) Let $x \in \mathcal{V}^j \subseteq \mathcal{V}^{i'}$. By the inductive hypothesis,

$$x = c_{i'} x_{i'} + \sum_{i: i' \prec i} c_i x_i = c_{i'} x_{i'} + \sum_{i: i=j \text{ or } j \prec i} c_i x_i.$$

The set is poset-upper-triangular so $x_i(i') = 0$ for all i with $i' \prec i$. Evaluating x at i' yields $x(i') = c_{i'} x_{i'}(i')$ which must equal 0 since $x \in \mathcal{V}^j$. By assumption $x_{i'}(i') \neq 0$ so $c_{i'} = 0$. This means \mathcal{V}^j is generated by $\{x_i\}_{j \prec i \text{ or } j=i}$ as desired.

Finally, evaluation at j yields $x(j) = c_j x_j(j)$. The element $x \in \mathcal{V}^j$ was chosen arbitrarily, so we conclude $\mathcal{V}^j(j)$ is generated by the single element $x_j(j)$ as claimed. \square

2.3. Bases for submodules of products. In this section we present the central problem of the manuscript, stated in purely algebraic and combinatorial language. Its geometric manifestation is reserved until Section 4. We also explain the core difficulty in addressing the problem, which motivates the poset pinball game introduced in Section 3.

Let \mathcal{J} be a countable graded poset with each rank finite. Let \mathcal{J} be a subset of \mathcal{J} with the partial order induced from \mathcal{J} . Let R and R' be integral domains, M_i a torsion-free R -module for each $i \in \mathcal{J}$, and M'_j a torsion-free R' -module for each $j \in \mathcal{J}$. Let $M \subseteq \prod_{i \in \mathcal{J}} M_i$ and $M' \subseteq \prod_{i \in \mathcal{J}} M'_i$ be R - and R' -submodules, respectively, of the given products. Suppose $\gamma : R \rightarrow R'$ is a ring homomorphism and that $\phi_i : M_i \rightarrow M'_i$ are surjective additive homomorphisms for each $i \in \mathcal{J}$ satisfying

$$(2.8) \quad \phi(rm) = \gamma(r)\phi(m) \quad \text{for all } r \in R, i \in \mathcal{J}, \text{ and } m \in M.$$

We also assume the homomorphism $\prod_{i \in \mathcal{J}} \phi_i : \prod_{i \in \mathcal{J}} M_i \rightarrow \prod_{i \in \mathcal{J}} M'_i$ restricts to a surjection $\phi : M \rightarrow M'$ so that the diagram

$$(2.9) \quad \begin{array}{ccc} M & \longrightarrow & \prod_{i \in \mathcal{J}} M_i \\ \phi \downarrow & & \downarrow \prod_{i \in \mathcal{J}} \phi_i \\ M' & \longrightarrow & \prod_{i \in \mathcal{J}} M'_i \end{array}$$

commutes, where the right vertical map is understood to be 0 on the components M_i for $i \notin \mathcal{J}$.

Now suppose $\mathcal{B} = \{x_\alpha\}$ is a poset-upper-triangular basis of M . (In many examples, such bases exist because of extra geometric structure; see Section 4.) Poset-upper-triangularity implies that the elements x_α have convenient vanishing properties when viewed in the product $\prod_{i \in \mathcal{J}} M_i$. The homomorphism ϕ is surjective, so there exists a subset of the image $\phi(\mathcal{B}) \subseteq M'$ which generates M' . We may then ask the following question (see Question 4.9 for the geometric version).

Question 2.8. *Is it possible to obtain a poset-upper-triangular basis for $M' \subseteq \prod_{i \in \mathcal{J}} M'_i$ with respect to the induced partial order on $\mathcal{J} \subseteq \mathcal{J}$ from a subset of $\phi(\mathcal{B})$?*

This turns out to be a difficult problem. The fundamental obstacle is that the maps in (2.9) do not necessarily behave well with respect to poset-flow-ups. More precisely, the intersection $\mathcal{L}_{\mathcal{J}}(i) \cap \mathcal{J}$ of a principal order filter $\mathcal{L}_{\mathcal{J}}(i)$ with the subset \mathcal{J} is *not* necessarily a principal order filter of \mathcal{J} when $i \notin \mathcal{J}$. As a consequence, the images $\phi(\mathcal{B}) = \{\phi(x_\alpha)\}$ need not even be poset-flow-up elements in M' in the sense of Definition 2.2. This means that poset-upper-triangularity with respect to $(\mathcal{J}, <)$ does not immediately translate via ϕ to poset-upper-triangularity with respect to $(\mathcal{J}, <)$. In the next section, we introduce the combinatorial game of **poset pinball**, which was created to address these difficulties.

3. POSET PINBALL: A COMBINATORIAL GAME ON DIRECTED GRAPHS

3.1. The poset pinball game. We now introduce a non-deterministic game which we call **poset pinball**, by analogy with pinball arcade games, which involve dropping balls on a tilted board. The game can be understood and played independently of the considerations in the previous section; however, the game was designed to address the algebraic difficulties outlined in Section 2.3 (and discussed from a geometric perspective in Section 4). We present several variants of poset pinball; which flavor one plays depends on the geometric, algebraic, or combinatorial context. Examples are in Section 3.2. In Section 3.3, we discuss principal order ideals and the role they play in poset pinball.

We begin with the basic structure of the game, common to all variants. Henceforth we assume that the poset \mathcal{J} is finite.

Poset pinball rules and terminology:

- (1) Let $(\mathcal{J}, <)$ be a finite partially ordered set. We identify \mathcal{J} with its Hasse diagram, so we think of \mathcal{J} as a directed acyclic graph with vertices the elements of \mathcal{J} and with a directed edge from i to i' precisely when i covers i' with respect to the partial order. We denote an edge from i to i' by $i \mapsto i'$.

- (2) The graph \mathcal{J} is the **pinball board**, or simply the **board**. The vertices are called **pinball slots**, or simply **slots**. At most one pinball can occupy a slot at any time.
- (3) Let \mathcal{J} be a fixed subset of \mathcal{J} . We call \mathcal{J} an **initial subset**. Note that \mathcal{J} inherits a partial order from \mathcal{J} .
- (4) We place **pinballs** at the initial subset, i.e., for each element $j \in \mathcal{J}$, we place a pinball at the slot corresponding to j .
- (5) The directed edges of the graph \mathcal{J} are called **pinball slides**, or simply **slides**. When released, a pinball may **roll down** along a slide, in the direction determined by the directed edge. Specifically, if $i \mapsto i'$ is an edge, a pinball at slot i may roll down to the slot i' .
- (6) During the game, we occasionally place **walls** across some slides. A wall across a slide *prevents* a pinball from rolling down that edge (slide). The initial board has no walls. A wall is never removed once it has been placed.
- (7) Fix a *total order* \prec on the initial subset \mathcal{J} subordinate to the induced partial order on \mathcal{J} . We write $\mathcal{J} = \{j_1 \prec j_2 \prec \dots \prec j_{|\mathcal{J}|}\}$ with respect to this total order.
- (8) We now define the procedure for allowing a pinball to **roll down (along slides)**. Suppose a pinball is at slot $i \in \mathcal{J}$. Consider the set of downward-pointing edges with i as top vertex. The pinball at slot i is allowed to roll down to i' as long as there is no wall across the slide $i \mapsto i'$. Hence we consider

$$(3.1) \quad \left\{ \begin{array}{l} i' \in \mathcal{J} : \quad \text{there exists an edge } i \mapsto i', \text{ and} \\ \quad \quad \quad \text{there is no wall across } i \mapsto i' \end{array} \right\}.$$

Choose an arbitrary element i' in the set in (3.1) and move the pinball to slot i' . We refer to this as **rolling along the slide** $i \mapsto i'$. Repeat the above process using the new slot i' in the role of i above, and continue in this manner. We say that the pinball can **roll no further** if at any stage the set in (3.1) is empty, namely there are no lower available slots. When a pinball starting at slot i has rolled down successive slides until it can roll no further, the final slot at which the pinball rests is called the **rolldown of i** and denoted $roll(i)$. We refer to this process of associating to i its rolldown $roll(i)$ as **rolling (or dropping) the pinball**. This procedure is *not* deterministic because of the choices made when rolling along each slide (just like real-life pinball!). Note also that the rolldown $roll(j)$ of a pinball which was originally at a slot $j \in \mathcal{J}$ might *not* be an element of \mathcal{J} .

- (9) We drop pinballs successively according to the total order \prec on \mathcal{J} . Hence we first drop the pinball from slot j_1 as described above. For every $k = 1, 2, \dots, |\mathcal{J}|$, after rolling the k -th pinball, we may place more walls along the slides of the board. Each version of pinball has a separate set of rules for placing walls; details for each variant are given in the description below. Once the first $k - 1$ pinballs are dropped, we drop the pinball at j_k and continue until all $|\mathcal{J}|$ pinballs are dropped.
- (10) Fix the board \mathcal{J} , the initial subset \mathcal{J} , the choice of total order \prec , and a particular outcome of poset pinball, which we write as $\{(j, roll(j)) : j \in \mathcal{J}\}$. Then we denote by $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ the set of slots in \mathcal{J} occupied by the rolldown elements, i.e.

$$\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec) := \{roll(j) : j \in \mathcal{J}\} \subseteq \mathcal{J}.$$

We call $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ the **rolldown set** for the given outcome of pinball. We also denote by $\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec)$ the set of rolldown elements for the first k pinballs, i.e.

$$\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec) := \{roll(j_\ell) : j_\ell \in \mathcal{J}, 1 \leq \ell \leq k\} = \{roll(j_1), roll(j_2), \dots, roll(j_k)\} \subseteq \mathcal{J}.$$

We refer to $\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec)$ as the **rolldown set up to step k** . We emphasize that since pinball is *not* deterministic, the sets $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ and $\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec)$ might *not* be uniquely determined by \mathcal{J}, \mathcal{J} and \prec .

The different versions of poset pinball are distinguished by how the walls are placed after each pinball in the initial subset rolls down. We now describe these variants of poset pinball.

Basic pinball. In this version, the walls are placed as follows. Let $k = 1, 2, \dots, |\mathcal{J}|$. Suppose the k -th pinball j_k has been dropped. We then place a wall across every edge of the form $i \mapsto roll(j_k)$. Hence the walls in basic pinball simply enforce the rule that at most one pinball may occupy a given slot at any time.

Upper-triangular pinball. This version takes into account the partial order structure on \mathcal{J} . Let $k = 1, 2, \dots, |\mathcal{J}|$. Suppose the k -th pinball j_k has been dropped. We then place a wall across:

- every edge of the form $i \mapsto roll(j_k)$ for $i \in \mathcal{J}$, and
- every edge of the form $i \mapsto i'$ for $i \in \mathcal{J}, i' \in \mathcal{L}_{j_k}$.

The rules of upper-triangular pinball ensure that for each $j \in \mathcal{J}$, the element j has a unique maximal rolldown in its principal order ideal, and that maximal rolldown is $\text{roll}(j)$.

Betti pinball. This version of pinball is motivated by the geometric applications discussed later, in which we allow the Betti numbers of an underlying topological space to impose additional constraints on the pinball game. Here we assume that \mathcal{J} is a finite graded poset with rank function $\rho : \mathcal{J} \rightarrow \mathbb{N}$. We also assume $\mathbf{b} = (b_0, b_1, \dots, b_n)$ is a sequence of nonnegative integers. (In geometric applications, these b_j are in fact the Betti numbers of a topological space, so we refer to \mathbf{b} as the *target Betti numbers*.) Let $k = 1, 2, \dots, |\mathcal{J}|$. Suppose the k -th pinball j_k has been dropped. Then the walls are placed as follows:

- Place a wall across any edge of the form $i \mapsto \text{roll}(j_k)$ for $i \in \mathcal{J}$.
- Let $\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec)$ denote the rolldown set up to step k . Let $j \in \{0, 1, \dots, n\}$ and suppose that

$$(3.2) \quad b_j = |\{i \in \mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec) : \rho(i) = j\}|,$$

i.e. there are exactly b_j rolldown elements of rank j at step k . Then place walls across every edge of the form $i \mapsto i'$ where $\text{deg}(i') = j$.

These rules ensure that the number of rolldown elements of rank j do not exceed the given target b_j .

We say that a game of Betti pinball is **successful** if, after all the pinballs in the initial subset \mathcal{J} are dropped, there are precisely b_j rolldowns of rank j for each j . In other words, after a successful game of Betti pinball, the ranks of the elements of the rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ precisely reflect the target Betti numbers \mathbf{b} .

Upper-triangular Betti pinball. This version adds both the walls for upper-triangular pinball and those for Betti pinball at each pinball step. We leave it to the reader to write the rules. As in Betti pinball, we assume we are given a graded poset \mathcal{J} and the data of *target Betti numbers* $\mathbf{b} = (b_0, b_1, b_2, \dots, b_n)$. Also as in Betti pinball, we say that a game of upper-triangular Betti pinball is **successful** if the ranks of the rolldowns precisely reflect the target Betti numbers.

3.2. Playing pinball: examples. Here we illustrate our poset pinball game with two concrete examples. In both cases, the ambient graded poset \mathcal{J} is the symmetric group S_4 equipped with the usual Bruhat order. We take the rank of a permutation w to be the standard **Bruhat length** of w . For simplicity, we do not draw the entire graph of \mathcal{J} in the figures below, but only those vertices and edges relevant in the game. We label each vertex by its corresponding permutation, factored into simple transpositions. The graph is drawn so that all directed edges point towards the bottom of the page, so the minimal permutation e is at the bottom of the figure. The initial subset \mathcal{J} is indicated by the circled vertices, and the final rolldown set is indicated by the vertices with squares around them. Each example is accompanied by a table recording each step of the poset pinball game as it was played.

Example 3.1. In our first example, the rolldown set is in fact unique for basic, upper-triangular, or Betti pinball with $\mathbf{b} = (1, 3)$. In particular, the partial order induced on \mathcal{J} by \mathcal{J} is already a total order, so there is a unique total order with respect to which to play pinball. Notice that the rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ in this example is a union of principal order ideals; we explore this phenomenon further in Section 3.3.

The initial subset is $\mathcal{J} = \{e, s_3, s_3s_2, s_3s_2s_1\}$. The final drop-down set is $\mathcal{R}(\mathcal{J}, \mathcal{J}) = \{e, s_3, s_2, s_1\}$. (The reader may wish to explore how the game changes if the Betti numbers are $\mathbf{b} = (1, 2, 1)$.)




FIGURE 3.1. Example of basic pinball.

(3.3)

<i>pinball step</i>	w_k	v_k
1	$w_1 = e$	$v_1 = e$
2	$w_2 = s_3$	$v_2 = s_3$
3	$w_3 = s_3 s_2$	$v_3 = s_2$
4	$w_4 = s_3 s_2 s_1$	$v_4 = s_1$

Example 3.2. In this example, we play Betti pinball with target Betti numbers $\mathbf{b} = (1, 3, 4, 3, 1)$. The final drop-down set $\mathcal{R}(\mathcal{J}, \mathcal{J})$ is indicated by the squared vertices. Dotted lines indicate a path in the partial order, with some intermediate vertices omitted for visual simplicity. Also for visual simplicity, not all edges in the poset are drawn above rank 2. In this example, the drop-down set $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ is not a union of principal order ideals because of the constraints imposed by the target Betti numbers.

The rolldown set in Figure 3.2 could also be the outcome of a game of upper-triangular Betti pinball. However, if we instead chose the rolldown roll $(s_1 s_2 s_3 s_1 s_2 s_1) = s_1 s_2 s_3 s_2$ then we would obtain a successful outcome of Betti pinball that is not a successful outcome of upper-triangular Betti pinball.




FIGURE 3.2. An example of Betti pinball.

(3.4)

<i>pinball step</i>	w_k	v_k
1	$w_1 = e = [1, 2, 3, 4]$	$v_1 = e = [1, 2, 3, 4]$
2	$w_2 = s_3 = [1, 2, 4, 3]$	$v_2 = s_3 = [1, 2, 4, 3]$
3	$w_3 = s_2 = [1, 3, 2, 4]$	$v_3 = s_2 = [1, 3, 2, 4]$
4	$w_4 = s_1 = [2, 1, 3, 4]$	$v_4 = s_1 = [2, 1, 3, 4]$
5	$w_5 = s_1 s_3 = s_3 s_1 = [2, 1, 4, 3]$	$v_5 = s_1 s_3 = s_3 s_1 = [2, 1, 4, 3]$
6	$w_6 = s_1 s_2 = [2, 3, 1, 4]$	$v_6 = s_1 s_2 = [2, 3, 1, 4]$
7	$w_7 = s_2 s_1 = [3, 1, 2, 4]$	$v_7 = s_2 s_1 = [3, 1, 2, 4]$
8	$w_8 = s_3 s_2 s_3 = [1, 4, 3, 2]$	$v_8 = s_2 s_3 = [1, 3, 4, 2]$
9	$w_9 = s_2 s_1 s_2 = [3, 2, 1, 4]$	$v_9 = s_2 s_1 s_2 = [3, 2, 1, 4]$
10	$w_{10} = s_3 s_2 s_1 s_3 = [4, 1, 3, 2]$	$v_{10} = s_3 s_2 s_1 = [4, 1, 2, 3]$
11	$w_{11} = s_1 s_2 s_3 s_1 s_2 = [3, 4, 2, 1]$	$v_{11} = s_1 s_2 s_3 = [2, 3, 4, 1]$
12	$w_{12} = s_1 s_2 s_3 s_1 s_2 s_1 = [4, 3, 2, 1]$	$v_{12} = s_1 s_3 s_2 s_1 = [4, 2, 1, 3]$

3.3. Principal order ideals and poset pinball. In this section, we briefly explore the role played by principal order ideals in poset pinball. We are motivated by our geometric applications, in which principal order ideals can correspond naturally to subvarieties in an ambient variety (cf. Remark 3.5).

We begin with a simple statement about basic pinball.

Proposition 3.3. *Let $(\mathcal{J}, <)$ be a finite poset and let $\mathcal{J} \subseteq \mathcal{J}$ be a subset. Let \prec be a total ordering on \mathcal{J} compatible with the partial order $<$ induced from \mathcal{J} . Suppose $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec) \subseteq \mathcal{J}$ is the rolldown set from a game of basic pinball played with board \mathcal{J} , initial set \mathcal{J} , and total order \prec . Then $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ is a union of principal order ideals of \mathcal{J} .*

Proof. A subset $\mathcal{K} \subseteq \mathcal{J}$ is a union of principal order ideals exactly if for all $i \in \mathcal{K}$, the principal order ideal \mathcal{L}_i is entirely contained in \mathcal{K} . Let $\mathcal{J} = \{j_1, j_2, \dots, j_{|\mathcal{J}|}\}$ be the totally-ordered initial subset. We will induct on $k = 1, 2, \dots, |\mathcal{J}|$ to show that each $\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec)$ is a union of principal order ideals. In the base case, the rolldown $roll(j_1)$ associated to j_1 must be minimal in \mathcal{J} , by definition of basic pinball. A minimal element $\{roll(j_1)\}$ is a principal order ideal, so the claim holds for $k = 1$. Now assume that $\mathcal{R}_{k-1}(\mathcal{J}, \mathcal{J}, \prec)$ is a union of principal order ideals. Let $roll(j_k) \leq j_k$ be a rolldown of j_k . If $i < roll(j_k)$ then i must be in the rolldown set $\mathcal{R}_{k-1}(\mathcal{J}, \mathcal{J}, \prec)$ by the rules of basic pinball. Hence \mathcal{L}_i is contained in

$$\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec) = \mathcal{R}_{k-1}(\mathcal{J}, \mathcal{J}, \prec) \cup \{roll(j_k)\}.$$

The rolldown set $\mathcal{R}_{k-1}(\mathcal{J}, \mathcal{J}, \prec)$ is itself a union of principal order ideals by the inductive hypothesis. Hence $\mathcal{R}_k(\mathcal{J}, \mathcal{J}, \prec)$ is also a union of principal order ideals. The case $k = |\mathcal{J}|$ proves the proposition. \square

The previous proposition only applies to basic pinball. In upper-triangular or Betti pinball, the additional walls placed during the game imply that the resulting rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ may not be a union of principal order ideals. Indeed, Example 3.2 is an instance of Betti pinball in which the associated rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ is not a union of principal order ideals. On the other hand, in many examples (like the example in the Introduction or Example 6.14), Betti pinball does produce rolldown sets which are unions of principal order ideals. This leads us to ask the following.

Question 3.4. *What are combinatorial conditions on $\mathcal{J}, \mathcal{J}, \prec$, and (in the case of Betti pinball) the target Betti numbers (b_0, \dots, b_n) which guarantee that outcomes of upper-triangular or Betti pinball are unions of principal order ideals?*

A concrete answer to this question would yield new perspectives on the geometric problems, such as computing cohomology rings or Betti numbers, that motivate our pinball game.

Remark 3.5. *One situation in which an answer to Question 3.4 is straightforward is when the initial subset \mathcal{J} is itself the principal order ideal \mathcal{L}_i of some element $i \in \mathcal{J}$. In this case, a successful outcome of any version of pinball is the rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{J}, \prec)$ that equals the original $\mathcal{J} = \mathcal{L}_i$. This situation can arise naturally in geometric contexts. We give two examples.*

- (1) *The poset \mathcal{J} is the Weyl group W of a complex reductive algebraic group G , identified with the T -fixed points of its flag variety G/B . In this case, a principal order ideal \mathcal{L}_w of $w \in W$ corresponds naturally to the T -fixed points in a Schubert subvariety of G/B .*
- (2) *Let X be a complex projective algebraic variety (possibly singular) equipped with a T -action that has isolated T -fixed points X^T . Choose a one-parameter subgroup $S : \mathbb{G}_m \rightarrow T$ with $X^S = X^T$. Then X is partitioned into locally closed subsets X_p defined by*

$$X_p := \{x \in X : \lim_{z \rightarrow 0} S(z) \cdot x = p\}.$$

The disjoint union $\bigsqcup X_p$ is called a Bialynicki-Birula decomposition of X [2]. Moreover Knutson states that X^T can be given a poset structure by taking the transitive closure of the rule that $p \leq q$ when $p \in \overline{X^q}$ [19]. The principal order ideal \mathcal{L}_p of $p \in X^T$ then corresponds naturally to the T -fixed points in $\overline{X_p}$.

4. POSET PINBALL FOR GKM-COMPATIBLE SUBSPACES

4.1. Background: GKM theory in equivariant cohomology. The algebraic questions discussed in the previous sections arise naturally in equivariant algebraic topology. Suppose G is a topological group and X is a topological space with a continuous G -action. Let pt denote the topological space consisting of one point, equipped with the trivial G -action, and let E_G^* denote a generalized equivariant cohomology theory with a commutative cup product. (Examples include Borel-equivariant cohomology $H_G^*(-; \mathfrak{r})$ for various coefficient rings \mathfrak{r} , topological equivariant K -theory in the sense of Atiyah and Segal, and equivariant cobordism;

cf. [25, Chapter XIII].) Then $E_G^*(\text{pt})$ is a commutative ring, and $E_G^*(X)$ is naturally an $E_G^*(\text{pt})$ -module for any G -space X via the map induced on cohomology by the G -equivariant map $X \rightarrow \text{pt}$.

We work in a situation in which $E_G^*(X)$ has a well-studied combinatorial description, often called ‘‘GKM theory’’ due to an influential manuscript of Goresky-Kottwitz-MacPherson [11]. We present one of many variations and generalizations of GKM theory in the literature (see [14] and references therein).

For the purposes of this manuscript only, we say that **the GKM package** holds for a G -space X when the following statements hold. Let G and X be as above.

- We assume that the G -fixed set X^G consists of countably many isolated points, i.e. $X^G \cong \bigcup_{i \in \mathcal{J}} F_i$ with $F_i \cong \text{pt}$ for all $i \in \mathcal{J}$.
- We assume that the indexing set \mathcal{J} for the fixed points $\bigcup_{i \in \mathcal{J}} F_i$ may be equipped with a graded partial order such that there are only finitely many elements of each rank.
- We assume that X is a stratified G -space $X = \bigcup_{i \in \mathcal{J}} X_i$ with $F_i \in X_i$ for each $i \in \mathcal{J}$ and that the cohomology $E_G^*(X) = \varprojlim E_G^*(X_i)$.
- We assume the restriction map

$$(4.1) \quad \iota^* : E_G^*(X) \rightarrow \prod_{i \in \mathcal{J}} E_G^*(F_i) \cong \prod_{i \in \mathcal{J}} E_G^*(\text{pt})$$

is injective.

- We assume there exist certain nonzero equivariant cohomology classes $e_{ij} \in E_G^*(\text{pt})$ satisfying:
 - if $e_{ij} \neq 1$ then i and j are comparable in \mathcal{J} ,
 - if there is a covering relation between $i > j$ in \mathcal{J} then $e_{ij} \neq 1$
 such that the image of ι^* in (4.1) is precisely

$$(4.2) \quad \text{image}(\iota^*) = \left\{ x \in \prod_{i \in \mathcal{J}} E_G^*(\text{pt}) \mid e_{ij} | (x(i) - x(j)) \text{ for all } j < i \text{ in the partial order.} \right\}$$

- We assume there exists a module basis $\{x_i\}_{i \in \mathcal{J}}$ for $E_G^*(X)$ that is indexed by the (isolated) fixed points F_i , where the $x_i \in E_G^*(X)$ are equivariant cohomology classes satisfying

$$(4.3) \quad x_i(i') = 0 \text{ for } i' \not\geq i,$$

and

$$(4.4) \quad x_i(i) \text{ generates the ideal } e_i E_G^*(\text{pt})$$

where $e_i := \prod_{j < i} e_{ij}$. (In this case $\{x_i\}_{i \in \mathcal{J}}$ is a free $E_G^*(\text{pt})$ -module basis of $E_G^*(X)$ as proven in, e.g., [14, Proposition 4.1].)

The vanishing condition in Equation (4.3) says exactly that the x_i are poset-flow-ups, so the $\{x_i\}_{i \in \mathcal{J}}$ are in fact a poset-upper-triangular module basis by Proposition 2.6.

For convenience, we often collect the information needed to determine $E_G^*(X)$ using Equation (4.2) in a directed, labeled graph called the **GKM graph of X** . The vertices of the GKM graph are the fixed points F_i , or equivalently the elements of the poset \mathcal{J} . There is an edge between F_i and F_j exactly when $e_{ij} \neq 1$. If it exists, the edge between F_i and F_j is labeled e_{ij} and is directed from F_i to F_j exactly when $i > j$. Note that the GKM graph of X contains the Hasse diagram of the poset \mathcal{J} , but possibly includes edges which are not poset covering relations.

We now recall some situations in which the GKM package holds.

Remark 4.1. (1) Let $G = T$ be a compact torus and let $E_G^* = H_T^*(-; \mathbb{F})$ be Borel-equivariant cohomology with coefficients in a field \mathbb{F} of characteristic zero. Let (M, ω, Φ) be a compact Hamiltonian T -manifold with moment map $\Phi : M \rightarrow \mathfrak{t}^*$. Suppose that M has finitely many (isolated) fixed points, and for every codimension 1 subtorus $K \subseteq T$, each connected component of the fixed submanifold M^K has (real) dimension less than or equal to 2. Assume in addition that there are finitely many one-dimensional T -orbits in M and that there exists a T -invariant Palais-Smale metric. Let $\Psi := \Phi^\xi$ denote a generic component of the moment map and consider its negative gradient flow with respect to the given Palais-Smale metric. Denote by $\lambda(p)$ the Morse index of Ψ at a critical point $p \in M^T$. It is known [10, Remark 4.3] that Ψ is index-increasing, i.e. if $p, q \in M^T$ and $\Psi(p) < \Psi(q)$, then $\lambda(p) < \lambda(q)$. We give the fixed point set M^T a partial order by defining $p < q$ precisely if $\Psi(p) < \Psi(q)$ and there exists a one-dimensional T -orbit O whose closure

contains p and q . In particular $p < q$ implies $\lambda(p) < \lambda(q)$. (The Hasse diagram of this poset M^T has an edge between p and q exactly when $p < q$ and $\lambda(q) = \lambda(p) + 2$, i.e. the Morse index increases from p to q by precisely 2.) By standard equivariant symplectic geometry arguments, the Morse index is always even. Thus the function $\rho : M^T \rightarrow \mathbb{Z}_{\geq 0}$ defined by $\rho(p) := \frac{1}{2}\lambda(p)$ is a well-defined rank function, and the fact that Ψ is index-increasing implies that ρ gives M^T the structure of a graded poset.

Moreover, in this situation it is also known that the map (4.1) is injective and that the description of the image of ι^* given in (4.2) is valid [11, Theorem 14.1 (9)]. Finally, in this situation, Goldin and Tolman prove that there exists a collection $\{x_p\}_{p \in M^T}$ of canonical classes [10, Definition 1.1] in $H_T^*(M; \mathbb{F})$ indexed by the T -fixed points M^T . These canonical classes give a module basis for $H_T^*(M; \mathbb{F})$ satisfying Equations (4.3) and (4.4) with respect to the partial order on M^T defined above [10, Proposition 4.4]. In particular, they give a poset-upper-triangular module basis with respect to the partial order on M^T . Moreover the x_p are homogeneous classes of degree $2\rho(p)$. (In fact, their canonical classes form a module basis for $H_T^*(M; \mathbb{Z})$ with integer coefficients, but we will not use that here.)

- (2) Suppose \mathcal{G} is a Kac-Moody group and \mathcal{P} a parabolic subgroup with corresponding flag variety $X = \mathcal{G}/\mathcal{P}$. For $G = T_{\mathcal{G}}/Z(\mathcal{G})$ where $T_{\mathcal{G}}$ is the maximal torus of \mathcal{G} and $Z(\mathcal{G})$ is its center, Equations (4.1) and (4.2) hold for many cases of E_G^* (see e.g. [14]). The set of T -fixed points of \mathcal{G}/\mathcal{P} may be identified with the quotient $W_{\mathcal{G}}/W_{\mathcal{P}}$ where $W_{\mathcal{G}}, W_{\mathcal{P}}$ are the Weyl groups of \mathcal{G}, \mathcal{P} respectively. The index set \mathcal{J} may be identified with this countable quotient and given a poset structure and rank function induced by the Bruhat order and Bruhat length on $W_{\mathcal{G}}$ respectively. In the cases of Borel-equivariant cohomology $E_T^* = H_T^*(-)$ or equivariant K -theory $E_T^* = K_T^*$, the **equivariant Schubert classes** $\{\sigma_w\}_{w \in W_{\mathcal{G}}/W_{\mathcal{P}}}$ corresponding to the Schubert varieties in \mathcal{G}/\mathcal{P} form a module basis satisfying Equations (4.3) and (4.4) (see e.g. [23] and also [34, 35]). Unlike the previous example, this one includes cases of infinite posets \mathcal{J} . Harada-Henriques-Holm give some explicit computations for $\Omega SU(2)$, an infinite-dimensional affine Grassmannian [14].

We close the section with a brief discussion of a more general class of spaces for which poset-upper-triangular bases satisfying conditions (4.3) and (4.4) exist in Borel-equivariant cohomology.

Theorem 4.2. *Let $S \cong S^1$ or $S \cong \mathbb{C}^*$ be a rank-one torus and let X be an S -space satisfying the following conditions:*

- (1) *the set of S -fixed points X^S is isolated;*
- (2) *the set $(X^S, <)$ can be equipped with a poset structure satisfying the conditions of Section 2.1;*
- (3) *the S -equivariant cohomology $H_S^*(X; \mathbb{F})$ is a free $H_S^*(\text{pt}; \mathbb{F})$ -module; and*
- (4) *the ring map $\iota_X^* : H_S^*(X; \mathbb{F}) \rightarrow H_S^*(X^S; \mathbb{F})$ induced by the inclusion $\iota_X : X^S \hookrightarrow X$ is injective.*

Then the module $H_S^(X; \mathbb{F})$ has a poset-upper-triangular basis with respect to any choice of total order compatible with the partial order on X^S .*

Proof. The space X has isolated S -fixed points and $H_S^*(X; \mathbb{F})$ is a free $H_S^*(\text{pt}; \mathbb{F})$ -module isomorphic to a submodule of $\bigoplus_{i \in X^S} H_S^*(\text{pt}; \mathbb{F})$. The ring $H_S^*(\text{pt}; \mathbb{F})$ is isomorphic to the polynomial ring $\mathbb{F}[t]$ in one variable, which is a PID. Hence the sets $\mathcal{V}^i(i)$ in Proposition 2.4 are generated by at most one nonzero element, so the index sets \mathcal{K}_i in Proposition 2.4 have at most one element. The claim follows from Proposition 2.6. \square

Remark 4.3. *Conditions (3) and (4) in Theorem 4.2 will recur in this manuscript, for instance in the definition of GKM-compatibility.*

Remark 4.4. *Many natural topological spaces satisfy the conditions of Theorem 4.2, including many of those described in Remark 4.1. The poset structure in Condition (2) of Theorem 4.2 is often induced from the S -action, as described in Remark 3.5 for complex projective varieties equipped with a T -action and in Remark 4.1 for Hamiltonian T -manifolds. The existence of a paving by complex affine cells, as is sometimes induced by a Bialynicki-Birula decomposition, implies Conditions (3) and (4).*

4.2. Subspaces of GKM spaces. GKM theory is a powerful combinatorial tool that can provide natural and computationally convenient bases for $E_G^*(X)$. However, there are many G -spaces for which the GKM package does not hold. This brings us to the central geometric problem of this manuscript. More specifically, we will describe a geometric framework within which we propose to exploit the GKM theory on an ambient space X in order to analyze the equivariant geometry of a subspace $Y \subseteq X$. We carried out this program in a special case in a previous paper [16]; one of the main goals of the current manuscript is to both generalize and formalize the techniques therein.

Let G be a topological group and suppose that X is a G -space. Throughout this section we assume that the G -action on X is such that the GKM package holds for X as described in Section 4.1. Let $X^G = \cup_{i \in \mathcal{J}} F_i$ denote the set of (isolated) fixed points and let $\{x_i\}_{i \in \mathcal{J}}$ denote the choice of poset-upper-triangular module generators of $E_G^*(X)$.

We wish to analyze subspaces Y of X using GKM theory on X . For this to be feasible, we need to place certain conditions on Y . We introduce the following terminology.

Definition 4.5. Let G be a topological group, X a G -space, and E_G^* an equivariant cohomology theory. Let $Y \subseteq X$ be a subspace of X and $H \subseteq G$ a topological subgroup of G preserving Y . We call the pair (Y, H) **GKM-compatible** with the pair (X, G) with respect to E_G^* if the following conditions hold.

(a) The H -fixed set of Y is the intersection of Y with X^G , i.e.

$$(4.5) \quad Y^H = Y \cap X^G.$$

(b) The H -equivariant cohomology $E_H^*(Y)$ is a free $E_H^*(\text{pt})$ -module.

(c) The ring map $\iota_Y^* : E_H^*(Y) \rightarrow E_H^*(Y^H)$ induced by the inclusion $\iota_Y : Y^H \hookrightarrow Y$ is injective.

When the groups are clear from context, we may simply say that Y is GKM-compatible with X . Similarly, when there is no ambiguity we often neglect to mention the choice of cohomology theory E_G^* .

Remark 4.6. In certain cases, Conditions (b) and (c) are related. For instance, if $G = T$ is a compact torus and $E_T^* = H_T^*(-, \mathbb{F})$ is Borel-equivariant cohomology with coefficients in a field \mathbb{F} of characteristic zero, then Condition (b) implies Condition (c).

Suppose now that (Y, H) is GKM-compatible with (X, G) with respect to E_G^* . Since we assume that $X^G = \cup_{i \in \mathcal{J}} F_i$ consists of isolated fixed points, the intersection $Y \cap X^G$ is indexed by some subset \mathcal{J} of \mathcal{J} . In this setting, the relationship between $E_G^*(X)$ to $E_H^*(Y)$ fits into the algebraic framework discussed in Section 2.3. Indeed, consider first the sequence of ring homomorphisms

$$(4.6) \quad E_G^*(X) \longrightarrow E_H^*(X) \longrightarrow E_H^*(Y)$$

where the first map is the forgetful map associated to the inclusion of groups $H \hookrightarrow G$ and the second is induced from the inclusion of spaces $Y \hookrightarrow X$. (For $X = Y = \text{pt}$, the composition (4.6) specializes to the usual forgetful map $E_G^*(\text{pt}) \rightarrow E_H^*(\text{pt})$.) Moreover, Condition (a) in the definition of GKM-compatibility means that the map (4.6) fits into a commutative diagram

$$(4.7) \quad \begin{array}{ccc} E_G^*(X) & \xrightarrow{\iota^*} & E_G^*(X^G) \cong \prod_{i \in \mathcal{J}} E_G^*(\text{pt}) \\ \downarrow & & \downarrow \pi \\ E_H^*(X) & \xrightarrow{\iota_Y^*} & E_H^*(Y^H = Y \cap X^G) \cong \prod_{i \in \mathcal{J}} E_H^*(\text{pt}) \end{array}$$

where the right arrow π is 0 on the components $i \notin \mathcal{J}$ and is the forgetful map $E_G^*(\text{pt}) \rightarrow E_H^*(\text{pt})$ on the components $i \in \mathcal{J}$. By assumption, the G -space X satisfies the GKM conditions, so the restriction map ι^* is injective. Finally, Condition (c) of the definition of GKM-compatibility assures us that ι_Y^* is an injection, and Condition (b) ensures that we can find a module basis for $E_H^*(Y)$. The commutative diagram (4.7) is thus an instance of the diagram (2.9) in Section 2.3, with $R = E_G^*(\text{pt})$, $R' = E_H^*(\text{pt})$, $M = E_G^*(X)$, and $M' = E_H^*(Y)$. The forgetful map $E_G^*(\text{pt}) \rightarrow E_H^*(\text{pt})$ satisfies the required property (2.8) for $\gamma : R \rightarrow R'$ by naturality. In contrast to Section 2.3, we do *not* assume here that the map $E_G^*(X) \rightarrow E_H^*(Y)$ is surjective. We discuss this further below.

We first give a rich class of examples of GKM-compatible subspaces of GKM spaces.

Remark 4.7. Let $X = G/B$ be the flag variety of a complex reductive algebraic group. As observed in Remark 4.1, X is GKM with respect to the standard action of the maximal torus T of G . In Section 5 we define a family of **Hessenberg varieties** $Y \subseteq X$ and show that in many cases, such as the **regular nilpotent Hessenberg varieties** in classical Lie type and the **Springer varieties** in Lie type A , there is a natural S^1 subtorus of T which preserves Y and makes (Y, S^1) GKM-compatible with the pair (X, T) with respect to Borel-equivariant cohomology with \mathbb{Q} coefficients.

Hessenberg varieties are our primary examples, but the following question arises naturally.

Question 4.8. What are other classes of GKM-compatible subspaces of GKM spaces?

We now give the main geometric question of this manuscript. Let $\{x_i\}_{i \in \mathcal{J}}$ be a poset-upper-triangular module basis of $E_G^*(X)$ and let \bar{x}_i denote the image of x_i under the composition $E_G^*(X) \rightarrow E_H^*(Y)$.

Question 4.9. *Suppose (X, G) is GKM and suppose (Y, H) is GKM-compatible with (X, G) . Under what circumstances can we exploit the GKM theory on X to explicitly construct a computationally convenient module basis for $E_H^*(Y)$ using the images $\{\bar{x}_i\}_{i \in \mathcal{J}}$ in $E_H^*(Y)$?*

Ideally we would like the module basis for $E_H^*(Y)$ to be a linear combination of elements in $\{\bar{x}_i\}_{i \in \mathcal{J}}$ or even to be a subset of $\{\bar{x}_i\}_{i \in \mathcal{J}}$. For this to be possible, we need that

$$(4.8) \quad \text{the ring homomorphism } E_G^*(X) \rightarrow E_H^*(Y) \text{ is a surjection.}$$

Condition (4.8) is not included in the definition of GKM-compatibility because in many applications, we can use other topological data, together with poset pinball, to deduce surjectivity. For instance, when $G = T$ and $E_T^* = H_T^*(-; \mathbb{F})$, the ring surjection in Condition (4.8) follows from a successful game of Betti poset pinball; and to play Betti pinball we need prior knowledge of the target Betti numbers. We give concrete examples of such arguments in the cases of Peterson varieties in classical Lie type in Section 5, and Springer varieties in type A in Section 6. They were also part of our arguments in previous work [16]. We discuss this in more detail in Section 4.3.

Remark 4.10. *In some situations, Condition (4.8) may be seen to hold directly without using Betti-number arguments. For instance, suppose the spaces X and Y in the discussion above are complex algebraic varieties. If there is a G -invariant affine paving of X which restricts to an affine paving of Y (i.e. a subset of the affine cells in the G -invariant affine paving of X yields an affine paving of Y), then Condition (4.8) holds for any subgroup $H \subseteq G$.*

4.3. Borel-equivariant cohomology $E_T^* = H_T^*$ with field coefficients. We now specialize to the case $G = T$ and $E_T^* = H_T^*(-; \mathbb{F})$ Borel-equivariant cohomology with coefficients in a field \mathbb{F} with characteristic zero. Let X denote an ambient T -space satisfying the GKM package and let (Y, T') be a GKM-compatible subspace. Moreover, let $\{x_i\}$ be a poset-upper-triangular basis for $H_T^*(X; \mathbb{F})$ indexed by the set of isolated fixed points $\mathcal{J} = X^T$ and let $\rho_X : \mathcal{J} \rightarrow \mathbb{N}$ be the rank function on the poset \mathcal{J} . Borel-equivariant cohomology is a graded theory, so we may speak of the *degree* of a class x_i . For most of the discussion we assume that

$$(4.9) \quad \text{the ordinary cohomology of the spaces } X \text{ and } Y \text{ vanish in odd degrees.}$$

In practice, this is not a very restrictive condition, but see Remark 4.17 below.

Remark 4.11. *Given the assumptions of this section, and in particular the vanishing condition (4.9), it follows that the Leray-Serre spectral sequence for Borel-equivariant cohomology of Y collapses, so $H_{T'}^*(Y; \mathbb{F})$ is a free $H_{T'}^*(\text{pt}; \mathbb{F})$ -module. By the localization theorem in Borel-equivariant cohomology (e.g. [1, Theorem (3.5)], [13, Theorem 11.4.4]) the inclusion $\iota : Y^{T'} \hookrightarrow Y$ induces an injection*

$$\iota^* : H_{T'}^*(Y; \mathbb{F}) \hookrightarrow H_{T'}^*(Y^{T'}; \mathbb{F}),$$

so Conditions (b) and (c) of GKM-compatibility are automatically satisfied.

We need a homogeneity condition on the classes x_i in the module basis, so we define the following.

Definition 4.12. Let X be as above. The poset-upper-triangular basis $\{x_i\}$ of $H_T^*(X; \mathbb{F})$ is *rank-homogeneous* if each x_i has homogenous degree with respect to the standard \mathbb{Z} -grading on Borel-equivariant cohomology, and if

$$\deg(x_i) = 2 \cdot \rho_X(i) \text{ for all } i \in \mathcal{J}.$$

For instance, the bases in Remark 4.1 for Borel-equivariant cohomology are all rank-homogeneous. We now tackle Question 4.9 in this setting, namely we build a module basis for $H_{T'}^*(Y; \mathbb{F})$ from the basis $\{x_i\}_{i \in \mathcal{J}}$ for $H_T^*(X; \mathbb{F})$. As we said at the end of Section 4.2, we often know the Betti numbers $b_j := \dim_{\mathbb{F}} H^{2j}(Y; \mathbb{F})$ of Y in advance but do not know that the ring map $H_T^*(X; \mathbb{F}) \rightarrow H_{T'}^*(Y; \mathbb{F})$ is surjective. Poset pinball can be useful in this situation: a successful game of Betti pinball using the target Betti numbers $b_j = \dim_{\mathbb{F}} H^{2j}(Y; \mathbb{F})$ can build a module basis for $H_{T'}^*(Y; \mathbb{F})$, from which we may deduce surjectivity. We explain this in the next two propositions. Recall we denote by \bar{x} the image in $H_{T'}^*(Y; \mathbb{F})$ of a class x in $H_T^*(X; \mathbb{F})$.

Proposition 4.13. *Let T be a compact torus and X a T -space for which the GKM package holds. Let $T' \subseteq T$ be a subtorus and suppose (Y, T') is GKM-compatible with (X, T) . We assume $H^*(Y; \mathbb{F})$ is finite-dimensional. Suppose $\{x_i\}_{i \in \mathcal{J}}$ is a set of rank-homogeneous poset-upper-triangular $H_T^*(\text{pt}, \mathbb{F})$ -module generators of $H_T^*(X, \mathbb{F})$. Suppose the ordinary cohomology of Y vanishes in odd degrees and let $b_j := \dim_{\mathbb{F}} H^{2j}(Y; \mathbb{F})$ be the even Betti numbers of Y . Suppose there exists a subset $\mathcal{K} \subseteq \mathcal{J}$ such that the images $\mathcal{B}' := \{\bar{x}_k\}_{k \in \mathcal{K}}$ in $H_{T'}^*(Y; \mathbb{F})$ under the ring map $H_T^*(X; \mathbb{F}) \rightarrow H_{T'}^*(Y; \mathbb{F})$ in (4.6)*

- (A) *are $H_{T'}^*(\text{pt}, \mathbb{F})$ -linearly independent in $H_{T'}^*(Y; \mathbb{F})$, and*
- (B) *for each $j \in \mathbb{Z}_{\geq 0}$ there exist precisely b_j elements in \mathcal{B}' of homogeneous degree $2j$.*

Then \mathcal{B}' is a $H_{T'}^(\text{pt}, \mathbb{F})$ -module basis for $H_{T'}^*(Y)$.*

Proof. We apply [16, Proposition A.1] using $R = H_T^*(\text{pt}; \mathbb{F})$, $M = H_{T'}^*(Y; \mathbb{F})$, and $V = H^*(Y; \mathbb{F})$, where conditions (A) and (B) above are the hypotheses of [16, Proposition A.1]. \square

If a subset \mathcal{K} in the above proposition exists, then the $\{\bar{x}_k\}_{k \in \mathcal{K}}$ form a module basis for $H_{T'}^*(Y; \mathbb{F})$, and so the ring map $H_T^*(X; \mathbb{F}) \rightarrow H_{T'}^*(Y; \mathbb{F})$ is surjective. In other words, we may deduce surjectivity from the Betti numbers—though it may be challenging to find a subset \mathcal{K} . The following proposition shows that poset pinball can sometimes accomplish this.

Proposition 4.14. *Suppose we have $T, X, T', Y, \{x_i\}_{i \in \mathcal{J}}, b_j := \dim_{\mathbb{F}} H^{2j}(Y; \mathbb{F})$ as in Proposition 4.13. Let \mathcal{J} be the T' -fixed points of Y and \mathcal{J} be the T -fixed points of X . Suppose that **either** of the following holds for poset pinball played with ambient poset \mathcal{J} and initial subset \mathcal{J} :*

- (1) *An instance of upper-triangular Betti pinball with target Betti numbers $\mathbf{b} = (b_0, b_1, b_2, \dots)$ is successful.*
- (2) *An instance of Betti pinball with target Betti numbers $\mathbf{b} = (b_0, b_1, b_2, \dots)$ is successful, and the classes corresponding to the roldown set $\{\bar{x}_{\text{roll}(j)}\}_{j \in \mathcal{J}}$ are $H_{T'}^*(\text{pt}; \mathbb{F})$ -linearly independent.*

Then the classes corresponding to the roldown set $\{\bar{x}_{\text{roll}(j)}\}_{j \in \mathcal{J}}$ are an $H_{T'}^(Y; \mathbb{F})$ -module basis for $H_{T'}^*(Y; \mathbb{F})$.*

Proof. If successful, Betti pinball yields the correct nonzero Betti numbers (b_0, b_1, b_2, \dots) by construction. Hence in either case, Condition (B) of Proposition 4.13 is satisfied by the classes corresponding to the roll-downs $\text{roll}(j)$ for $j \in \mathcal{J}$. It remains to verify Condition (A). In the first case, the classes corresponding to the roll-downs are poset-upper-triangular by the rules of upper-triangular pinball; by Proposition 2.6 they are linearly independent. In the second, linear independence is assumed. The result follows. \square

Remark 4.15. *Alternatively, if we know in advance that the map $H_T^*(X; \mathbb{F}) \rightarrow H_T^*(Y; \mathbb{F})$ is surjective, then it would be possible to deduce the Betti numbers of Y from a successful game of upper-triangular pinball together with an argument that the classes corresponding to the roll-downs satisfy the conditions of Proposition 2.4.*

Remark 4.16. *Betti pinball often yields a set of elements in $H_{T'}^*(Y; \mathbb{F})$ which are not upper-triangular with respect to the original partial order $<$ on \mathcal{J} . In fact, Example 6.14 shows a case in which no successful game of Betti pinball yields a poset-upper-triangular set of elements in $H_{T'}^*(Y; \mathbb{F})$. However we can frequently find a total order \prec compatible with the original partial order, with respect to which the classes associated to the roldown set are upper-triangular. Applying Proposition 2.6 with respect to the total order \prec we conclude that these classes are linearly independent. If Betti pinball is successful in this case, Proposition 4.14 guarantees that the set forms a module basis. In our view, this indicates that except in particularly nice settings such as the partial flag varieties G/P , the notion of poset-upper-triangularity that is geometrically natural may not coincide with the notion that is combinatorially natural (as articulated by Billey).*

Remark 4.17. *It is sometimes possible to apply the theory of GKM spaces and GKM-compatible subspaces without Assumption (4.9) on the ordinary cohomology of X and Y . For instance “mod 2” GKM theory (e.g. [15] and references therein) can deal with real projective spaces $\mathbb{R}P^n$ which have non-vanishing cohomology in odd degree (working with coefficients in $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$). It would be possible to rephrase our theorems and the game of poset pinball to account for these ‘mod 2’ GKM spaces and Borel-equivariant cohomology with $\mathbb{Z}/2\mathbb{Z}$ coefficients, but we have chosen to simplify exposition by assuming (4.9).*

We close this section by addressing Question 4.9 via **matchings** with respect to different integer functions on the underlying posets. This is a complementary approach to pinball that is easier to use in some contexts. Betti pinball does not determine the degree $\rho_X(\text{roll}(j))$ of the roldown of the vertex $j \in \mathcal{J}$. However, in

some cases, geometric considerations on the subspace Y naturally give rise to a function $\deg_Y : \mathcal{J} \rightarrow \mathbb{Z}_{\geq 0}$ with the property that

$$b_k = \dim H^{2k}(Y; \mathbb{F}) = |\{j \in \mathcal{J} : \deg_Y(j) = k\}|.$$

For example, as in the setting of Remark 4.7, many varieties Y have a paving by complex affine cells; in this situation, it is natural to define $\deg_Y(j)$ to be the complex dimension of the affine cell of Y containing the T' -fixed point of Y associated to $j \in \mathcal{J}$. Given \deg_Y we could define a new version of poset pinball in which we require that roll downs satisfy $\rho_X(\text{roll}(j)) = \deg_Y(j)$. However, instead of taking this approach, we construct module bases for $H_{T'}^*(Y; \mathbb{F})$ directly, using *matchings* compatible with rank functions.

Theorem 4.18. *Let T be a compact torus and X a T -space for which the GKM package holds. Let $T' \subseteq T$ be a subtorus and suppose (Y, T') is GKM-compatible with (X, T) . We assume $H^*(Y; \mathbb{F})$ is finite-dimensional. Suppose $\{x_i\}_{i \in \mathcal{J}}$ is a set of rank-homogeneous poset-upper-triangular $H_{T'}^*(\text{pt}, \mathbb{F})$ -module generators of $H_{T'}^*(X, \mathbb{F})$ with $\min(x_i) = i$ for each i . Suppose the ordinary cohomology of Y vanishes in odd degrees and let $b_k := \dim_{\mathbb{F}} H^{2k}(Y; \mathbb{F})$ be the even Betti numbers of Y . Suppose $\deg_Y : \mathcal{J} \rightarrow \mathbb{Z}_{\geq 0}$ is a function with*

$$|\{j \in \mathcal{J} : \deg_Y(j) = k\}| = b_k.$$

Suppose there is an injection $f : \mathcal{J} \rightarrow \mathcal{J}$ with $\deg_Y(j) = \rho_X(f(j))$ and a total ordering \prec compatible with the partial order $<$ with respect to which $\{\bar{x}_{f(j)}\}$ is upper-triangular. Then $\{\bar{x}_{f(j)}\}$ is a module basis for $H_{T'}^(Y; \mathbb{F})$.*

If it exists, we call the map $f : \mathcal{J} \rightarrow \mathcal{J}$ a *matching*.

Proof. By hypothesis, the set $\{\bar{x}_{f(j)}\}_{j \in \mathcal{J}}$ is upper-triangular with respect to the total ordering \prec compatible with $<$. Proposition 2.6 then implies that $\{\bar{x}_{f(j)}\}_{j \in \mathcal{J}}$ is linearly independent. The matching condition implies that for each $k \in \mathbb{Z}_{\geq 0}$ there exist precisely b_k elements in $\{\bar{x}_{f(j)}\}_{j \in \mathcal{J}}$ of homogeneous degree $2k$. Thus the set $\{\bar{x}_{f(j)}\}_{j \in \mathcal{J}}$ satisfies the hypotheses of [16, Proposition A.1], and the result follows. \square

Remark 4.19. *Module bases constructed from a matching may be very different from bases obtained by poset pinball if the degree function \deg_Y is not compatible with the partial order on \mathcal{J} . On the other hand, if the degree function $\deg_Y : \mathcal{J} \rightarrow \mathbb{Z}_{\geq 0}$ also satisfies*

$$(4.10) \quad \deg_Y(j) \leq \rho_X(j) \text{ for all } j \in \mathcal{J},$$

namely $2 \deg_Y(j)$ is bounded by the cohomology degree of x_j in $H^(X)$, then a matching basis could arise as a poset pinball basis.*

5. EXAMPLE: PETERSON VARIETIES AND OTHER REGULAR NILPOTENT HESSENBERG VARIETIES

Regular nilpotent Hessenberg varieties are a family of subvarieties of the flag varieties G/B which fit into the geometric framework of Section 4.2, so we can analyze their equivariant cohomology using poset pinball. We discuss facts about regular nilpotent Hessenberg varieties in different Lie types in Section 5.1. Then in Section 5.2 we explicitly calculate the Borel-equivariant cohomology of Peterson varieties, a collection of regular nilpotent Hessenberg varieties. To do this, we find a module basis, and show the basis may be obtained either via poset pinball (as in Proposition 4.14) or by a matching (as in Theorem 4.18). We studied the case of Lie type A previously [16]; the results here generalize that earlier work to all classical Lie types. In this section we always work in Borel-equivariant cohomology with coefficients in a field \mathbb{F} of characteristic zero.

5.1. Background on regular nilpotent Hessenberg varieties. Let G be a complex reductive linear algebraic group, and let B and $T \subseteq B$ denote choices of a Borel subgroup and a maximal torus of G , respectively. We denote by \mathfrak{g} and \mathfrak{b} the associated Lie algebras of G and B . The homogeneous space G/B is a generalized flag variety. A linear subspace $H \subseteq \mathfrak{g}$ is called a **Hessenberg space** if

- H contains the Lie algebra \mathfrak{b} , and
- H is closed under Lie bracket with \mathfrak{b} , i.e. $[H, \mathfrak{b}] \subseteq H$.

Let $N \in \mathfrak{g}$. The **Hessenberg variety** $\mathcal{H}(N, H)$ associated to N and H is the subvariety of G/B defined by

$$(5.1) \quad \mathcal{H}(N, H) := \{gB : \text{Ad}(g^{-1})(N) \in H\} \subseteq G/B.$$

When N is regular (also called principal) nilpotent, then the Hessenberg variety $\mathcal{H}(N, H)$ is called a **regular nilpotent Hessenberg variety**.

Let Φ denote the set of roots of \mathfrak{g} and $\Phi^+ \subset \Phi$ be the set of positive roots corresponding to \mathfrak{b} . Let $\Delta = \{\alpha_i\}$ denote the set of simple roots in Φ^+ . If $\alpha \in \Phi$ is a root, let \mathfrak{g}_α be its corresponding root space. Fix a basis element E_α for each \mathfrak{g}_α . Let W denote the Weyl group associated to G . We use the natural action of the maximal torus T on G/B given by left multiplication on cosets. The fixed point set $(G/B)^T$ may be naturally identified with the Weyl group W .

We begin with some useful facts.

Lemma 5.1. (1) *Any regular nilpotent Lie algebra element $N \in \mathfrak{g}$ is G -conjugate to the regular nilpotent element of the form*

$$(5.2) \quad N_0 := \sum_{\alpha_i \in \Delta} E_{\alpha_i}.$$

(2) *Suppose $H \subseteq \mathfrak{g}$ is a Hessenberg space $H \subseteq \mathfrak{g}$ and $N_1, N_2 \in \mathfrak{g}$ are G -conjugate. The corresponding varieties $\mathcal{H}(N_1, H)$ and $\mathcal{H}(N_2, H)$ are isomorphic, with explicit isomorphism*

$$gB \in \mathcal{H}(N_1, H) \leftrightarrow g_2gB \in \mathcal{H}(N_2, H),$$

where $g_2 \in G$ satisfies $N_1 = \text{Ad}(g_2^{-1})(N_2)$.

(3) *If $N \in \mathfrak{g}$ is a sum of simple root vectors, there exists a circle subgroup S^1 of the maximal torus T such that the restriction of the natural T -action on G/B to the S^1 -subgroup preserves $\mathcal{H}(N, H)$. Moreover, the points in $\mathcal{H}(N, H)$ that are fixed by this S^1 -action satisfy*

$$(5.3) \quad (\mathcal{H}(N, H))^{S^1} = \mathcal{H}(N, H) \cap (G/B)^T.$$

Proof. Part (1) is a standard result (see e.g. Collingswood-McGovern [7, Theorem 4.1.6]).

Part (2) is a straightforward consequence of Definition (5.1).

To prove (3), we explicitly construct the required subgroup S^1 . By definition \mathfrak{g}_α is an eigenspace for the action of $\text{Ad } T$ with eigenfunction $\alpha : T \rightarrow \mathbb{C}^*$. This means that $\text{Ad } t(x) = \alpha(t)x$ for all $x \in \mathfrak{g}_\alpha$ and that α is a character of T , which we think of as an element of \mathfrak{t}^* . (See also [17, 16.4].) The characters $\alpha_1, \alpha_2, \dots, \alpha_n$ form a maximal \mathbb{Z} -linearly independent set in \mathfrak{t}^* by definition of simple roots, so the map $\phi : T \rightarrow (\mathbb{C}^*)^n$ given by $\phi(t) = (\alpha_1(t), \alpha_2(t), \dots, \alpha_n(t))$ is an isomorphism of linear algebraic groups.

In particular, the preimage of the diagonal subgroup $\{(c, c, \dots, c) \mid c \in \mathbb{C}^*\}$ is a rank-one subtorus $S \cong \mathbb{C}^*$ of T whose elements t_c are parametrized by c . The elements of S also satisfy

$$(\text{Ad } t_c) \left(\sum E_{\alpha_i} \right) = \sum c E_{\alpha_i} = c \left(\sum E_{\alpha_i} \right)$$

for all $c \in \mathbb{C}^*$ and any sum of simple root vectors, since each $E_\alpha \in \mathfrak{g}_\alpha$. In particular $\text{Ad } (g^{-1}) (\text{Ad } (t_c^{-1})N)$ and $\text{Ad } (g^{-1})N$ differ by a multiple of the nonzero scalar c . Since H is a vector space we have

$$t_c gB \in \mathcal{H}(N, H) \iff gB \in \mathcal{H}(N, H).$$

We now confirm that $(G/B)^S = (G/B)^T$. We saw that the composition of the maps S and α_i send $c \mapsto t_c \mapsto \alpha_i(t_c) = c$ so the composition has degree one for each simple root α_i . Under the natural pairing of characters and one-parameter subgroups [17, 16.1], we have

$$(5.4) \quad \langle S, \alpha_i \rangle = 1 \text{ for all simple roots } \alpha_i.$$

This implies that S is a regular subgroup [17, 24.4], from which $(G/B)^S = (G/B)^T$ follows [17, Section 24, Exercise 6]. This in turn implies Equation (5.3). Finally, to obtain a real rank-one torus, we may restrict to the unit-length elements in \mathbb{C}^* . \square

For the rest of this section, we assume that $N = N_0$, which by Lemma 5.1 results in no loss of generality. We will also assume that the S^1 -action on $\mathcal{H}(N_0, H)$ is that constructed in Lemma 5.1.

Our next goal is to explicitly describe the S^1 -fixed points in $\mathcal{H}(N_0, H)$. By Equation (5.3), this is equivalent to identifying the T -fixed points in G/B that lie in $\mathcal{H}(N_0, H)$. The next proposition does this in arbitrary Lie type. We need the following notation. Given a Hessenberg space H , let \mathcal{M}_H denote the set of roots defined by the condition

$$(5.5) \quad H = \mathfrak{b} \oplus \bigoplus_{\alpha \in \mathcal{M}_H} \mathfrak{g}_\alpha.$$

For each $w \in W = N(T)/T$, choose a representative $\tilde{w} \in N(T)$. The coset $\tilde{w}B$ is independent of the choice of representative \tilde{w} since $T \subseteq B$, so we denote it wB . Recall the T -fixed points in G/B are the flags $\{wB : w \in W\}$.

Proposition 5.2. *Let \mathfrak{g} be of arbitrary Lie type. Let $H \subseteq \mathfrak{g}$ be a Hessenberg space and let $\mathcal{H}(N_0, H)$ be the regular nilpotent Hessenberg variety corresponding to H and N_0 . Then the flag $wB \in (G/B)^T$ is in the regular nilpotent Hessenberg variety $\mathcal{H}(N_0, H)$ if and only if $w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+$.*

Proof. The element wB is in $\mathcal{H}(N_0, H)$ if and only if $\text{Ad}(\tilde{w}^{-1})(N_0) \in H$ for any representative $\tilde{w} \in N(T)$ of $w \in W$. Since $\text{Ad}(\tilde{w}^{-1})(N_0) = \sum_{\alpha_i \in \Delta} E_{w^{-1}\alpha_i}$ we have $wB \in \mathcal{H}(N_0, H)$ exactly if $w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+$. \square

We next recall a result which allows us to deduce the Betti numbers of $\mathcal{H}(N_0, H)$. The original and stronger result, restated below, is that certain nilpotent Hessenberg varieties are *paved by (complex) affines*.

Lemma 5.3. (**[31, Theorem 6.1] and [32, Theorem 4.3]**) *Assume either that*

- (1) *the Lie algebra \mathfrak{g} is of classical Lie type and $N = N_0$ is the regular nilpotent element $N_0 = \sum_{\alpha_i \in \Delta} E_{\alpha_i}$ or*
- (2) *the Lie algebra \mathfrak{g} is of Lie type A and N is a nilpotent linear operator in Jordan canonical form.*

Let $H \subseteq \mathfrak{g}$ be a Hessenberg space and let $\mathcal{H}(N, H)$ denote the Hessenberg variety corresponding to H and N . Then $\mathcal{H}(N, H)$ has a paving by complex affines obtained by intersecting with an appropriate Bruhat decomposition $\bigcup \mathcal{C}_w$ of G/B . The homology classes corresponding to the subspaces $\overline{\mathcal{C}_w \cap \mathcal{H}(N, H)}$ generate $H_(\mathcal{H}(N, H))$.*

Moreover in Case (1), the intersection $\mathcal{C}_w \cap \mathcal{H}(N_0, H)$ is nonempty exactly when $w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+$. The degree of the homology class corresponding to w is

$$2|\{\alpha \in \Phi^+ : w^{-1}(\alpha) \in \mathcal{M}_H\}|.$$

In particular, the homology of $\mathcal{H}(N_0, H)$ is \mathbb{Z} -torsion-free, and nonzero only in even degree. It follows that the $2j^{\text{th}}$ Betti number $b_{2j} = \dim_{\mathbb{F}} H^{2j}(\mathcal{H}(N_0, H); \mathbb{F})$ is

$$(5.6) \quad b_{2j} = |\{w \in W : w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+ \text{ and } j = |\{\alpha \in \Phi^+ : w^{-1}(\alpha) \in \mathcal{M}_H\}|||}.$$

We saw in Remark 4.1 that the T -space G/B satisfies the GKM package of Section 4.1 and that the cohomology ring $H_T^*(G/B; \mathbb{F})$ has a well-known set of poset-upper-triangular generators with respect to the Bruhat order on $(G/B)^T \cong W$: the **equivariant Schubert classes** $\{\sigma_v\}_{v \in W}$. Our goal is to construct computationally convenient module bases for $H_{S^1}^*(\mathcal{H}(N_0, H); \mathbb{F})$ using the equivariant Schubert classes, according to the point of view laid out in previous sections. For this we need the following preliminary observation, which we state in more generality than we use here.

Theorem 5.4. *Suppose N is the regular nilpotent operator $\sum_{\alpha_i \in \Delta} E_{\alpha_i}$ in classical Lie type, or a nilpotent linear operator in Jordan canonical form in Lie type A . The pair $(\mathcal{H}(N, H), S^1)$ is GKM-compatible with the pair $(G/B, T)$ with respect to Borel-equivariant cohomology $H^*(-; \mathbb{F})$ with coefficients in a field \mathbb{F} of characteristic zero.*

Proof. A matrix in Jordan canonical form in Lie type A is by definition a sum of simple root vectors such as the E_{α_i} above. Equation (5.3) of Lemma 5.1 Part (3) gives the first condition of GKM compatibility, namely Equation (4.5) in the case of $\mathcal{H}(N, H)$. The complex paving by affines of $\mathcal{H}(N, H)$ described in Lemma 5.3 implies that the ordinary cohomology of $\mathcal{H}(N, H)$ is zero in odd degrees. The result follows from the argument in Remark 4.11. \square

In order to effectively compute a module basis for $H_{S^1}^*(\mathcal{H}(N_0, H); \mathbb{F})$, we need more information about the components $\sigma_v(w)$ of the equivariant Schubert classes in the direct sum

$$(5.7) \quad H_T^*(G/B; \mathbb{F}) \hookrightarrow H_T^*((G/B)^T; \mathbb{F}) \cong \bigoplus_{w \in W} H_T^*(\text{pt}; \mathbb{F}) \cong \bigoplus_{w \in W} \text{Sym}_{\mathbb{F}}(\mathfrak{t}^*),$$

where $\text{Sym}_{\mathbb{F}}(\mathfrak{t}^*)$ denotes the ring of polynomials with coefficients in the field \mathbb{F} on the Lie algebra \mathfrak{t} . Billey gave a complete description of the polynomial $\sigma_v(w)$ for any $v, w \in W$ in arbitrary Lie type [3, Theorem 4]. We will only need the following consequences of her formula.

Proposition 5.5. (Corollaries of Billey's formula [3, Theorem 4]) *Let $v, w \in W$. Let $\sigma_v \in H_T^*(G/B; \mathbb{F})$ be the equivariant Schubert class corresponding to v . Then:*

- Given a reduced word decomposition of w , the component $\sigma_v(w)$ is a sum of terms, with one summand for each reduced subword of w that equals v . In particular,

$$\sigma_v(w) = 0$$

if $w \not> v$ in Bruhat order.

- Suppose $w > v$ in Bruhat order. Each summand in $\sigma_v(w)$ is a monomial in the positive roots Φ^+ with a positive integer coefficient.

In this context, the commutative diagram (4.7) becomes

$$(5.8) \quad \begin{array}{ccc} H_T^*(G/B; \mathbb{F}) & \hookrightarrow & H_T^*((G/B)^T; \mathbb{F}) \cong \bigoplus_{w \in W} \text{Sym}_{\mathbb{F}}(\mathfrak{t}^*) \\ \pi_{G/B} \downarrow & & \downarrow \pi \\ H_{S^1}^*(\mathcal{H}(N_0, H); \mathbb{F}) & \hookrightarrow & H_{S^1}^*((\mathcal{H}(N_0, H))^{S^1}; \mathbb{F}) \cong \bigoplus_{\substack{w \in W \text{ such that} \\ w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+}} \text{Sym}_{\mathbb{F}}(\text{Lie}(S^1)^*) \end{array}$$

where the left vertical arrow $\pi_{G/B}$ is the composition of the natural maps $H_T^*(G/B; \mathbb{F}) \rightarrow H_{S^1}^*(G/B; \mathbb{F})$ and $H_{S^1}^*(G/B; \mathbb{F}) \rightarrow H_{S^1}^*(\mathcal{H}(N_0, H); \mathbb{F})$ as in Section 4.2. For $v \in W$ let

$$p_v := \pi_{G/B}(\sigma_v) \in H_{S^1}^*(\mathcal{H}(N_0, H); \mathbb{F})$$

denote the image of a Schubert class σ_v under $\pi_{G/B}$. Given this setup, the following proposition—which holds in arbitrary Lie type—is a straightforward consequence of Proposition 5.5.

Proposition 5.6. *Let $H \subseteq \mathfrak{g}$ be a Hessenberg space and let $\mathcal{H}(N_0, H)$ denote the regular nilpotent Hessenberg variety corresponding to H and N_0 . Let $w \in W$ satisfy $w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+$. For $v \in W$,*

- $p_v(w) = 0$ if $w \not> v$ in Bruhat order, and
- $p_v(w) \neq 0$ if $w > v$ in Bruhat order.

Proof. Since S^1 is a torus of rank one, the symmetric algebra $\text{Sym}_{\mathbb{F}}(\text{Lie}(S^1)^*)$ may be identified with a polynomial ring in one variable. Denote this variable t . Given $S^1 \subseteq T$ constructed in Lemma 5.1 Part (3), consider the natural map $H_T^*(\text{pt}; \mathbb{F}) \rightarrow H_{S^1}^*(\text{pt}; \mathbb{F})$. The induced map $\text{Sym}_{\mathbb{F}}(\mathfrak{t}^*) \rightarrow \text{Sym}_{\mathbb{F}}(\text{Lie}(S^1)^*)$ sends each simple root $\alpha_i \in \mathfrak{t}^* \subseteq \text{Sym}_{\mathbb{F}}(\mathfrak{t}^*)$ to $t \in \text{Sym}_{\mathbb{F}}(\text{Lie}(S^1)^*)$ by Equation (5.4). Moreover, the arrow labeled π in Equation (5.8) is defined by restricting the class $(p(w))_{w \in W} \in H_T^*((G/B)^T; \mathbb{F})$ to the components indexed by Weyl group elements with $w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+$. In other words π sends $(p(w))_{w \in W}$ to $(p(w))_{w \in W: w^{-1}\Delta \subseteq \mathcal{M}_H \cup \Phi^+}$. Proposition 5.5 now implies that $p_v(w)$ is either zero or a polynomial in t with positive integer coefficients, so the claim follows. \square

5.2. The S^1 -equivariant cohomology of the Peterson variety in classical Lie types. In this section, we explicitly build module bases for the S^1 -equivariant cohomology of **Peterson varieties** in classical Lie type. These are special cases of the regular nilpotent Hessenberg varieties in Section 5.1, for which the Hessenberg space is chosen to be

$$(5.9) \quad H_{\Delta} := \mathfrak{b} \oplus_{\alpha \in -\Delta} \mathfrak{g}_{\alpha}.$$

In other words $\mathcal{M}_H = -\Delta$. For notational simplicity, we fix \mathfrak{g} and denote the corresponding Peterson variety

$$\mathcal{Y} := \mathcal{H}(N_0, H_{\Delta}).$$

The set of S^1 -fixed points \mathcal{Y}^{S^1} is a key ingredient in the combinatorial constructions from Section 3, since it corresponds to the initial subset $\mathcal{J} \subseteq \mathcal{J}$ with $\mathcal{J} = W \cong (G/B)^T$. Proposition 5.2 characterizes the Weyl group elements $w \in W$ whose flags wB are in an arbitrary regular nilpotent Hessenberg variety; Proposition 5.8 refines this characterization for Peterson varieties. We need the following lemma, which we give for convenience, though it is probably familiar to experts. We follow the notation of Section 5.1. Also, given a subset $J \subseteq \Delta$ we denote by W_J the corresponding Weyl group, and by Φ_J (respectively Φ_J^+ or Φ_J^-) the corresponding root system (respectively positive or negative roots).

Lemma 5.7. *Let Φ be a finite root system of arbitrary Lie type with Weyl group W . Let $\Delta = \{\alpha_i\}$ denote the simple roots in a choice of positive roots Φ^+ of Φ . Let $w \in W$ and define*

$$(5.10) \quad J := \{\alpha_i : w^{-1}(\alpha_i) < 0\}.$$

Then

$$(5.11) \quad w^{-1}\Delta \subseteq -\Delta \cup \Phi^+$$

if and only if w is the maximal element of the Weyl group W_J .

Proof. Bourbaki proves that if w_J is the (unique) maximal element of W_J then $\ell(w_J) = \Phi_J^+$ and w_J^{-1} sends Δ_J to $-\Delta_J$ [4, Corollary 3 in VI.1.1.6]. Every element of Φ_J^+ is a linear combination of the simple roots Δ_J with nonnegative coefficients, so $w_J^{-1}(\Phi_J^+) = \Phi_J^-$ by linearity. The length of w_J is the number of positive roots that w_J^{-1} sends to negative roots [4, Corollary 2 in VI.1.1.6] so $w_J^{-1}(\Phi^+ - \Phi_J^+) \subseteq \Phi^+$. We conclude $w_J^{-1}\Delta \subseteq -\Delta \cup \Phi^+$ as desired.

Conversely, suppose $w \in W$ and $w^{-1}\Delta \subseteq -\Delta \cup \Phi^+$. We first show that $w^{-1}(\Phi^+ - \Phi_J^+) \subseteq \Phi^+$. Choose $\alpha = \sum_{\alpha_i \in \Delta} c_i \alpha_i$ to be an arbitrary positive root with $w^{-1}(\alpha) < 0$. (In particular, each coefficient c_i is non-negative.) Since w^{-1} is linear on Φ we obtain

$$(5.12) \quad w^{-1}(\alpha) = \sum_{\alpha_i \in \Delta} c_i w^{-1}(\alpha_i) = \sum_{\alpha_k \notin J} c_k w^{-1}(\alpha_k) + \sum_{\alpha_j \in J} c_j w^{-1}(\alpha_j) < 0.$$

Now let $\alpha_i \in \Delta$ with $\alpha_i \notin J$, so $w^{-1}(\alpha_i) > 0$. If the coefficient c_i of α_i is strictly positive then

$$(5.13) \quad 0 < c_i w^{-1}(\alpha_i) \leq \sum_{\alpha_k \notin J} c_k w^{-1}(\alpha_k) < \sum_{\alpha_j \in J} -c_j w^{-1}(\alpha_j),$$

where the last inequality follows from Equation (5.12). Each summand in Equation (5.13) is non-negative by definition of J and the fact that the coefficients c_j, c_k are non-negative. For each $j \in J$ the root $-w^{-1}(\alpha_j)$ is in Δ by the hypothesis that $w^{-1}\Delta \subseteq -\Delta \cup \Phi^+$. Equation (5.13) now implies that $w^{-1}(\alpha_i)$ is a linear combination of $\{w^{-1}(\alpha_j)\}_{\alpha_j \in J}$. This contradicts the fact that $w^{-1}(\Delta)$ is a linearly independent set of roots. We conclude that $c_i = 0$ for all $i \notin J$, from which it follows that $w^{-1}(\Phi^+ - \Phi_J^+) \subseteq \Phi^+$.

To complete the proof, we will show that $w^{-1}w_J = e$. We saw that $w^{-1}(\Phi^+ - \Phi_J^+) \subseteq \Phi^+$ and as before we know that $w^{-1}(\Phi_J^+) \subseteq \Phi^-$ by linearity. Hence $\ell(w) = |\Phi_J^+| = \ell(w_J)$. Recall that $w_J = w_J^{-1}$ [4, Corollary 3 in VI.1.1.6]. We already saw that $w_J(\Phi_J^+) = -\Phi_J^+$ and so $w_J(\Phi^+ - \Phi_J^+) = \Phi^+ - \Phi_J^+$. We now show that $w^{-1}w_J(\alpha_i) > 0$ for all $i = 1, 2, \dots, n$. If $i \in J$ then since

$$w^{-1}w_J(\Phi_J^+) = w^{-1}(-\Phi_J^+) \subseteq \Phi^+$$

we know $w^{-1}w_J(\alpha_i) > 0$. If $i \notin J$ then $w^{-1}w_J(\alpha_i) = w^{-1}(\alpha)$ for some $\alpha \in \Phi^+ - \Phi_J^+$. The element w^{-1} has length $|\Phi_J^+|$ and sends exactly the positive roots Φ_J^+ to negative roots, so $w^{-1}(\alpha) > 0$. Thus $w^{-1}w_J$ sends no simple root to a negative root. It follows that $w^{-1}w_J = e$ [17, 10.2, Corollary to Lemma C], as desired. \square

Combining the previous lemma and Proposition 5.2 immediately gives the following.

Proposition 5.8. *Let \mathfrak{g} be of arbitrary Lie type. Fix the Hessenberg space $H_\Delta := \mathfrak{b} \oplus_{\alpha \in -\Delta} \mathfrak{g}_\alpha$ and let \mathcal{Y} be the Peterson variety corresponding to H_Δ and N_0 . Let $\Phi^+ \subseteq \Phi$ be the positive roots corresponding to \mathfrak{b} in the root system Φ of \mathfrak{g} and let $\Delta \subseteq \Phi^+$ denote the simple roots. Then the S^1 -fixed points \mathcal{Y}^{S^1} of \mathcal{Y} are in one-to-one correspondence with subsets J of Δ , with explicit bijection given by*

$$(5.14) \quad J \subseteq \Delta \longleftrightarrow \text{the maximal element } w_J \text{ in } W_J \longleftrightarrow w_J B \in \mathcal{Y}^{S^1}.$$

Now assume \mathfrak{g} has classical Lie type. In the next result, we combine the concrete characterization of \mathcal{Y}^{S^1} with the Betti numbers from Lemma 5.3 to construct a roldown $roll(w) \in W$ for each element w in the initial subset $\mathcal{J} = \mathcal{Y}^{S^1}$. We show that the roldowns can arise from a successful game of upper-triangular Betti pinball, so the set $\{p_{roll(w)}\}_{w \in \mathcal{Y}^{S^1}}$ forms an $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis for $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$.

We denote by $s_i \in W$ the simple reflection corresponding to the simple root α_i in Δ .

Theorem 5.9. *Let \mathfrak{g} be of classical Lie type. Fix the Hessenberg space $H_\Delta := \mathfrak{b} \oplus_{\alpha \in -\Delta} \mathfrak{g}_\alpha$ and let \mathcal{Y} be the Peterson variety corresponding to H_Δ and N_0 . For each subset $J \subseteq \Delta$ let w_J be maximal Weyl group element of W_J . Suppose $J = \{\alpha_{i_1}, \alpha_{i_2}, \dots, \alpha_{i_s} : i_1 < i_2 < \dots < i_s\}$. Let*

$$(5.15) \quad v_J := s_{i_1} s_{i_2} \dots s_{i_s} \in W.$$

Then the association $w_J \mapsto v_J$ for $J \subseteq \Delta$ is a possible outcome of a successful game of upper-triangular Betti pinball, where $v_J = roll(w_J)$. In particular, the equivariant cohomology classes $\{p_{v_J}\} \subseteq H_{S^1}^(\mathcal{Y})$ form a $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis for $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$.*

Proof. We prove the claim by playing upper-triangular Betti pinball. The board \mathcal{J} is the Weyl group W , identified with the set of T -fixed points on G/B and equipped with Bruhat order. We use the rank function on $\mathcal{J} = W$ given by $\rho(w) = \ell(w)$, the Bruhat length. The initial subset is $\mathcal{J} = \{w_J : J \subseteq \Delta\} \cong \mathcal{Y}^{S^1}$. The inclusion $J' \subseteq J$ implies that $w_{J'} \in W_{J'}$. This in turn implies $w_{J'} \leq w_J$ since w_J is maximal in $W_{J'}$. In other words, Bruhat order induces the partial order on the initial subset \mathcal{J} given by

$$(5.16) \quad w_{J'} < w_J \Leftrightarrow J' \subset J.$$

Fix any total order $<$ of \mathcal{J} subordinate to this partial order.

Specializing Formula (5.6) in Lemma 5.3 to Peterson varieties, we see that the nonzero Betti numbers of \mathcal{Y} are $b_{2j} = \binom{|\Delta|}{j}$, namely the number of subsets $J \subseteq \Delta$ with $|J| = j$. These b_{2j} are our target Betti numbers.

We now play upper-triangular Betti pinball. We will show that for each w_J our choice $v_J := \text{roll}(w_J)$ satisfies all the rules for basic pinball, upper-triangular pinball, and Betti pinball. Indeed, for all J we have $v_J < w_J$ in Bruhat order by construction, so v_J is a possible basic pinball rolldown for w_J . Second we prove that at each step of upper-triangular pinball, no wall on the board prevents w_J from rolling down to v_J . It suffices to show that if $w_{J'} < w_J$ then the element $v_J \not< w_{J'}$ in Bruhat order. The reflection $s_{\alpha_i} < v_J$ precisely when $\alpha_i \in J$ by construction of the element v_J . Hence $v_J < w_{J'}$ if and only if $J \subseteq J'$, from which it follows that if $w_{J'} < w_J$ then $v_J \not< w_{J'}$. Third, we saw above that $\rho(v_J) = \ell(v_J) = |J|$ and there are precisely $\binom{|\Delta|}{j}$ subsets J with degree $|J|$, so the v_J are also rolldowns in Betti pinball.

Finally, since the equivariant Schubert classes $\{\sigma_w\}_{w \in W}$ are a rank-homogeneous poset-upper-triangular basis with respect to Bruhat order, we conclude from Proposition 4.14 that the classes $\{p_{v_J} : J \subseteq \Delta\}$ form a $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis for $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$, as desired. \square

Remark 5.10. A subset of the images $\{p_w\}_{w \in W}$ of the equivariant Schubert classes generate the ring $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$, so the ring map

$$\pi_{G/B} : H_T^*(G/B; \mathbb{F}) \rightarrow H_{S^1}^*(\mathcal{Y}; \mathbb{F})$$

is surjective when \mathcal{Y} is the Peterson variety corresponding to G of classical Lie type. (This is the map from Diagram (5.8), which is Diagram (4.6) for the special case of regular nilpotent Hessenberg varieties.) We note that Carrell and Kaveh have shown that surjectivity of $\pi_{G/B}$ is equivalent to the statement that $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$ is generated by the Chern classes of B -equivariant vector bundles [6].

We can also construct the module basis $\{p_{v_J}\}_{J \subseteq \Delta}$ in the above theorem from a matching compatible with degrees, as discussed in Section 4.3. The additional ingredient which enables this construction is the geometric data of the dimensions of the affine cells that pave \mathcal{Y} , as recorded in Lemma 5.3.

Theorem 5.11. Let $\mathfrak{g}, \mathcal{Y}, \{w_J : J \subseteq \Delta\}, \{v_J : J \subseteq \Delta\}$ be as in Theorem 5.9. Then the $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis $\{p_{v_J} : J \subseteq \Delta\}$ of $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$ can be obtained via a matching compatible with degrees in the sense of Theorem 4.18.

Proof. Define a degree function $\text{deg}_{\mathcal{Y}} : \mathcal{J} \rightarrow \mathbb{Z}_{\geq 0}$ by

$$\text{deg}_{\mathcal{Y}}(w_J) = (\text{complex dimension of the affine cell } \overline{c_{w_J}} \cap \mathcal{Y} \text{ associated to } w_J \text{ in Lemma 5.3}).$$

Lemmas 5.3 and 5.7 together show that $\text{deg}_{\mathcal{Y}}(w_J) = |J|$. Take the rank function $\rho : \mathcal{J} \rightarrow \mathbb{Z}_{\geq 0}$ to be the usual Bruhat length, namely $\rho(w) = \ell(w)$. (Bruhat length of w equals half the cohomology degree of the Schubert class σ_w .) This means that $\text{deg}_{\mathcal{Y}}(w_J) = \rho(v_J)$. In particular the map $f : \mathcal{J} \rightarrow \mathcal{J}$ given by $f(w_J) = v_J$ satisfies $\text{deg}_{\mathcal{Y}}(w_J) = \rho(v_J) = |J|$. The proof of Theorem 5.9 showed that the $\{p_{v_J}\}$ are a rank-homogeneous poset-upper-triangular basis with respect to any total ordering compatible with $<$. The result follows from Theorem 4.18. \square

In previous work [16], the authors constructed an $H_{S^1}^*(\text{pt}; \mathbb{F})$ -module basis for $H_{S^1}^*(\mathcal{Y}; \mathbb{F})$ without reference to poset pinball, in the case when \mathfrak{g} has Lie type A . In fact, the formula for v_J given in Equation (5.15) generalizes to arbitrary Lie type the explicit formulas for what was called $v_{\mathcal{A}}$ in earlier work [16, Equation (2.7) and Definition 4.1]. We deduce that the basis discussed in [16] in fact arises from poset pinball.

Moreover, our previous paper [16] used the poset pinball basis $\{p_{v_J}\}$ in Lie type A to explicitly analyze the structure constants of $H_{S^1}^*(\mathcal{Y})$ via a kind of Monk's formula in equivariant cohomology. We conclude this section with a question for future work.

Question 5.12. What is an explicit combinatorial formula for the structure constants of $H_{S^1}^*(\mathcal{Y})$ with respect to the basis $\{p_{v_J}\}$ in each classical Lie type?

6. EXAMPLE: SPRINGER VARIETIES IN TYPE A

In this section we analyze a special class of nilpotent Hessenberg varieties in Lie type A : *Springer varieties*, and in particular the *subregular* Springer varieties. The flag variety $GL_n(\mathbb{C})/B$ can be identified with

$$\mathcal{F}lags(\mathbb{C}^n) = \{V_\bullet : 0 \subseteq V_1 \subseteq V_2 \subseteq \cdots \subseteq V_{n-1} \subseteq \mathbb{C}^n \text{ such that } \dim_{\mathbb{C}}(V_i) = i\}.$$

Suppose $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is a nilpotent linear operator and \mathfrak{b} is the standard Borel subalgebra of upper-triangular matrices in \mathfrak{g} . The **Springer variety** \mathcal{S}_N **associated to** N is the Hessenberg variety associated to N and the Hessenberg space $H = \mathfrak{b}$, namely

$$(6.1) \quad \mathcal{S}_N := \mathcal{H}(N, \mathfrak{b}).$$

In Lie type A , this can be expressed as

$$\mathcal{S}_N := \{V_\bullet : NV_i \subseteq V_i \text{ for all } 1 \leq i \leq n\} \subseteq \mathcal{F}lags(\mathbb{C}^n).$$

Springer discovered that the symmetric group S_n acts on the ordinary cohomology $H^*(\mathcal{S}_N; \mathbb{C})$ for any Springer variety [29]. This representation is graded by the degree of the cohomology classes. Springer also showed that the top-dimensional cohomology group is an irreducible representation, and that any irreducible representation of S_n arises in this way. Indeed, the irreducible representation corresponding to a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_s)$ arises from the top-dimensional cohomology of the Springer variety \mathcal{S}_N for N with Jordan canonical form given by Jordan blocks of size $\lambda_1, \lambda_2, \dots, \lambda_s$.

In this section, we use poset pinball to construct an explicit module basis for the S^1 -equivariant cohomology with complex coefficients of subregular Springer varieties of Lie type A . Moreover, we construct an S_n -representation on this explicit module basis and obtain an equivariant Springer representation. Goresky and MacPherson give a related construction for a different torus action [12, Section 7].

6.1. An S^1 -action and the S^1 -fixed points of Springer varieties. In this section, we describe an S^1 -action on arbitrary Springer varieties of Lie type A and make some initial observations on their fundamental properties. For instance, we will see that the fixed points $\mathcal{S}_N^{S^1}$ may be identified with the set of permutations whose descents are in positions given by the partition of n determined by the Jordan canonical form of N . By contrast, Carrell obtains a similar result in general Lie type using a different torus action [5]. In Section 6.3, we will specialize to a particular class of nilpotent operators called the *subregular* operators.

Lemma 5.1 shows that for any $g \in GL_n(\mathbb{C})$ the Springer variety \mathcal{S}_N is homeomorphic to $\mathcal{S}_{g^{-1}Ng}$. We will assume without loss of generality that N is in Jordan canonical form, with Jordan blocks weakly decreasing in size. We denote by λ_N both the partition of n and the Young diagram corresponding to this decomposition of N into Jordan blocks.

In Lemma 5.1 Part (3) we define a circle subgroup of the standard maximal torus T^n of diagonal matrices in $U(n, \mathbb{C})$. It can be described very explicitly in this setting as

$$(6.2) \quad S^1 := \left\{ \left[\begin{array}{cccc} t^n & 0 & \cdots & 0 \\ 0 & t^{n-1} & & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & & t \end{array} \right] \middle| t \in \mathbb{C}, \|t\| = 1 \right\} \subseteq T^n \subseteq U(n, \mathbb{C}).$$

The maximal torus T^n acts canonically on $GL(n, \mathbb{C})/B \cong \mathcal{F}lags(\mathbb{C}^n)$ so $S^1 \subseteq T^n$ also acts naturally. In this case Lemma 5.1 states

$$\mathcal{F}lags(\mathbb{C}^n)^{S^1} = \mathcal{F}lags(\mathbb{C}^n)^{T^n}.$$

We now show that the subgroup S^1 in Equation (6.2) preserves the Springer variety \mathcal{S}_N .

Lemma 6.1. *Let N be a nilpotent operator in Jordan canonical form. Then the subgroup S^1 in Equation (6.2) preserves the Springer variety $\mathcal{S}_N \subseteq \mathcal{F}lags(\mathbb{C}^n)$.*

Proof. Suppose $V_\bullet \in \mathcal{S}_N$ and let $\gamma(t)$ denote the diagonal matrix with entries $t^n, t^{n-1}, t^{n-2}, \dots, t$ along the diagonal, as in Equation (6.2). Observe that

$$N(\gamma(t)V_i) \subseteq \gamma(t)V_i$$

if and only if

$$(\gamma(t)^{-1}N\gamma(t))V_i \subseteq V_i.$$

A simple calculation shows that

$$\gamma(t)^{-1}N\gamma(t) = tN.$$

Moreover $(tN)V_i \subseteq V_i$ if and only if $NV_i \subseteq V_i$ since V_i is linear. We conclude that tV_\bullet is in \mathcal{S}_N as desired. \square

The following proposition is a summary of results in the literature, phrased in our language.

Proposition 6.2. *For each nilpotent operator $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ the Springer variety \mathcal{S}_N has no odd-dimensional cohomology. If N is in Jordan canonical form and S^1 is as in Equation (6.2) then the pair (\mathcal{S}_N, S^1) is GKM-compatible with $(GL_n(\mathbb{C})/B, T)$ with respect to Borel-equivariant cohomology $H^*(-; \mathbb{F})$.*

Proof. Spaltenstein proved that the ordinary cohomology of Springer varieties is zero in odd degrees [28]. The result follows from the argument in Remark 4.11. \square

We now compute the fixed points of the Springer variety \mathcal{S}_N with respect to this S^1 -action. Given a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_s)$ of n the permutation $w \in S_n$ has descents in positions given by λ if

$$w(i) > w(i+1) \Rightarrow i \in \{\lambda_1, \lambda_1 + \lambda_2, \lambda_1 + \lambda_2 + \lambda_3, \dots, \lambda_1 + \lambda_2 + \lambda_3 + \dots + \lambda_s\}.$$

For example, the permutation $w = (24581736)$ has descents in the positions given by $\lambda = (4, 2, 2)$.

Theorem 6.3. *Let $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a nilpotent operator in Jordan canonical form whose Jordan blocks weakly decrease in size. Let λ_N be the corresponding partition of n . The S^1 -fixed points of \mathcal{S}_N are in bijection with the set*

$$\{w \in S_n : w^{-1} \text{ has descents in the positions given by } \lambda_N\}.$$

The bijection sends the permutation w to the fixed point wB , where w also denotes the permutation matrix whose i^{th} column is the standard basis vector $e_{w(i)}$ for all i .

Proof. Since $\mathcal{F}lags(\mathbb{C}^n)^{T^n} = \mathcal{F}lags(\mathbb{C}^n)^{S^1}$ it suffices to find the intersection

$$\mathcal{F}lags(\mathbb{C}^n)^{T^n} \cap \mathcal{S}_N.$$

The T^n -fixed points of $\mathcal{F}lags(\mathbb{C}^n)$ consist precisely of the permutation flags $\{wB : w \in S_n\}$ where w is a permutation matrix in $GL(n, \mathbb{C})$ whose i^{th} column has the standard basis vector $e_{w(i)}$. The definition of Springer varieties in Equation (6.1) says that wB is in \mathcal{S}_N exactly when $w^{-1}Nw$ is upper-triangular. Let $E_{i,j}$ denote the $n \times n$ matrix with 1 in the (i, j) -th entry and 0 in all other entries. The matrix N is in Jordan canonical form, so $N = \sum_{i \notin A} E_{i,i+1}$ where

$$A = \{(\lambda_N)_1, (\lambda_N)_1 + (\lambda_N)_2, \dots, (\lambda_N)_1 + (\lambda_N)_2 + (\lambda_N)_3 + \dots + (\lambda_N)_s\}.$$

(In other words, the sum is over pairs $i, i+1$ in the same part of the partition λ_N .) This means wB is in \mathcal{S}_N if and only if

$$w^{-1} \sum_{i \notin A} E_{i,i+1} w = \sum_{i \notin A} E_{w^{-1}(i), w^{-1}(i+1)} \in \mathfrak{b},$$

or equivalently

$$i \notin A \Rightarrow w^{-1}(i) < w^{-1}(i+1).$$

As desired, this implies w^{-1} has descents in the positions given by the partition λ_N . \square

Example 6.4. *Let $n = 4$ and take N to be the matrix with 2 Jordan blocks each of size 2, so $N = E_{12} + E_{34}$. By Theorem 6.3, the S^1 -fixed points of \mathcal{S}_N are the inverses of the following permutations, written in one-line notation:*

$$1234, 1324, 1423, 2314, 2413, 3412.$$

Informally, the matrix $w^{-1}Nw$ is the sum of $E_{w^{-1}(i), w^{-1}(i+1)}$ over i such that $i, i+1$ are in the same part of the partition λ_N . For example, the matrix corresponding to the fixed point $w^{-1} = 2314$ is $E_{23} + E_{14}$ while the matrix corresponding to $w^{-1} = 3412$ is $E_{34} + E_{12}$.

6.2. The subregular Springer representation. A nilpotent linear operator $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is **subregular** if the partition associated to its Jordan canonical form is $(n-1, 1)$, namely it has one Jordan block of size $n-1$ and one of size 1. If N is subregular, the Springer variety \mathcal{S}_N is called a **subregular Springer variety** and Springer's representation on $H^*(\mathcal{S}_N; \mathbb{C})$ is the **subregular Springer representation**. In this section, we collect results about the subregular Springer variety and representation.

Theorem 6.3 implies that the S^1 -fixed points of the subregular Springer variety are in bijective correspondence with permutations whose descents are given by the partition $(n-1, 1)$. The one-line notation of such a permutation w increases in the first $n-1$ entries. Since the last entry can be any integer between 1 and n , the fixed points in one-line notation are precisely

$$\mathcal{S}_N^{S^1} = \{w_i := 1\ 2\ 3 \cdots i-1\ \hat{i}\ i+1 \cdots n\ i \text{ for each } i \text{ with } 1 \leq i \leq n\},$$

where \hat{i} indicates that the integer i is skipped. Note that w_n is the identity element in S_n .

We describe Garsia-Procesi's construction of the subregular Springer representation, using a classical description of irreducible representations of S_n . A filling of the Young diagram $(n-1, 1)$ with the numbers $1, 2, \dots, n$ without repetition is *row-strict* if each row increases left-to-right. (Hence a row-strict subregular filling is either a Young tableau or has $1, 2, 3, \dots, n-1$ in the top row and 1 in the bottom row.)

Let $M^{(n-1,1)}$ denote the complex vector space whose basis is the set of row-strict fillings with shape $(n-1, 1)$. Define an S_n -action on $M^{(n-1,1)}$ as follows. Given $w \in S_n$ and a row-strict filling T of shape $(n-1, 1)$, define the filling $w(T)$ by:

- For each $i = 1, 2, 3, \dots, n$, place $w(i)$ in the box where T had entry i .
- Reorder each row so it increases left-to-right.

By construction $w(T)$ is row-strict of shape $(n-1, 1)$. This is a well-defined action of S_n on the set of row-strict fillings of shape $(n-1, 1)$, and extends by \mathbb{C} -linearity to a representation of S_n on $M^{(n-1,1)}$. (These claims generalize to arbitrary partitions λ of n , see e.g. [8, Section 7.2].)

Combining several results (see Garsia-Procesi's summary [9, page 84] and e.g. Fulton's text for background [8, Section 7.2]) yields the following.

Proposition 6.5. (Garsia-Procesi) *The subregular Springer representation on $H^*(\mathcal{S}_{(n-1,1)}; \mathbb{C})$ is isomorphic, as an ungraded representation, to the S_n -representation $M^{(n-1,1)}$ defined above. Moreover, suppose $M^{(n-1,1)}$ is given the grading inherited from the cohomology ring $H^*(\mathcal{S}_{(n-1,1)}; \mathbb{C})$. Then the set of $n-1$ vectors*

$$v_j := \begin{array}{|c|c|c|c|} \hline 1 & 2 & \cdots & n \\ \hline j & & & \\ \hline \end{array} - \begin{array}{|c|c|c|c|} \hline 2 & 3 & \cdots & n \\ \hline 1 & & & \\ \hline \end{array}$$

form a \mathbb{C} -basis $\{v_2, v_3, \dots, v_n\}$ for the top-degree graded piece of $M^{(n-1,1)}$. A basis for the zero-degree graded piece is given by the vector

$$v_0 := \sum_{j=1}^n \begin{array}{|c|c|c|c|} \hline 1 & 2 & \cdots & n \\ \hline j & & & \\ \hline \end{array}.$$

We can explicitly compute the character of the subregular Springer representation from this description. The representation preserves degrees, so we analyze the characters on each degree separately. Denote the character of the Springer representation on $H^{2i}(\mathcal{S}_N; \mathbb{C})$ by $\chi^i : S_n \rightarrow \mathbb{Z}$. For instance, if $w \in S_n$ the integer $\chi^1(w)$ is the trace of the linear operator on $H^2(\mathcal{S}_N; \mathbb{C})$ corresponding to w .

Corollary 6.6. *Let S_n act on $H^*(\mathcal{S}_N; \mathbb{C}) \cong M^{(n-1,1)}$ via the subregular Springer representation. Then*

$$\chi^1(w) = \#\{\text{fixed points of } w\} - 1 = \#\{j \in \{1, 2, \dots, n\} : w(j) = j\} - 1$$

for all $w \in S_n$. Also for all $w \in S_n$

$$\chi^0(w) = 1.$$

In particular the subregular Springer representation in the zero-degree piece is the trivial representation.

Proof. The first part of the claim is an exercise from Sagan [26, Exercise 2.12.4], and a nice exercise for the reader, given the explicit basis for $H^2(\mathcal{S}_N; \mathbb{C})$ described in Proposition 6.5.

The second part can be seen by inspection. By definition $w \cdot v_0 = v_0$ for all $w \in S_n$. This means $\chi^0(w) = 1$ for all w , so the zero-degree piece is the trivial representation, as desired. \square

6.3. Lifting the Springer action to the S^1 -equivariant cohomology of subregular Springer varieties. We now lift the Springer representation on the ordinary cohomology $H^*(\mathcal{S}_N; \mathbb{C})$ to an action of S_n on the S^1 -equivariant cohomology $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ in the case when N is subregular. To define the lift, we first build a convenient $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module basis of $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ using upper-triangular Betti pinball.

In this context, the commutative diagram (4.7) becomes:

$$(6.3) \quad \begin{array}{ccc} H_T^*(GL(n, \mathbb{C})/B; \mathbb{C}) & \hookrightarrow & H_T^*((GL(n, \mathbb{C})/B)^T; \mathbb{C}) \cong \prod_{w \in S_n} H_T^*(\text{pt}; \mathbb{C}) \cong \prod_{w \in S_n} \text{Sym}(\mathfrak{t}^*) \\ \downarrow & & \downarrow \\ H_{S^1}^*(\mathcal{S}_N; \mathbb{C}) & \hookrightarrow & H_{S^1}^*(\mathcal{S}_N^{S^1}; \mathbb{C}) \cong \prod_{w \in \mathcal{S}_N^{S^1}} H_{S^1}^*(\text{pt}; \mathbb{C}) \cong \prod_{w \in \mathcal{S}_N^{S^1}} \text{Sym}(\text{Lie}(S^1)^*) \end{array}$$

where the left vertical arrow is the composition

$$H_T^*(GL(n, \mathbb{C})/B; \mathbb{C}) \longrightarrow H_{S^1}^*(GL(n, \mathbb{C})/B; \mathbb{C}) \longrightarrow H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$$

and the right vertical arrow is zero on each component corresponding to $w \notin \mathcal{S}_N^{S^1}$. As before, we denote the equivariant Schubert class corresponding to w in $H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$ by σ_w for each $w \in S_n$.

For each $i = 1, 2, \dots, n-1$, let s_i be the permutation on $\{1, 2, \dots, n\}$ that exchanges i and $i+1$ and leaves the other numbers fixed. (In general Lie type s_i is the reflection s_{α_i} for a choice of simple roots.) The proof of the next theorem is similar to that of Theorem 5.9.

Theorem 6.7. *Let N be a subregular nilpotent linear operator $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ and let \mathcal{S}_N denote its associated subregular Springer variety. Then the association*

$$w_i^{-1} \mapsto v_i := \begin{cases} e & \text{if } i = n \\ s_i & \text{if } 1 \leq i \leq n-1 \end{cases}$$

is an outcome of a successful game of upper-triangular Betti pinball, where $v_i = \text{roll}(w_i^{-1})$. In particular, the classes $\{p_{v_i}\}_{i=1}^n = \{p_e, p_{s_1}, p_{s_2}, \dots, p_{s_{n-1}}\}$ form an $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module basis for $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$.

Proof. Recall that w_n is the identity $e \in S_n$. For each $i = 1, 2, \dots, n-1$ note that

$$(6.4) \quad w_i = s_{n-1} s_{n-2} \cdots s_{i+1} s_i$$

is a reduced word decomposition of w_i . Thus for each $i = 1, 2, \dots, n-1$, we have $w_i > s_i$ and $w_i \not> s_j$ for any $j < i$. Moreover $w_n^{-1} < w_{n-1}^{-1} < \cdots < w_1^{-1}$ so Bruhat order totally orders the S^1 -fixed points. In particular the choice of total order required to play pinball is uniquely determined in this case.

Now we play upper-triangular Betti pinball with board $S^n = (GL(n, \mathbb{C})/B)^{S^1}$ and initial subset $(\mathcal{S}_N)^{S^1}$. We saw that $v_i < w_i^{-1}$ for all i . In upper-triangular pinball, walls are never placed between w_i^{-1} and v_i because if $w_j < w_i^{-1}$ then $j > i$ and so $v_i \not< w_j^{-1}$. Finally $\ell(v_n) = 0$ and $\ell(v_i) = 1$ for $i = 1, 2, \dots, n-1$. Comparing with the Betti numbers of \mathcal{S}_N in Proposition 6.5, we conclude that the $\{v_i\}_{i=1}^n$ are a successful outcome of upper-triangular Betti pinball. Applying Proposition 4.14 gives the claim. \square

With a basis for $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ in hand, we may now discuss our construction of an S_n -representation, for which we depend on a previous construction by Kostant and Kumar of a W -representation on $H_T^*(G/B; \mathbb{C})$ when G is any Kac-Moody group [21]. We work in Lie type A , for which $W = S_n$ and $G/B \cong \text{Flags}(\mathbb{C}^n)$.

In the case of S_n acting on $H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$, Kostant and Kumar's action is defined as follows. As before, denote the u -th coordinate of a class $\sigma \in H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$ by $\sigma(u)$. Let $w, u \in W$, and let σ be any element of $H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$. The element $w \cdot \sigma$ is defined by the equation

$$(6.5) \quad (w \cdot \sigma)(u) := \sigma(uw).$$

This Kostant-Kumar action is defined componentwise, so it commutes with the natural maps induced on $H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$ and $H_T^*((GL(n, \mathbb{C})/B)^T; \mathbb{C})$ by $S^1 \hookrightarrow T$. This implies that Kostant-Kumar's action descends to an action on $H_{S^1}^*(GL(n, \mathbb{C})/B; \mathbb{C})$. (We warn the reader that not all S_n -actions on $H_T^*(G/B; \mathbb{C})$ descend to $H_{S^1}^*(G/B; \mathbb{C})$; Tymoczko analyzes another natural S_n -action that does not [33].)

The main result of this section is that Kostant-Kumar's action gives rise to an S_n -action on $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ when \mathcal{S}_N is a subregular nilpotent Springer variety, and that this action lifts the Springer representation to

$H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$. The first step is to show that Kostant-Kumar's action on $H_{S^1}^*(GL(n, \mathbb{C})/B; \mathbb{C})$ preserves the $H_{S^1}^*(\text{pt}; \mathbb{C})$ -submodule spanned by the elements corresponding to the rolldowns from Theorem 6.7.

The following proposition gives a special case of a more general formula of Kostant-Kumar [21, Proposition 4.24.g]. The interested reader may also prove it using the following special cases of Billey's formula for the classes σ_{s_j} :

$$\sigma_{s_j}(w) = \alpha_j \text{ for } w \in \{s_j\} \cup \{s_j s_i : i \neq j, i = 1, \dots, n-1\} \cup \{s_i s_j : i \neq j \pm 1, i \neq j, i = 1, \dots, n-1\}$$

and

$$\sigma_{s_j}(w) = \alpha_i + \alpha_j \text{ for } w = s_{j-1} s_j \text{ or } s_{j+1} s_j.$$

Proposition 6.8. (Kostant-Kumar) *For each i, j with $1 \leq i, j \leq n-1$ we have*

- if $i \neq j$ then

$$s_i \cdot \sigma_{s_j} = \sigma_{s_j},$$

- if $i = j$ then

$$s_j \cdot \sigma_{s_j} = \begin{cases} \alpha_j \sigma_e - \sigma_{s_j} + \sigma_{s_{j+1}} & \text{if } j = 1, \\ \alpha_j \sigma_e - \sigma_{s_j} + \sigma_{s_{j-1}} + \sigma_{s_{j+1}} & \text{if } j = 2, 3, \dots, n-2, \\ \alpha_j \sigma_e - \sigma_{s_j} + \sigma_{s_{j-1}} & \text{if } j = n-1, \end{cases}$$

- and for all $w \in S_n$

$$w \sigma_e = \sigma_e.$$

Our choice of S^1 induces the linear projection $\mathfrak{t}^* \rightarrow \text{Lie}(S^1)^*$ which sends the simple roots α_i to t , where t denotes the polynomial variable in $\text{Sym}(\text{Lie}(S^1)^*)$. The following corollary is immediate from this observation together with the formulae in Proposition 6.8.

Corollary 6.9. *The Kostant-Kumar action of S_n on $H_{S^1}^*(GL_n(\mathbb{C})/B; \mathbb{C})$ preserves the $H_{S^1}^*(\text{pt}; \mathbb{C})$ -submodule that is spanned by the images of the classes $\{\sigma_e, \sigma_{s_1}, \sigma_{s_2}, \dots, \sigma_{s_{n-1}}\}$.*

Proof. By definition, Kostant-Kumar's action of each $w \in S_n$ is an $H_T^*(\text{pt}; \mathbb{C})$ -module homomorphism, in the sense that if $f \in \text{Sym}(\mathfrak{t}^*)$ and $\sigma \in H_T^*(GL_n(\mathbb{C})/B; \mathbb{C})$ then $w \cdot (f\sigma) = (f)(w \cdot \sigma)$. Proposition 6.8 thus implies that the $H_T^*(\text{pt}; \mathbb{C})$ -span of $\{\sigma_e, \sigma_{s_1}, \sigma_{s_2}, \dots, \sigma_{s_{n-1}}\}$ is an S_n -subrepresentation of $H_T^*(GL_n(\mathbb{C})/B; \mathbb{C})$. The ring homomorphism $H_T^*(\text{pt}; \mathbb{C}) \rightarrow H_{S^1}^*(\text{pt}; \mathbb{C})$ is a surjection and the additive homomorphism $H_T^*(GL_n(\mathbb{C})/B; \mathbb{C}) \rightarrow H_{S^1}^*(GL_n(\mathbb{C})/B; \mathbb{C})$ respects multiplication in the sense of Equation (2.8). Hence the images of the classes $\{\sigma_e, \sigma_{s_1}, \sigma_{s_2}, \dots, \sigma_{s_{n-1}}\}$ span an $H_{S^1}^*(\text{pt}; \mathbb{C})[S_n]$ -submodule of $H_{S^1}^*(GL_n(\mathbb{C})/B; \mathbb{C})$. \square

As a consequence, we deduce that the formulae given in the following corollary give well-defined actions on the ordinary and equivariant cohomology rings of the subregular Springer varieties. For each $w \in S_n$ we denote by p_w the image in $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ of σ_w under the left vertical arrow of (6.3).

Corollary 6.10. *Let N be a subregular nilpotent linear operator $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ and let \mathcal{S}_N denote its associated subregular Springer variety. Kostant-Kumar's S_n -action on $H_T^*(G/B; \mathbb{C})$, described in Proposition 6.8, naturally induces an S_n -representation on $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ as follows. For each i, j with $1 \leq i, j \leq n-1$ define:*

- if $i \neq j$ then

$$s_i \cdot p_{s_j} = p_{s_j},$$

- if $i = j$ then

$$s_j \cdot p_{s_j} = \begin{cases} t p_e - p_{s_j} + p_{s_{j+1}} & \text{if } j = 1, \\ t p_e - p_{s_j} + p_{s_{j-1}} + p_{s_{j+1}} & \text{if } j = 2, 3, \dots, n-2, \\ t p_e - p_{s_j} + p_{s_{j-1}} & \text{if } j = n-1, \end{cases}$$

- for all $w \in S_n$

$$w \cdot p_e = p_e.$$

This is a well-defined S_n -action on $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$. Moreover, this action induces a well-defined S_n -representation on the ordinary cohomology

$$H^*(\mathcal{S}_N; \mathbb{C}) \cong H_{S^1}^*(\mathcal{S}_N; \mathbb{C}) / (t)H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$$

by setting $t = 0$ in the previous formulae.

Proof. From Theorem 6.7 we know that $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ is a free $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module with basis $\{p_e, p_{s_1}, \dots, p_{s_{n-1}}\}$. In addition $H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$ is a free $H_T^*(\text{pt}; \mathbb{C})$ -module with module basis given by the equivariant Schubert classes $\{\sigma_w\}_{w \in S_n}$. By Proposition 6.8, Kostant-Kumar's action preserves the $H_T^*(\text{pt}; \mathbb{C})$ -submodule of $H_T^*(GL(n, \mathbb{C})/B; \mathbb{C})$ generated by the degree-0 and degree-2 classes $\{\sigma_e, \sigma_{s_1}, \dots, \sigma_{s_{n-1}}\}$. By definition of the classes $\{p_w\}$, this submodule maps isomorphically onto $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ under the natural map

$$H_T^*(GL(n, \mathbb{C})/B; \mathbb{C}) \rightarrow H_{S^1}^*(\mathcal{S}_N; \mathbb{C}).$$

The action of S_n on $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ is defined via this isomorphism. The explicit formulas to be proven follow immediately from Proposition 6.8 and the definition of the classes p_w .

By Proposition 6.2 the S^1 -equivariant cohomology of \mathcal{S}_N is a free $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module. Let $M = H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ denote the S^1 -equivariant cohomology of the Springer variety \mathcal{S}_N considered as an $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module and let t be the degree 2 polynomial variable in $H_{S^1}^*(\text{pt}; \mathbb{C}) \cong \mathbb{C}[t]$. The ordinary cohomology of \mathcal{S}_N is isomorphic to the quotient $M/(t)M$ [11, Equation 1.2.4]. The $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module structure on the quotient factors through \mathbb{C} via the ring homomorphism $H_{S^1}^*(\text{pt}; \mathbb{C}) \cong \mathbb{C}[t] \rightarrow \mathbb{C}$ taking t to 0. In particular the images of the $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module basis $\{p_e, p_{s_1}, \dots, p_{s_{n-1}}\}$ in $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ form a \mathbb{C} -module basis for $H^*(\mathcal{S}_N; \mathbb{C})$. The S_n -action defined on the free $H_{S^1}^*(\text{pt}; \mathbb{C})$ -module $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ is $H_{S^1}^*(\text{pt}; \mathbb{C})$ -linear and thus, via the quotient map taking t to 0, yields a well-defined action on the free \mathbb{C} -module $H^*(\mathcal{S}_N; \mathbb{C})$ as desired. \square

We refer to the S_n -actions on both $H_{S^1}^*(\mathcal{S}_N; \mathbb{C})$ and $H^*(\mathcal{S}_N; \mathbb{C})$ as **Kostant-Kumar representations**. We now compute the character of the Kostant-Kumar representation on the complex vector spaces $H^{2i}(\mathcal{S}_N; \mathbb{C})$, denoted by ψ^i . We then compare the Kostant-Kumar representation with the Springer representation.

Proposition 6.11. *Let N be a subregular nilpotent linear operator $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ with Springer variety \mathcal{S}_N . Let $\psi^i : W \rightarrow \mathbb{Z}$ denote the character of the Kostant-Kumar representation on $H^{2i}(\mathcal{S}_N; \mathbb{C})$. Then for each $w \in S_n$*

$$\psi^1(w) = \#\{\text{fixed points of } w\} - 1 = \#\{j \in \{1, 2, \dots, n\} : w(j) = j\} - 1.$$

The S_n -representation on $H^0(\mathcal{S}_N; \mathbb{C})$ is the trivial representation, hence for each $w \in S_n$

$$\psi^0(w) = 1.$$

Proof. For the purposes of this proof we use cycle notation for permutations, so e.g. $(1, 2, 3, 4)$ sends 1 to 2, 2 to 3, 3 to 4, and 4 to 1. Each element of S_n may be written as a product of disjoint cycles, where the product is denoted by concatenation. The character is a class function, so it suffices to compute $\psi^1(w)$ on a representative of each conjugacy class. Thus we may assume without loss of generality that w has the form

$$(6.6) \quad w = (1, 2, \dots, \mu_1)(\mu_1 + 1, \mu_1 + 2, \dots, \mu_2) \cdots (\mu_{j-1} + 1, \mu_{j-1} + 2, \dots, \mu_j = n),$$

for some $\mu_1 < \mu_2 < \cdots < \mu_j$ where cycles may have length 1.

Choose a, b with $1 \leq a < b \leq n$. A cycle $(a, a+1, \dots, b-1, b)$ of length at least 2 has reduced word decomposition $s_a s_{a+1} \cdots s_{b-1}$. Using this word and the formula in Corollary 6.10 we easily check that

$$(6.7) \quad (a, a+1, \dots, b-1, b) \cdot p_k = \begin{cases} -p_k + (\mathbb{Z}\text{-linear combination of } \{p_j\}_{j \neq k}), & \text{if } k = a, \\ \mathbb{Z}\text{-linear combination of } \{p_j\}_{j \neq k}, & \text{if } a+1 \leq k \leq b-1, \\ p_k + (\mathbb{Z}\text{-linear combination of } \{p_j\}_{j \neq k}), & \text{else.} \end{cases}$$

For any $a = 1, 2, \dots, n$, a cycle (a) of length 1 corresponds to a fixed point of w . The cycle (a) also denotes the identity element in S_n so

$$(6.8) \quad (a) \cdot p_k = p_k \text{ for all } k = 1, 2, \dots, n-1.$$

For each $k = 1, 2, \dots, n-1$ consider the basis element p_k . The index k appears in precisely one of the cycles in Equation 6.6. From Equations (6.7) and (6.8) we conclude that, as desired,

$$\psi^1(w) = (\text{number of cycles of length 1}) - 1 = \#\{j \in \{1, 2, \dots, n\} : w(j) = j\} - 1.$$

Finally, the class p_e generates $H^0(\mathcal{S}_N; \mathbb{C})$ and $w p_e = p_e$ for all $w \in S_n$. So $H^0(\mathcal{S}_N; \mathbb{C})$ is the trivial 1-dimensional representation, and $\psi^0(w) = 1$ for all $w \in S_n$. \square

Finally, we observe that the Kostant-Kumar S_n -representation on $H^*(\mathcal{S}_N; \mathbb{C})$ agrees with the Garsia-Procesi description of the Springer representation [9]. In fact, since S_n -representations are uniquely determined by their characters, the following is immediate from Corollary 6.6 and Proposition 6.11.

Corollary 6.12. *Let N be a subregular nilpotent linear operator $N : \mathbb{C}^n \rightarrow \mathbb{C}^n$ and let \mathcal{S}_N denote its associated subregular Springer variety. The S_n -representation on $H_{S_1}^*(\mathcal{S}_N; \mathbb{C})$ constructed in Corollary 6.10 lifts the classical subregular Springer representation on $H^*(\mathcal{S}_N; \mathbb{C})$ to $H_{S_1}^*(\mathcal{S}_N; \mathbb{C})$ via the homomorphism $H_{S_1}^*(\mathcal{S}_N; \mathbb{C}) \rightarrow H^*(\mathcal{S}_N; \mathbb{C})$.*

Remark 6.13. *The extent to which the constructions in Section 6.3 can be generalized to other classes of Springer varieties is an open question. We have preliminary experimental evidence that suggests that module bases constructed via poset pinball for other Springer varieties are not in general poset-upper-triangular with respect to Bruhat order. This may make computations using them more difficult. We leave further exploration to future work.*

We conclude the paper with a concrete example of Betti pinball in the Springer case which does yield a module basis, albeit not a poset-upper-triangular one.

Example 6.14. *In our last example, we play Betti pinball with target Betti numbers $\mathbf{b} = (1, 3, 5, 3)$. For visual simplicity, not all edges in the poset are drawn above rank 2. The reader can verify that the rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{I})$ in this case is a union of principal order ideals. The double lines indicate covering relations in the partial order which cause the failure of poset-upper-triangularity of the rolldown set. Specifically, poset-upper-triangularity fails in this example because of the drop-downs labeled v_8, v_{10}, v_{11} , and v_{12} in the table.*

This example is a pinball game arising from the Springer variety associated to the nilpotent matrix for the partition $(2, 1, 1)$ of $n = 4$. In this case we can check by direct computation that the rolldown set $\mathcal{R}(\mathcal{J}, \mathcal{I})$ yields a module basis for the submodule corresponding to \mathcal{J} , even though it is not poset-upper-triangular.




FIGURE 6.1. An example of Betti pinball for which the rolldown set is not poset-upper-triangular, but is a union of principal order ideals.

(6.9)

<i>pinball step</i>	w_k	v_k
1	$w_1 = e = [1, 2, 3, 4]$	$v_1 = e = [1, 2, 3, 4]$
2	$w_2 = s_3 = [1, 2, 4, 3]$	$v_2 = s_3 = [1, 2, 4, 3]$
3	$w_3 = s_2 = [1, 3, 2, 4]$	$v_3 = s_2 = [1, 3, 2, 4]$
4	$w_4 = s_2 s_3 = [1, 3, 4, 2]$	$v_4 = s_2 s_3 = [1, 3, 4, 2]$
5	$w_5 = s_3 s_2 = [1, 4, 2, 3]$	$v_5 = s_3 s_2 = [1, 4, 2, 3]$
6	$w_6 = s_2 s_1 = [3, 1, 2, 4]$	$v_6 = s_1 = [2, 1, 3, 4]$
7	$w_7 = s_3 s_2 s_3 = [1, 4, 3, 2]$	$v_7 = s_3 s_2 s_3 = [1, 4, 3, 2]$
8	$w_8 = s_2 s_1 s_3 = [3, 1, 4, 2]$	$v_8 = s_1 s_3 = s_3 s_1 = [2, 1, 4, 3]$
9	$w_9 = s_3 s_2 s_1 = [4, 1, 2, 3]$	$v_9 = s_2 s_1 = [3, 1, 2, 4]$
10	$w_{10} = s_2 s_1 s_3 s_2 = [3, 4, 1, 2]$	$v_{10} = s_1 s_2 = [2, 3, 1, 4]$
11	$w_{11} = s_3 s_2 s_1 s_3 = [4, 1, 3, 2]$	$v_{11} = s_3 s_2 s_1 = [4, 1, 2, 3]$
12	$w_{12} = s_3 s_2 s_1 s_3 s_2 = [4, 3, 1, 2]$	$v_{12} = s_1 s_3 s_2 = [2, 4, 1, 3]$

REFERENCES

[1] M. F. Atiyah and R. Bott. The moment map and equivariant cohomology. *Topology*, 23(1):1–28, 1984.

[2] A. Białyński-Birula. Some properties of the decompositions of algebraic varieties determined by actions of a torus. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 24(9):667–674, 1976.

[3] S. Billey. Kostant polynomials and the cohomology ring of G/B . *Duke Math. J.*, 96:205–224, 1999.

[4] N. Bourbaki. *Éléments de mathématique*. Masson, Paris, 1981. Groupes et algèbres de Lie. Chapitres 4, 5 et 6. [Lie groups and Lie algebras. Chapters 4, 5 and 6].

[5] J. B. Carrell. Orbits of the Weyl group and a theorem of DeConcini and Procesi. *Compositio Math.*, 60(1):45–52, 1986.

[6] J. B. Carrell and K. Kaveh. On the equivariant cohomology of subvarieties of a B -regular variety. *Transform. Groups*, 13(3-4):495–505, 2008.

[7] D. H. Collingwood and W. M. McGovern. *Nilpotent orbits in semisimple Lie algebras*. Van Nostrand Reinhold Mathematics Series. Van Nostrand Reinhold Co., New York, 1993.

[8] W. Fulton. *Young tableaux*, volume 35 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.

[9] A. M. Garsia and C. Procesi. On certain graded S_n -modules and the q -Kostka polynomials. *Adv. Math.*, 94(1):82–138, 1992.

[10] R. Goldin and S. Tolman. Towards generalizing Schubert calculus in the symplectic category, April 2009, math.SG/0904.1245.

[11] M. Goresky, R. Kottwitz, and R. MacPherson. Equivariant cohomology, Koszul duality, and the localization theorem. *Invent. Math.*, 131:25–83, 1998.

[12] M. Goresky and R. MacPherson. On the spectrum of the equivariant cohomology ring. *Canad. J. Math.*, 62(2):262–283, 2010.

[13] V. W. Guillemin and S. Sternberg. *Supersymmetry and equivariant de Rham theory*. Mathematics Past and Present. Springer-Verlag, Berlin, 1999.

[14] M. Harada, A. Henriques, and T. S. Holm. Computation of generalized equivariant cohomologies of Kac-Moody flag varieties. *Adv. Math.*, 197(1):198–221, 2005.

[15] M. Harada and T. S. Holm. The equivariant cohomology of hypertoric varieties and their real loci. *Comm. Anal. Geom.*, 13(3):527–559, 2005.

[16] M. Harada and J. Tymoczko. A positive Monk formula in the S^1 -equivariant cohomology of type A peterson varieties, <http://arxiv.org/abs/0908.3517>.

[17] J. E. Humphreys. *Linear algebraic groups*. Springer-Verlag, New York, 1975. Graduate Texts in Mathematics, No. 21.

[18] T. Ikeda and H. Naruse. Excited Young diagrams and equivariant Schubert calculus. *Trans. Amer. Math. Soc.*, 361(10):5193–5221, 2009.

[19] A. Knutson. A compactly supported formula for equivariant localization, and, simplicial complexes of Białyński-Birula decompositions, <http://arxiv.org/abs/0801.4092>.

[20] A. Knutson and T. Tao. Puzzles and (equivariant) cohomology of grassmannians. *Duke Math. J.*, 119(2):221–260, 2003.

[21] B. Kostant and S. Kumar. The nil Hecke ring and cohomology of G/P for a Kac-Moody group G . *Advances in Math.*, 62:187–237, 1986.

[22] V. Kreiman. Schubert classes in the equivariant K -theory and equivariant cohomology of the Grassmannian, <http://arxiv.org/abs/math/0512204>.

[23] S. Kumar. *Kac-Moody groups, their flag varieties and representation theory*, volume 204 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2002.

[24] V. Lakshmibai, K. N. Raghavan, and P. Sankaran. Equivariant Giambelli and determinantal restriction formulas for the Grassmannian. *Pure Appl. Math. Q.*, 2(3, part 1):699–717, 2006.

- [25] J. P. May. *Equivariant homotopy and cohomology theory*, volume 91 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1996.
- [26] B. E. Sagan. *The symmetric group*. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA, 1991. Representations, combinatorial algorithms, and symmetric functions.
- [27] P. Slodowy. *Four lectures on simple groups and singularities*, volume 11 of *Communications of the Mathematical Institute, Rijksuniversiteit Utrecht*. Rijksuniversiteit Utrecht Mathematical Institute, Utrecht, 1980.
- [28] N. Spaltenstein. The fixed point set of a unipotent transformation on the flag manifold. *Nederl. Akad. Wetensch. Proc. Ser. A 79=Indag. Math.*, 38(5):452–456, 1976.
- [29] T. A. Springer. Trigonometric sums, Green functions of finite groups and representations of Weyl groups. *Invent. Math.*, 36:173–207, 1976.
- [30] R. P. Stanley. *Enumerative combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.
- [31] J. S. Tymoczko. Linear conditions imposed on flag varieties. *Amer. J. Math.*, 128(6):1587–1604, 2006.
- [32] J. S. Tymoczko. Paving Hessenberg varieties by affines. *Selecta Math. (N.S.)*, 13(2):353–367, 2007.
- [33] J. S. Tymoczko. Permutation actions on equivariant cohomology of flag varieties. In *Toric topology*, volume 460 of *Contemp. Math.*, pages 365–384. Amer. Math. Soc., Providence, RI, 2008.
- [34] M. Willems. Cohomologie et K -théorie équivariantes des variétés de Bott-Samelson et des variétés de drapeaux. *Bull. Soc. Math. France*, 132(4):569–589, 2004.
- [35] M. Willems. Cohomologie équivariante des tours de Bott et calcul de Schubert équivariant. *J. Inst. Math. Jussieu*, 5(1):125–159, 2006.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, 1280 MAIN STREET WEST, HAMILTON, ONTARIO L8S4K1, CANADA

E-mail address: Megumi.Harada@math.mcmaster.ca

URL: <http://www.math.mcmaster.ca/Megumi.Harada/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, 14 MACLEAN HALL, IOWA CITY, IOWA 52242-1419, U.S.A.

E-mail address: tymoczko@uiowa.edu

URL: <http://www.math.uiowa.edu/~tymoczko/>