

5-2018

Constant symplectic 2-groupoids

Rajan Amit Mehta

Xiang Tang

Follow this and additional works at: https://scholarworks.smith.edu/mth_facpubs

 Part of the [Mathematics Commons](#)

CONSTANT SYMPLECTIC 2-GROUPOIDS

RAJAN AMIT MEHTA AND XIANG TANG

ABSTRACT. We propose a definition of symplectic 2-groupoid which includes integrations of Courant algebroids that have been recently constructed. We study in detail the simple but illustrative case of constant symplectic 2-groupoids. We show that the constant symplectic 2-groupoids are, up to equivalence, in one-to-one correspondence with a simple class of Courant algebroids that we call constant Courant algebroids. Furthermore, we find a correspondence between certain Dirac structures and Lagrangian sub-2-groupoids.

1. INTRODUCTION

In [10], Liu, Weinstein, and Xu introduced the notion of a *Courant algebroid*, axiomatizing the brackets studied by Courant and Weinstein [2, 3] and Dorfman [4]. Recognizing the similarities and relationships between Lie algebroids and Courant algebroids, they asked whether there is a groupoid-like object that can be viewed as the “integration” of a Courant algebroid.

In [15], Ševera outlined a construction by which, given a Courant algebroid, one could formally produce a 2-groupoid as a moduli space of maps of dg-manifolds. The space of 2-simplices of this 2-groupoid possesses a symplectic form, suggesting that the integration of a Courant algebroid should be called a *symplectic 2-groupoid*. In the case of an exact Courant algebroid, the symplectic 2-groupoid arising from Ševera’s construction (the *Liu-Weinstein-Xu 2-groupoid*) was explicitly described and shown to be smooth in [11].

In all but the most trivial cases, the 2-groupoids arising from Ševera’s construction are infinite-dimensional, so it is reasonable to look for finite-dimensional models. In different (but overlapping) special cases, such models were independently found in [9, 12, 16], each with its own shortcoming. In [16], there is no 2-form constructed on the integration. In [9], the integration has a symplectic structure but is only defined locally¹. The construction in [12] is global, but the 2-form constructed there is degenerate.

Based on the class of examples discovered there, the authors in [12] suggested a definition of symplectic 2-groupoid where the 2-form is allowed to be degenerate, but where the degeneracy is controlled in a certain way by the simplicial structure. However, the definition given there fails to serve as a good general definition. Specifically, [12, Definition 6.7] implies that when the 2-form is genuinely nondegenerate at a point x of the unit space X_0 , the tangent space of X_1 at x is isomorphic to the tangent space of X_0 at x , which forces X_1 to have the same dimension of X_0 . This excludes too many interesting examples, including the case of constant

2010 *Mathematics Subject Classification.* 53D17, 58H05.

Key words and phrases. Dirac, symplectic, 2-groupoids, Courant algebroids.

¹Recently, Ševera and Širaň [17] described a general construction for local integration and showed that the local integrations can be glued together up to coherent homotopy.

symplectic 2-groupoids to be presented in this paper. This observation suggests that the definition requires revision.

In Definition 2.7, we give a new definition of symplectic 2-groupoid which still includes the examples constructed in [12] (see Section 2.4). This definition is partly inspired by the notion of a shifted symplectic structure [13], and we are particularly indebted to Getzler [6], who recast many of those ideas in the concrete language of simplicial manifolds. However, we stress that our definition is in several ways more strict than the one coming from [13]. Specifically, our definition only includes a 2-form on the space of 2-simplices X_2 , and we require this 2-form to be closed on the nose. Additionally, our nondegeneracy requirement is more strict.

The strictness of our definition reflects the fact that the solution to the integration problem will not be Morita invariant. For example, the notion of symplectic groupoid [1] is not Morita invariant, and this strictness is necessary in order to obtain a Lie-theoretic correspondence with Poisson manifolds. On the other hand, the notion of 1-shifted symplectic structure agrees with Xu’s [18] weaker notion of *quasi-symplectic groupoid*.

In the second half of this article, we consider the case of *constant symplectic 2-groupoids*, i.e. symplectic 2-groupoids with a linear structure with respect to which the 2-form is constant. Besides being a basic test case for the definition, constant symplectic 2-groupoids should be useful for understanding the general case since they appear as first-order approximations of arbitrary symplectic 2-groupoids. Specifically, given a symplectic 2-groupoid (X_\bullet, ω) and a point $x \in X_0$, the “tangent space at x ” $(T|_x X_\bullet, \omega_x)$ is a constant symplectic 2-groupoid. In this sense, constant symplectic 2-groupoids play the same role in the study of symplectic 2-groupoids as symplectic vector spaces play in the study of symplectic manifolds.

We find that there is a relationship between constant symplectic 2-groupoids and a certain class of Courant algebroids that we call *constant Courant algebroids*. Specifically, the main results are as follows:

- (1) (Theorem 4.3) There is a one-to-one correspondence between constant Courant algebroids and equivalence classes of constant symplectic 2-groupoids.
- (2) (Theorem 5.6) Under the above correspondence, constant Dirac structures are in one-to-one correspondence with wide linear Lagrangian sub-2-groupoids.

In other words, constant Courant algebroids integrate to constant symplectic 2-groupoids, and constant Dirac structures integrate to certain Lagrangian sub-2-groupoids. These results provide evidence in support of our definition of symplectic 2-groupoids as being the correct answer to the question posed by Liu, Weinstein, and Xu.

Organization of the paper. In Section 2, we define symplectic 2-groupoids and show that the class of examples from [12] satisfies the definition. In Section 3, we study constant symplectic 2-groupoids and find a minimal description of them in terms of linear algebra data. In Section 4, we similarly study constant Courant algebroids and show that they are in correspondence with constant symplectic 2-groupoids. Finally, in Section 5, we consider linear Lagrangian sub-2-groupoids and describe the correspondence with constant Dirac structures.

Acknowledgments. We would like to thank Ezra Getzler for inspiring discussions and explanations about symplectic structures on differentiable n -stacks. We would

like to thank Damien Calaque for a discussion about the relation between integration of Courant algebroids and derived symplectic geometry. The research of the second author is partially supported by NSF grant DMS 1363250.

2. SYMPLECTIC 2-GROUPOIDS

2.1. Lie n -groupoids and differential forms. We start by recalling the definition of a Lie n -groupoid (see [5, 8, 19]).

Definition 2.1. A *simplicial manifold* is a sequence $X_\bullet = \{X_q\}$, $q \geq 0$, of manifolds equipped with surjective submersions $f_i^q : X_q \rightarrow X_{q-1}$ (called *face maps*), $i = 0, \dots, q$, and embeddings $\sigma_i^q : X_q \rightarrow X_{q+1}$ (called *degeneracy maps*), $i = 0, \dots, q$, such that

$$(2.1) \quad f_i^{q-1} f_j^q = f_{j-1}^{q-1} f_i^q, \quad i < j,$$

$$(2.2) \quad \sigma_i^{q+1} \sigma_j^q = \sigma_{j+1}^{q+1} \sigma_i^q, \quad i < j,$$

$$(2.3) \quad f_i^{q+1} \sigma_j^q = \begin{cases} \sigma_{j-1}^{q-1} f_i^q, & i < j, \\ \text{id}, & i = j, j+1, \\ \sigma_j^{q-1} f_{i-1}^q, & i > j+1. \end{cases}$$

For $q \geq 1$ and $0 \leq k \leq q$, recall that a (q, k) -*horn* of X_\bullet consists of a q -tuple $(x_0, \dots, x_{k-1}, x_{k+1}, \dots, x_q)$, where $x_i \in X_{q-1}$, satisfying the *horn compatibility equations*

$$(2.4) \quad f_i^{q-1} x_j = f_{j-1}^{q-1} x_i$$

for $i < j$. The space of all (q, k) -horns is denoted $\Lambda_{q,k} X$.

The natural *horn maps* $\lambda_{q,k} : X_q \rightarrow \Lambda_{q,k} X$ are defined as

$$\lambda_{q,k}(x) = (f_0^q x, \dots, \widehat{f_k^q x}, \dots, f_q^q x)$$

for $x \in X_q$. It is immediate from (2.1) that $\lambda_{q,k}(x)$ satisfies the horn compatibility equations (2.4); in fact, the purpose of the horn compatibility equations is to axiomatize the properties satisfied by $\lambda_{q,k}(x)$.

Definition 2.2. A *Lie n -groupoid* is a simplicial manifold such that the horn maps $\lambda_{q,k}$ are

- (1) surjective submersions for all $q \geq 1$, and
- (2) diffeomorphisms for all $q > n$.

Given a simplicial manifold X_\bullet , we consider the bigraded space of differential forms $\Omega^\bullet(X_\bullet)$. There are two natural commuting differentials on $\Omega^\bullet(X_\bullet)$. One is the de Rham differential $d : \Omega^p(X_q) \rightarrow \Omega^{p+1}(X_q)$, and the other is the simplicial coboundary operator $\delta : \Omega^p(X_q) \rightarrow \Omega^p(X_{q+1})$, given by $\delta\alpha = \sum_{i=0}^{q+1} (-1)^i (f_i^{q+1})^* \alpha$.

Definition 2.3. A form $\alpha \in \Omega^p(X_q)$ is *multiplicative* if $\delta\alpha = 0$. A form $\alpha \in \Omega^p(X_q)$ is *normalized* if $(\sigma_{q-1}^i)^* \alpha = 0$ for all $i = 0, \dots, q-1$.

We note that the normalization condition holds vacuously in the case $q = 0$.

The space of normalized forms is denoted $\Omega_\nu^\bullet(X_\bullet)$. It is not hard to check that $\Omega_\nu^\bullet(X_\bullet)$ is closed under d and δ .

2.2. The tangent complex. Let X_\bullet be a simplicial manifold. For each $q > 0$, let $\sigma^q : X_0 \rightarrow X_q$ be defined as $\sigma^q = \sigma_0^{q-1} \cdots \sigma_0^0$. We can think of the image of σ^q as being the “unit space” in X_q .

For $x \in X_0$ let $T_{x,q}X$ denote the tangent space $T_{\sigma^q(x)}X_q$. There is a natural simplicial structure on $T_{x,\bullet}X$ where the face and degeneracy maps are restrictions of the differentials of f_i^q and σ_i^q . We note that $T_{x,\bullet}X$ is a simplicial vector space, in the sense that each $T_{x,q}X$ is a vector space and all of the face and degeneracy maps are linear.

There is a natural boundary map $\partial_q : T_{x,q}X \rightarrow T_{x,q-1}X$, given by

$$\partial_q(v) = \sum_{i=0}^q (-1)^i (f_i^q)_* v.$$

It follows from (2.1) that $\partial^2 = 0$.

The Dold-Kan correspondence, c.f. [7], associates a chain complex $\hat{T}_{x,\bullet}X$ to the simplicial vector space $T_{x,\bullet}X$. This chain complex can be explicitly described as follows. The *normalized tangent space* $\hat{T}_{x,q}X$ is defined to be the quotient of $T_{x,q}X$ by the sum of the degenerate subspaces:

$$\hat{T}_{x,q}X := T_{x,q}X \Big/ \sum_{i=0}^{q-1} (\sigma_i^{q-1})_* T_{x,q-1}X.$$

In particular, $\hat{T}_{x,0} = T_{x,0}$.

The boundary map ∂ descends to the normalized tangent spaces, so we have a chain complex

$$\cdots \rightarrow \hat{T}_{x,q}X \xrightarrow{\partial} \hat{T}_{x,q-1}X \xrightarrow{\partial} \cdots \xrightarrow{\partial} \hat{T}_{x,1}X \xrightarrow{\partial} \hat{T}_{x,0}X,$$

which is called the *tangent complex* of the simplicial manifold X_\bullet at $x \in X_0$. Taken together, the tangent complexes at every $x \in X_0$ form a complex of vector bundles over X_0 , called the tangent complex of X_\bullet .

Proposition 2.4. *If X_\bullet is a Lie n -groupoid, then $\hat{T}_{x,q}$ is trivial for $q > n$.*

In this paper, we will only make use of Proposition 2.4 in the case $n = 2$. A proof in this case is essentially contained in Section 3.1; in particular, see Remark 3.1. We leave the general case to the reader.

Example 2.5. In the case where X_\bullet is the nerve of a Lie groupoid $G \rightrightarrows M$, then the tangent complex can be identified with the 2-term complex $A \xrightarrow{\rho} TM$, where A is the Lie algebroid of G and ρ is the anchor map.

2.3. Forms on Lie 2-groupoids. We will now restrict our attention to the case of Lie 2-groupoids. From Proposition 2.4, we know that the tangent complex of a Lie 2-groupoid X_\bullet is a 3-term complex

$$\hat{T}_2X \xrightarrow{\partial} \hat{T}_1X \xrightarrow{\partial} T_0X.$$

Given a normalized 2-form $\omega \in \Omega_v^2(X_2)$, we can obtain the following bilinear pairings on the tangent complex at any $x \in X_0$ (we learned of these from Getzler [6]):

(1) For $v \in T_{x,0}X$ and $w \in T_{x,2}X$, let

$$(2.5) \quad \tilde{A}_\omega(v, w) := \omega(\sigma_*^2 v, w).$$

It follows from the assumption that ω is normalized that \tilde{A}_ω descends to a well-defined bilinear pairing A_ω between $T_{x,0}X$ and $\hat{T}_{x,2}X$.

(2) For $\theta, \eta \in T_{x,1}X$, let

$$(2.6) \quad \tilde{B}_\omega(\theta, \eta) := \omega((\sigma_1^1)_*\theta, (\sigma_0^1)_*\eta) + \omega((\sigma_1^1)_*\eta, (\sigma_0^1)_*\theta).$$

It follows from the assumption that ω is normalized that \tilde{B}_ω descends to a well-defined symmetric bilinear form B_ω on $\hat{T}_{x,1}X$.

Definition 2.6. A normalized 2-form $\omega \in \Omega_\nu^2(X_2)$ is called *simplicially nondegenerate* if the induced pairings A_ω and B_ω are nondegenerate at all $x \in X_0$.

Definition 2.7. A *symplectic 2-groupoid* is a Lie 2-groupoid X_\bullet equipped with a closed, multiplicative, normalized, and simplicially nondegenerate 2-form $\omega \in \Omega_\nu^2(X_2)$.

Definition 2.8. Two closed, multiplicative, normalized, and simplicially nondegenerate 2-forms $\omega, \omega' \in \Omega_\nu^2(X_2)$ are *equivalent* if there exists a closed and normalized 2-form $\alpha \in \Omega_\nu^2(X_1)$ such that $\omega' - \omega = \delta\alpha$ and $A_{\delta\alpha} = B_{\delta\alpha} = 0$.

Remark 2.9. By a straightforward calculation using (2.5) and (2.3), one can see that $A_{\delta\alpha} = 0$ if and only if

$$(2.7) \quad \alpha((\sigma_0^0)_*v, \partial w) = 0$$

for all $v \in T_{x,0}X$ and $w \in T_{x,2}X$. Similarly, $B_{\delta\alpha} = 0$ if and only if

$$(2.8) \quad \alpha(\theta, (\sigma_0^0)_*\partial\eta) + \alpha(\eta, (\sigma_0^0)_*\partial\theta) = 0$$

for all $\theta, \eta \in T_{x,1}X$.

Remark 2.10. It is known [1] that any multiplicative 2-form on a Lie groupoid is automatically normalized; thus the normalization condition does not explicitly appear in the definition of symplectic groupoid. However, in the case of Lie 2-groupoids, normalization does not automatically follow from multiplicativity. For example, for any manifold M , let $X_k = M$ for all k , with all the face and degeneracy maps being the identity. It is immediate that any nonzero 2-form ω on $X_2 = M$ is multiplicative but not normalized.

2.4. Example: an integration of $A \oplus A^*$. In [10], Liu, Weinstein, and Xu constructed a Courant algebroid $A \oplus A^*$ associated to any Lie bialgebroid (A, A^*) . This construction leads to a large and important class of Courant algebroids. In [12], the authors described a method of integrating Courant algebroids of the form $A \oplus A^*$ by first integrating (if possible) the Lie bialgebroid (A, A^*) to a symplectic double Lie groupoid D and then applying the bar functor to obtain a Lie 2-groupoid $\overline{W}ND$ equipped with a closed 2-form. The fact that this 2-form is degenerate was the first clue that the correct notion of symplectic 2-groupoid should allow for 2-forms that have some degeneracy.

We will now prove that the integration of the standard Courant algebroid² $TM \oplus T^*M$ in [12] satisfies Definition 2.7.

²The extension of the proof to the general case $\overline{W}ND$, where D is a symplectic double Lie groupoid, is similar.

We first recall the Lie 2-groupoid obtained in [12] as integration of $TM \oplus T^*M$. For any $k \geq 1$, let $T_{(k)}^*M$ denote the direct sum of k copies of T^*M . For each $q \geq 0$, define X_q as

$$X_q = M \times T^*M \times \cdots \times T_{(q)}^*M.$$

Let $\tau_i^q : T_{(q)}^*M \rightarrow T^*M$ be the projection onto the i -th component, let $p_q : T_{(q)}^*M \rightarrow M$ denote the bundle projection map, and let $\iota^q : M \rightarrow T_{(q)}^*M$ be the zero section map. We will omit the index q in p_q and ι^q when $q = 1$. Finally, for $i = 1, 2$, let $\iota_i^2 : T^*M \rightarrow T_{(2)}^*M$ be defined as

$$\iota_1^2(\xi) = (\xi, \iota(p(\xi))), \quad \iota_2^2(\xi) = (\iota(p(\xi)), \xi).$$

The face and degeneracy maps between $X_0 = M$ and $X_1 = M \times T^*M$ are defined as

$$\sigma_0^0(x) = (x, \iota(x)), \quad f_0^1(x, \xi) = x, \quad f_1^1(x, \xi) = p(\xi),$$

for $x \in M$ and $\xi \in T^*M$. The degeneracy maps $\sigma_i^1 : X_1 \rightarrow X_2$ are defined as

$$\sigma_0^1(x, \xi) = (x, \xi, \iota_1^2(\xi)), \quad \sigma_1^1(x, \xi) = (x, \iota(x), \iota_2^2(\xi)),$$

and the face maps $f_i^2 : X_2 \rightarrow X_1$ are defined as

$$\begin{aligned} f_0^2(x, \xi, \xi^2) &:= (x, \xi), \\ f_1^2(x, \xi, \xi^2) &:= (x, \tau_1^2(\xi^2) + \tau_2^2(\xi^2)), \\ f_2^2(x, \xi, \xi^2) &:= (p(\xi), \tau_2^2(\xi^2)), \end{aligned}$$

for $x \in M$, $\xi \in T^*M$, and $\xi^2 \in T_{(2)}^*M$.

As we will only need the structure maps up to level 2, we stop here and refer the reader to [12] for the general definitions of the simplicial structure maps.

There is a natural map $d : X_2 = M \times T^*M \times T_{(2)}^*M \rightarrow T^*M \times T^*M$ given by

$$d(m, \xi, \xi^2) := (\xi, \tau_1^2(\xi^2)).$$

Let ω_0 be the canonical symplectic form on T^*M . Then the 2-form $\omega \in \Omega^2(X_2)$ is defined as the pullback by d of $(\omega_0, -\omega_0)$. It is clear that ω is closed, we proved in [12, Proposition 6.2] that ω is multiplicative, and it can be easily checked that ω is normalized.

Proposition 2.11. *The 2-form ω on $X_2 = M \times T^*M \times T_{(2)}^*M$ is simplicially nondegenerate. Therefore, (X_\bullet, ω) is a symplectic 2-groupoid.*

Proof. We start by describing the tangent spaces $T_{x,q}X$. We observe that $T_{x,0}X = T_xM$, that $T_{x,1}X$ can be identified with $T_xM \oplus T_xM \oplus T_x^*M$, and that $T_{x,2}X$ can be similarly identified with $T_xM \oplus T_xM \oplus T_x^*M \oplus T_xM \oplus T_x^*M \oplus T_x^*M$. Using these identifications, we can describe the degeneracy maps $(\sigma_0^0)_* : T_{x,0}X \rightarrow T_{x,1}X$ and $(\sigma_i^1)_* : T_{x,1}X \rightarrow T_{x,2}X$ as

$$(\sigma_0^0)_*(v) = (v, v, 0),$$

$$(\sigma_0^1)_*(v_1, v_2, \xi) = (v_1, v_2, \xi, v_2, \xi, 0), \quad (\sigma_1^1)_*(v_1, v_2, \xi) = (v_1, v_1, 0, v_2, 0, \xi),$$

for $v, v_1, v_2 \in T_xM$ and $\xi \in T_x^*M$.

At $\sigma^2(x)$, the 2-form ω is given in terms of the above identifications by

$$\omega\left((v_1, v_2, \xi_1, v_3, \xi_2, \xi_3), (v'_1, v'_2, \xi'_1, v'_3, \xi'_2, \xi'_3)\right) = \xi'_1(v_2) - \xi_1(v'_2) - \xi'_2(v_3) + \xi_2(v'_3).$$

From this, we can compute the pairing \tilde{A}_ω between $T_{x,0}X$ and $T_{x,2}X$ to be

$$\tilde{A}_\omega\left(v, (v_1, v_2, \xi_1, v_3, \xi_2, \xi_3)\right) := \xi_1(v) - \xi_2(v)$$

and the pairing \tilde{B}_ω on $T_{x,1}$ to be

$$\tilde{B}_\omega\left((v_1, v_2, \xi), (v'_1, v'_2, \xi')\right) := \xi'(v_1 - v_2) + \xi(v'_1 - v'_2).$$

One can now see that the kernels of \tilde{A}_ω and \tilde{B}_ω consist of sums of vectors in the images of $(\sigma_i^q)_*$. Thus the induced pairing A_ω between $T_{x,0}X$ and $\hat{T}_{x,2}X$ and the induced bilinear form B_ω on $\hat{T}_{x,1}X$ are both nondegenerate. \square

Remark 2.12. The fact that $\hat{T}_{x,2}X$ pairs nondegenerately with $T_{x,0}X = T_xM$ means that $\hat{T}_{x,2}X$ is isomorphic to T_x^*M . The isomorphism is explicitly given by composing the map $T_x^*M \rightarrow T_{x,2}X$, $\xi \mapsto (0, 0, \xi, 0, 0, 0)$ with the quotient map $T_{x,2}X \rightarrow \hat{T}_{x,2}X$.

Similarly, $\hat{T}_{x,1}X$ is isomorphic to $T_xM \oplus T_x^*M$, with the isomorphism given by the map $T_xM \oplus T_x^*M \rightarrow T_{x,1}X$, $(v, \xi) \mapsto (v, 0, \xi)$. Under this isomorphism, the pairing B_ω agrees with the standard symmetric pairing on $TM \oplus T^*M$, which is an important part of the Courant algebroid structure.

3. CONSTANT SYMPLECTIC 2-GROUPOIDS

A *constant symplectic 2-groupoid* is a symplectic 2-groupoid (V_\bullet, ω) where V_\bullet is a simplicial vector space and $\omega \in \Omega_\nu^2(V_2)$ is constant. In this section, we will study constant symplectic 2-groupoids and obtain a fairly simple description of them. We will later see that there is a correspondence between constant symplectic 2-groupoids and a certain class of Courant algebroids that we call *constant Courant algebroids*.

3.1. Linear 2-groupoids. A *linear 2-groupoid* is a Lie 2-groupoid V_\bullet such that each V_q is a vector space, and where the face and degeneracy maps are all linear. Linear 2-groupoids are known to be equivalent, via the Dold-Kan correspondence, to 3-term chain complexes of vector spaces. Since there are different possible choices of convention and we will require explicit formulas, we will give a brief description of this correspondence.

Suppose that V_\bullet is a linear 2-groupoid. A 3-term chain complex (W_\bullet, ∂) is constructed as follows.

First, we set $W_0 := V_0$. Next, we observe that $f_0^1 : V_1 \rightarrow W_0$ is a surjection with right inverse σ_0^0 . Thus we have a split short exact sequence

$$(3.1) \quad W_1 \longrightarrow V_1 \begin{array}{c} \xleftarrow{\sigma_0^0} \\ \xrightarrow{f_0^1} \end{array} W_0,$$

where $W_1 := \ker f_0^1$, giving us the natural decomposition $V_1 = W_1 \oplus W_0$. Using this decomposition, we define a linear map $\partial_1 : W_1 \rightarrow W_0$ given by $\partial_1 w_1 = -f_1^1(w_1, 0)$. (The sign is chosen to agree with the tangent complex; see Remark 3.1.) We then have the following formulas for the face and degeneracy maps between V_1 and V_0 :

$$f_0^1(w_1, w_0) = w_0, \quad f_1^1(w_1, w_0) = w_0 - \partial_1 w_1, \quad \sigma_0^0(w_0) = (0, w_0).$$

Now consider the horn space $\Lambda_{2,2}V$, consisting of pairs $(v_1, v'_1) \in V_1 \times V_1$ such that $f_0^1(v_1) = f_0^1(v'_1)$. Given such a pair, we may use the decomposition $V_1 = W_1 \oplus W_0$

to write $v_1 = (w_1, w_0)$, $v'_1 = (w'_1, w_0)$. This allows us to make the identification $\Lambda_{2,2}V = W_1 \oplus W_1 \oplus W_0$, where the pair (v_1, v'_1) is identified with (w_1, w'_1, w_0) .

Let $\Phi_2 : \Lambda_{2,2}V \rightarrow V_2$ be defined by

$$(3.2) \quad \Phi_2(v_1, v'_1) = \sigma_0^1 v_1 + \sigma_1^1 (v'_1 - v_1).$$

One can see that Φ_2 is a right inverse of the horn map $\lambda_{2,2}$, so we have a split short exact sequence

$$(3.3) \quad W_2 \longrightarrow V_2 \begin{array}{c} \xleftarrow{\Phi_2} \\ \xrightarrow{\lambda_{2,2}} \end{array} \Lambda_{2,2}V,$$

where $W_2 := \ker \lambda_{2,2}$, giving us a natural decomposition $V_2 = W_2 \oplus \Lambda_{2,2}V = W_2 \oplus W_1 \oplus W_1 \oplus W_0$. Under this decomposition, we have by construction that

$$\begin{aligned} f_0^2(w_2, w_1, w'_1, w_0) &= (w_1, w_0), \\ f_1^2(w_2, w_1, w'_1, w_0) &= (w'_1, w_0). \end{aligned}$$

Using (3.2), we also see that

$$(3.4) \quad \sigma_0^1(w_1, w_0) = \Phi_2(w_1, w_1, w_0) = (0, w_1, w_1, w_0),$$

$$(3.5) \quad \sigma_1^1(w_1, w_0) = \Phi_2(0, w_1, w_0) = (0, 0, w_1, w_0).$$

To obtain a formula for f_2^2 , we first define a map $\partial_2 : W_2 \rightarrow W_1$, given by

$$(3.6) \quad f_2^2(w_2, 0, 0, 0) = (\partial_2 w_2, 0).$$

Note that (3.6) well-defines ∂_2 , since $f_0^1 f_2^2(w_2, 0, 0, 0) = f_1^1 f_0^2(w_2, 0, 0, 0) = 0$. We then make the following calculation:

$$(3.7) \quad \begin{aligned} f_2^2(0, w_1, w'_1, w_0) &= f_2^2 \Phi_2(w_1, w'_1, w_0) \\ &= f_2^2 \sigma_0^1(w_1, w_0) + f_2^2 \sigma_1^1(w'_1 - w_1, 0) \\ &= \sigma_0^0 f_1^1(w_1, w_0) + (w'_1 - w_1, 0) \\ &= (w'_1 - w_1, w_0 - \partial_1 w_1). \end{aligned}$$

Putting (3.6) and (3.7) together, we have

$$f_2^2(w_2, w_1, w'_1, w_0) = (\partial_2 w_2 + w'_1 - w_1, w_0 - \partial_1 w_1).$$

We observe that $\partial_1 \partial_2 w_2 = -f_1^1 f_2^2(w_2, 0, 0, 0) = -f_1^1 f_1^2(w_2, 0, 0, 0) = 0$, so

$$W_2 \xrightarrow{\partial_2} W_1 \xrightarrow{\partial_1} W_0$$

is a 3-term chain complex.

Now consider the horn space $\Lambda_{3,3}V$, consisting of triples $(v_2, v'_2, v''_2) \in V_2 \times V_2 \times V_2$ such that $f_0^2 v_2 = f_0^2 v'_2$, $f_1^2 v_2 = f_0^2 v''_2$, and $f_1^2 v'_2 = f_1^2 v''_2$. In terms of the decomposition $V_2 = W_2 \oplus W_1 \oplus W_1 \oplus W_0$, we may write $v_2 = (w_2, w_1, w'_1, w_0)$, $v'_2 = (w'_2, w_1, w''_1, w_0)$, $v''_2 = (w''_2, w'_1, w'''_1, w_0)$. This allows us to make the identification $\Lambda_{3,3}V = W_2 \oplus W_2 \oplus W_2 \oplus W_1 \oplus W_1 \oplus W_1 \oplus W_0$, where (v_2, v'_2, v''_2) is identified with $(w_2, w'_2, w''_2, w_1, w'_1, w''_1, w_0)$.

Let $\Phi_3 : \Lambda_{3,3}V \rightarrow V_3$ be given by

$$\Phi_3(v_2, v'_2, v''_2) = \sigma_0^2 v_2 + \sigma_1^2 (v'_2 - v_2) + \sigma_2^2 (v''_2 - v'_2 + v_2 - \sigma_0^1 f_1^2 v_2).$$

One can see that Φ_3 inverts the horn map $\lambda_{3,3}$, which is assumed to be an isomorphism since V_\bullet is a Lie 2-groupoid. Implicitly using Φ_3 to identify V_3 with $\Lambda_{3,3}V$, we then obtain the following formulas for the face maps:

$$(3.8) \quad f_0^3(w_2, w'_2, w''_2, w_1, w'_1, w''_1, w_0) = v_2 = (w_2, w_1, w'_1, w_0),$$

$$(3.9) \quad f_1^3(w_2, w'_2, w''_2, w_1, w'_1, w''_1, w_0) = v'_2 = (w'_2, w_1, w''_1, w_0),$$

$$(3.10) \quad f_2^3(w_2, w'_2, w''_2, w_1, w'_1, w''_1, w_0) = v''_2 = (w''_2, w'_1, w''_1, w_0),$$

$$(3.11) \quad f_3^3(w_2, w'_2, w''_2, w_1, w'_1, w''_1, w_0) = (w''_2 - w'_2 + w_2, \partial_2 w_2 + w'_1 - w_1, \partial_2 w'_2 + w''_1 - w_1, w_0 - \partial_1 w_1).$$

We have seen that, given a linear 2-groupoid V_\bullet , we can obtain a 3-term chain complex (W_\bullet, ∂) . Conversely, given a 3-term chain complex, one can construct a linear 2-groupoid by setting $V_0 = W_0$, $V_1 = W_1 \oplus W_0$, etc., with the face and degeneracy maps given in low degrees by the above formulas. As V_\bullet is a Lie 2-groupoid, the higher simplicial maps are completely determined by the data in low degrees.

Remark 3.1. From the short exact sequences (3.1) and (3.3), we can see that the 3-term chain complex (W_\bullet, ∂) is naturally isomorphic to the tangent complex of V_\bullet . As a result, we obtain an alternative description of the tangent complex of a Lie 2-groupoid X_\bullet , where $\hat{T}_{x,2}X = \ker(\lambda_2^2)_* \subseteq T_{x,2}X$, $\hat{T}_{x,1}X = \ker(\lambda_1^1)_* = \ker(f_0^1)_* \subseteq T_{x,1}X$, and where the boundary map is $\partial_q = (-1)^q (f_q^1)_*$.

3.2. Constant multiplicative 2-forms. Let V_\bullet be a linear 2-groupoid. In this subsection, we will obtain a description of constant multiplicative 2-forms $\omega \in \Omega_\nu^2(V_2)$ in terms of data on the associated 3-term complex (W_\bullet, ∂) .

Recall from Section 3.1 that V_2 can be naturally decomposed as $W_2 \oplus W_1 \oplus W_1 \oplus W_0$. With respect to this decomposition, we may write any constant $\omega \in \Omega^2(V_2)$ as a sum of bilinear forms $C_{11} \in \wedge^2 W_2^*$, $C_{12}, C_{13} \in W_2^* \otimes W_1^*$, $C_{14} \in W_2^* \otimes W_0^*$, $C_{22}, C_{33} \in \wedge^2 W_1^*$, $C_{23} \in W_1^* \otimes W_1^*$, $C_{24}, C_{34} \in W_1^* \otimes W_0^*$, and $C_{44} \in \wedge^2 W_0^*$. Then we can write ω in the form of a block matrix

$$\omega = \begin{bmatrix} C_{11} & C_{12} & C_{13} & C_{14} \\ C_{21} & C_{22} & C_{23} & C_{24} \\ C_{31} & C_{32} & C_{33} & C_{34} \\ C_{41} & C_{42} & C_{43} & C_{44} \end{bmatrix},$$

where $C_{ij}(w, w') = -C_{ji}(w', w)$.

Now suppose that ω is normalized. From (3.5), we immediately see that C_{33} , C_{34} , and C_{44} vanish. From (3.4), we then see that $C_{24} = 0$, since

$$0 = \omega((0, w_1, w_1, w_0), (0, 0, 0, w'_0)) = C_{24}(w_1, w'_0)$$

for all $w_1 \in W_1$ and $w_0, w'_0 \in W_0$, and that

$$0 = \omega((0, w_1, w_1, 0), (0, w'_1, w'_1, 0)) = C_{22}(w_1, w'_1) + C_{23}(w_1, w'_1) + C_{32}(w_1, w'_1)$$

for all $w_1, w'_1 \in W_1$. Thus, ω is of the form

$$(3.12) \quad \omega = \begin{bmatrix} C_{11} & C_{12} & C_{13} & C_{14} \\ C_{21} & C_{22} & C_{23} & 0 \\ C_{31} & C_{32} & 0 & 0 \\ C_{41} & 0 & 0 & 0 \end{bmatrix},$$

where

$$(3.13) \quad C_{22} + C_{23} + C_{32} = 0.$$

Conversely, it is straightforward to check that, if ω is of the form (3.12) and satisfies (3.13), then ω is normalized.

Now suppose that ω is normalized and multiplicative. In the following series of lemmas, we find further relations that hold between the bilinear forms C_{ij} . The proofs utilize the decomposition $V_3 = W_2 \oplus W_2 \oplus W_2 \oplus W_1 \oplus W_1 \oplus W_1 \oplus W_0$ constructed in Section 3.1, as well as the formulas (3.8)–(3.11) for the face maps f_i^3 .

Lemma 3.2. $C_{13}(w_2, w_1) = C_{32}(w_1, \partial w_2)$ for all $w_2 \in W_2$ and $w_1 \in W_1$.

Proof. Let $u, v \in V_3$ be defined as $u = (w_2, 0, 0, 0, 0, 0, 0)$ and $v = (0, 0, 0, 0, 0, w_1, 0)$. Then

$$\begin{aligned} 0 = \delta\omega(u, v) &= \sum_{i=0}^3 (-1)^i \omega(f_i^3 u, f_i^3 v) \\ &= -\omega((w_2, \partial w_2, 0, 0), (0, 0, w_1, 0)) \\ &= -C_{13}(w_2, w_1) - C_{23}(\partial w_2, w_1) \\ &= -C_{13}(w_2, w_1) + C_{32}(w_1, \partial w_2). \quad \square \end{aligned}$$

Lemma 3.3. $C_{12}(w_2, w_1) = C_{32}(\partial w_2, w_1)$ for all $w_2 \in W_2, w_1 \in W_1$.

Proof. Let $u, v \in V_3$ be defined as $u = (0, w_2, 0, 0, 0, 0, 0)$ and $v = (0, 0, 0, 0, w_1, 0, 0)$. Then

$$\begin{aligned} 0 = \delta\omega(u, v) &= -\omega((-w_2, 0, \partial w_2, 0), (0, w_1, 0, 0)) \\ &= C_{12}(w_2, w_1) - C_{32}(\partial w_2, w_1). \quad \square \end{aligned}$$

Lemma 3.4. $C_{11}(w_2, w'_2) = -C_{32}(\partial w_2, \partial w'_2)$ for all $w_2, w'_2 \in W_2$.

Proof. Let $u, v \in V_3$ be defined as $u = (0, 0, w_2, 0, 0, 0, 0)$ and $v = (w'_2, 0, 0, 0, 0, 0, 0)$. Then

$$\begin{aligned} 0 = \delta\omega(u, v) &= -\omega((w_2, 0, 0, 0), (w'_2, \partial w'_2, 0, 0)) \\ &= -C_{11}(w_2, w'_2) - C_{12}(w_2, \partial w'_2). \end{aligned}$$

By Lemma 3.3, we see that $C_{11}(w_2, w'_2) = -C_{12}(w_2, \partial w'_2) = -C_{32}(\partial w_2, \partial w'_2)$. \square

Lemmas 3.2–3.4, together with (3.13), show that ω is completely determined by C_{41} and C_{32} .

Theorem 3.5. *Let V_\bullet be a linear 2-groupoid with associated 3-term chain complex (W_\bullet, ∂) . There is a one-to-one correspondence between constant multiplicative 2-forms $\omega \in \Omega_\nu^2(V_2)$ and pairs (C_{41}, C_{32}) , where C_{41} is a bilinear pairing of W_0 with W_2 and C_{32} is a bilinear form on W_1 , such that*

$$(3.14) \quad C_{41}(\partial w_1, w_2) = C_{32}(\partial w_2, w_1) + C_{32}(w_1, \partial w_2)$$

for all $w_2 \in W_2$ and $w_1 \in W_1$.

Proof. In the discussion above, we have already seen how to obtain C_{41} and C_{32} from ω . To see that (3.14) holds, let $u, v \in V_3$ be defined as $u = (0, 0, 0, w_1, 0, 0, 0)$ and $v = (0, 0, w_2, 0, 0, 0, 0)$. Then

$$(3.15) \quad \begin{aligned} 0 = \delta\omega(u, v) &= -\omega((0, -w_1, -w_1, -\partial w_1), (w_2, 0, 0, 0)) \\ &= -C_{12}(w_2, w_1) - C_{13}(w_2, w_1) + C_{41}(\partial w_1, w_2). \end{aligned}$$

Equation (3.14) then follows from Lemmas 3.3 and 3.4.

In the other direction, given C_{41} and C_{32} satisfying (3.14), we can construct ω of the form (3.12), with the other components given by (3.13) and Lemmas 3.2–3.4. The skew-symmetry of C_{22} follows from (3.13), and the skew-symmetry of C_{11} follows from Lemma 3.4 and (3.14). We have already observed that such an ω will be normalized. It is long but straightforward to check that ω is multiplicative. \square

3.3. Simplicial nondegeneracy. Let V_\bullet be a linear 2-groupoid equipped with a constant 2-form $\omega \in \Omega_\nu^2(V_2)$. We will now describe the pairings A_ω and B_ω from Section 2.3 in terms of the components in (3.12).

Recall from Remark 3.1 that the 3-term complex (W, ∂) associated to V_\bullet is isomorphic to the tangent complex of V_\bullet . Thus we can view A_ω and B_ω as the restrictions of (2.5) and (2.6), respectively, to W_i . Therefore

$$\begin{aligned} A_\omega(w_0, w_2) &= \omega((0, 0, 0, w_0), (w_2, 0, 0, 0)) = C_{41}(w_0, w_2), \\ B_\omega(w_1, w'_1) &= \omega((0, 0, w_1, 0), (0, w'_1, w'_1, 0)) + \omega((0, 0, w'_1, 0), (0, w_1, w_1, 0)) \\ &= C_{32}(w_1, w'_1) + C_{32}(w'_1, w_1), \end{aligned}$$

for $w_0 \in W_0$, $w_1, w'_1 \in W_1$, and $w_2 \in W_2$. The following result is immediate.

Proposition 3.6. *ω is simplicially nondegenerate if and only if C_{41} and the symmetric part of C_{32} are both nondegenerate.*

Remark 3.7. Proposition 3.6 allows us to clearly see the difference between simplicial nondegeneracy and the ordinary notion of nondegeneracy for 2-forms. From (3.12), it is clear that ω is nondegenerate in the ordinary sense if and only if C_{41} and C_{32} are both nondegenerate. Thus the difference is that ordinary nondegeneracy considers C_{32} in its entirety, whereas simplicial nondegeneracy only considers the symmetric part of C_{32} . In the case where C_{32} is symmetric, the two notions agree; however, since C_{32} need not be symmetric in general, it is easy to find examples of 2-forms that are simplicially nondegenerate but not nondegenerate, and vice versa.

3.4. A minimal description of constant symplectic 2-groupoids. Putting Theorem 3.5 and Proposition 3.6 together, we see that there is a one-to-one correspondence between constant symplectic 2-groupoids and 3-term chain complexes (W_\bullet, ∂) equipped with a nondegenerate bilinear pairing C_{41} of W_0 with W_2 and a bilinear form C_{32} on W_1 whose symmetric part is nondegenerate, satisfying (3.14). Using the nondegeneracy of the pairings, we can further simplify the description.

Theorem 3.8. *There is a one-to-one correspondence between constant symplectic 2-groupoids and tuples $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial, r)$, where*

- W_1 and W_0 are vector spaces,
- $\langle \cdot, \cdot \rangle$ is a nondegenerate symmetric bilinear form on W_1 ,
- $\partial : W_1 \rightarrow W_0$ is a linear map such that the image of ∂^* in $W_1^* \cong W_1$ is isotropic, and

- r is an element of $\wedge^2 W_1^*$.

Proof. Given the data $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial, r)$, the corresponding 3-term chain complex is

$$(3.16) \quad W_0^* \xrightarrow{\partial^*} W_1 \xrightarrow{\partial} W_0,$$

where we are implicitly using $\langle \cdot, \cdot \rangle$ to identify W_1 with W_1^* . The equation $\partial \circ \partial^* = 0$ is equivalent to the requirement that the image of ∂^* be isotropic. We take C_{41} to be the canonical pairing of W_0^* with W_0 , and we set $C_{32} := \frac{1}{2} \langle \cdot, \cdot \rangle + r$. The equation (3.14) automatically holds. One can easily check that this gives a one-to-one correspondence. \square

3.5. Equivalences. In this subsection, we will describe equivalences between constant symplectic 2-groupoids in terms of the description given in Theorem 3.8.

From Definition 2.8 and Remark 2.9, we can see that any equivalence between constant symplectic 2-groupoid structures on a linear 2-groupoid V_\bullet is given by a 2-form $\alpha \in \Omega_\nu^2(V_1)$ satisfying (2.7) and (2.8). Since the simplicial coboundary operator δ is linear, we may assume without loss of generality that α is constant.

Let α be a constant normalized 2-form on V_1 . In terms of the decomposition $V_1 = W_1 \oplus W_0$, we can write α in block form as

$$\alpha = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & 0 \end{bmatrix},$$

where $B_{12}(w_1, w_0) = -B_{21}(w_0, w_1)$. The vanishing of the lower right block is a consequence of the assumption that α is normalized. A straightforward calculation then shows that (2.7) and (2.8) reduce in this case to the conditions

$$(3.17) \quad B_{21}(w_0, \partial w_2) = 0,$$

$$(3.18) \quad B_{21}(\partial w_1, w'_1) + B_{21}(\partial w'_1, w_1) = 0,$$

for all $w_2 \in W_2$, $w_1, w'_1 \in W_0$, and $w_0 \in W_0$.

Since $\delta\alpha$ is obviously multiplicative, it has block form

$$\delta\alpha = \begin{bmatrix} A_{11} & A_{12} & A_{13} & A_{14} \\ A_{21} & A_{22} & A_{23} & 0 \\ A_{31} & A_{32} & 0 & 0 \\ A_{41} & 0 & 0 & 0 \end{bmatrix}.$$

By Theorem 3.5, $\delta\alpha$ is completely determined by A_{41} and A_{32} . We calculate

$$\begin{aligned} A_{41}(w_0, w_2) &= \delta\alpha((0, 0, 0, w_0), (w_2, 0, 0, 0)) \\ &= \alpha((0, w_2), (\partial w_2, 0)) \\ &= B_{21}(w_0, \partial w_2), \end{aligned}$$

$$\begin{aligned} A_{32}(w_1, w'_1) &= \delta\alpha((0, 0, w_1, 0), (0, w'_1, 0, 0)) \\ &= \alpha((w_1, 0), (-w'_1, -\partial w'_1)) \\ &= -B_{11}(w_1, w'_1) - B_{12}(w_1, \partial w'_1). \end{aligned}$$

From this, we see that equation (3.17) holds if and only if $A_{41} = 0$, and that (3.18) holds if and only if A_{32} is skew-symmetric. Furthermore, we observe that there is no restriction on B_{11} , so every skew-symmetric pairing on W_1 appears as A_{32} for some choice of α satisfying (3.17) and (3.18).

It is clear from the above discussion that, in terms of the data of Theorem 3.8, equivalences between constant symplectic groupoids act transitively on r and do not affect any of the other data. Thus we have the following result.

Theorem 3.9. *There is a one-to-one correspondence between equivalence classes of constant symplectic 2-groupoids and tuples $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial)$, where*

- W_1 and W_0 are vector spaces,
- $\langle \cdot, \cdot \rangle$ is a nondegenerate symmetric bilinear form on W_1 , and
- $\partial : W_1 \rightarrow W_0$ is a linear map such that the image of ∂^* in $W_1^* \cong W_1$ is isotropic.

Remark 3.10. In each equivalence class of constant symplectic 2-groupoids, there is exactly one representative for which C_{32} is symmetric (or equivalently, in terms of the data of Theorem 3.8, where $r = 0$). In this case, we will say that the constant symplectic 2-groupoid is *symmetric*. From Remark 3.7, we can see that if (V_\bullet, ω) is a symmetric constant symplectic 2-groupoid, then $\omega \in \Omega^2(V_2)$ is genuinely nondegenerate, and therefore V_2 is genuinely symplectic.

4. INTEGRATION OF CONSTANT COURANT ALGEBROIDS

In this section, we will describe a simple class of Courant algebroids that we call *constant Courant algebroids*. We will see that constant Courant algebroids are in correspondence with equivalence classes of constant symplectic 2-groupoids.

4.1. Constant Courant algebroids. We first recall the definition of Courant algebroid.

Definition 4.1. A *Courant algebroid* is a vector bundle $E \rightarrow M$ equipped with a nondegenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$, a bundle map $\rho : E \rightarrow TM$ (called the *anchor*), and a bracket $\llbracket \cdot, \cdot \rrbracket$ (called the *Courant bracket*) on $\Gamma(E)$ such that

- (1) $\llbracket e_1, f e_2 \rrbracket = \rho(e_1)(f)e_2 + f \llbracket e_1, e_2 \rrbracket$,
- (2) $\rho(e_1)(\langle e_2, e_3 \rangle) = \langle \llbracket e_1, e_2 \rrbracket, e_3 \rangle + \langle e_2, \llbracket e_1, e_3 \rrbracket \rangle$,
- (3) $\llbracket \llbracket e_1, e_2 \rrbracket, e_3 \rrbracket = \llbracket e_1, \llbracket e_2, e_3 \rrbracket \rrbracket - \llbracket e_2, \llbracket e_1, e_3 \rrbracket \rrbracket$,
- (4) $\llbracket e_1, e_2 \rrbracket + \llbracket e_2, e_1 \rrbracket = \mathcal{D}(e_1, e_2)$,

for all $f \in C^\infty(M)$ and $e_i \in \Gamma(E)$, where $\mathcal{D} : C^\infty(M) \rightarrow \Gamma(E)$ is defined by

$$\langle \mathcal{D}f, e \rangle = \rho(e)(f).$$

A well-known but important consequence of Definition 4.1 is that, for any Courant algebroid $E \rightarrow M$, the sequence

$$(4.1) \quad T^*M \xrightarrow{\rho^*} E^* \cong E \xrightarrow{\rho} TM$$

is a 3-term chain complex of vector bundles over M . One should expect that a symplectic 2-groupoid integrating E should be such that its tangent complex is isomorphic (or at least quasi-isomorphic) to (4.1).

Definition 4.2. A *constant Courant algebroid* is a Courant algebroid of the form $W_1 \times W_0 \rightarrow W_0$, where W_0 and W_1 are vector spaces, such that

- (1) the pairing $\langle \cdot, \cdot \rangle$ is independent of the basepoint in W_0 ,
- (2) the anchor $\rho : W_1 \times W_0 \rightarrow TW_0 = W_0 \times W_0$ is independent of the basepoint, and
- (3) $\llbracket w_1, w'_1 \rrbracket = 0$ for all $w_1, w'_1 \in W_1$, viewed as constant sections.

Let $E = W_1 \times W_0 \rightarrow W_0$ be a constant Courant algebroid. The space of sections $\Gamma(E)$ can be naturally identified with $C^\infty(W_0) \otimes W_1$. Any bilinear pairing on $\Gamma(E)$ is completely determined by its restriction to constant sections, which gives a symmetric map $W_1 \otimes W_1 \rightarrow C^\infty(W_0)$. The requirement that the pairing be independent of basepoint implies the image of this map consists of constant functions; in other words, the pairing is given by a nondegenerate symmetric bilinear form on W_1 . Similarly, the requirement that the anchor be independent of basepoint implies that it is given by

$$(4.2) \quad \rho(w_1, w_0) = (\partial w_1, w_0)$$

for some linear map $\partial : W_1 \rightarrow W_0$.

The fact that $\rho \circ \rho^* = 0$ (see (4.1)) implies that $\partial \circ \partial^* = 0$ or, equivalently, the image of ∂^* in $W_1^* \cong W_1$ is isotropic.

The vanishing of the bracket of constant sections, together with axioms (1) and (4) in Definition 4.1, imply that the bracket is completely determined by the pairing and anchor. Specifically,

$$(4.3) \quad \llbracket f w_1, g w_1' \rrbracket = f \rho(w_1)(g) w_1' - g \rho(w_1')(f) w_1 + g \langle w_1, w_1' \rangle \mathcal{D}f$$

for $f, g \in C^\infty(W_0)$ and $w_1, w_1' \in W_1$.

Theorem 4.3. *There is a one-to-one correspondence between constant Courant algebroids and equivalence classes of constant symplectic 2-groupoids.*

Proof. Given a constant Courant algebroid, we have seen how to obtain the data $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial)$ for an equivalence class of constant symplectic 2-groupoids, as described in Theorem 3.9. On the other hand, given the data $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial)$, we can construct the constant Courant algebroid $E = W_1 \times W_0 \rightarrow W_0$, where the pairing on $\Gamma(E)$ agrees on constant sections with the pairing on W_0 , and the anchor and bracket are given by (4.2) and (4.3). It is a long but straightforward check that the conditions of Definition 4.1 hold. (We note that, alternatively, the Courant algebroid axioms can be checked quickly using the supergeometric formulation of [14] in local coordinates.) \square

5. INTEGRATION OF CONSTANT DIRAC STRUCTURES

5.1. Linear sub-2-groupoids. Let V_\bullet be a linear 2-groupoid. A *linear sub-2-groupoid* of V_\bullet is a simplicial subspace $L_\bullet \subseteq V_\bullet$ that is also a 2-groupoid.

Let $L_\bullet \subseteq V_\bullet$ be a linear sub-2-groupoid. Set $U_0 := L_0$, $U_1 := \ker f_0^1|_{L_1}$, and $U_2 := \ker \lambda_{2,2}|_{L_2}$. By comparing with the constructions of Section 3.1, we can see that U_\bullet is a subcomplex of (W_\bullet, ∂) . Conversely, given a subcomplex $U_\bullet \subseteq (W_\bullet, \partial)$, we can form a linear sub-2-groupoid L_\bullet , where $L_0 = U_0$, $L_1 = U_1 \oplus U_0$, and $L_2 = U_2 \oplus U_1 \oplus U_1 \oplus U_0$. This gives us the following result.

Proposition 5.1. *There is a one-to-one correspondence between linear sub-2-groupoids $L_\bullet \subseteq V_\bullet$ and subcomplexes $U_\bullet \subseteq (W_\bullet, \partial)$.*

5.2. Linear Lagrangian sub-2-groupoids. Now suppose that (V_\bullet, ω) is the symmetric constant symplectic 2-groupoid corresponding (via Theorem 3.9) to the data $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial)$. In this case, $V_2 = W_0^* \oplus W_1 \oplus W_1 \oplus W_0$, and the block form (3.12) is such that C_{41} is the natural pairing of W_0^* with W_0 and C_{32} is the pairing $\langle \cdot, \cdot \rangle$. It follows that ω is a genuine symplectic form on V_2 (see Remark 3.10).

If U_\bullet is a subcomplex of (W_\bullet, ∂) , then we can consider the space $L_2^\omega \subseteq V_2$ that is symplectic orthogonal to $L_2 = U_2 \oplus U_1 \oplus U_0$. The following gives a description of L_2^ω in terms of U_\bullet .

Lemma 5.2. $L_2^\omega = \text{Ann}(U_0) \oplus U_1^\perp \oplus U_1^\perp \oplus \text{Ann}(U_2)$.

Proof. For any $w_2 \in W_0^*$, $w_1, w'_1 \in W_1$, $w_0 \in W_0$, and $(u_2, u_1, u'_1, u_0) \in L_2$, we use (3.13) and Lemmas 3.2–3.4 together with the symmetry of C_{32} to derive the formulas

$$\begin{aligned}
 (5.1) \quad \omega((w_2, 0, 0, 0), (u_2, u_1, u'_1, u_0)) &= C_{11}(w_2, u_2) + C_{12}(w_2, u_1) + C_{13}(w_2, u'_1) + C_{14}(w_2, u_0) \\
 &= C_{32}(\partial^* w_2, u_1 + u'_1 - \partial^* u_2) - C_{41}(u_0, w_2) \\
 &= \langle \partial^* w_2, u_1 + u'_1 - \partial^* u_2 \rangle - w_2(u_0) \\
 &= w_2(\partial u_1 + \partial u'_1 - u_0),
 \end{aligned}$$

$$\begin{aligned}
 (5.2) \quad \omega((0, w_1, 0, 0), (u_2, u_1, u'_1, u_0)) &= C_{21}(w_1, u_2) + C_{23}(w_1, u'_1) \\
 &= -C_{32}(\partial^* u_2, w_1) - C_{32}(u'_1, w_1) \\
 &= -\langle w_1, u'_1 + \partial^* u_2 \rangle,
 \end{aligned}$$

$$\begin{aligned}
 (5.3) \quad \omega((0, 0, w'_1, 0), (u_2, u_1, u'_1, u_0)) &= C_{31}(w'_1, u_2) + C_{32}(w'_1, u_1) \\
 &= -C_{32}(w'_1, \partial^* u_2) + C_{32}(w'_1, u_1) \\
 &= \langle w'_1, u_1 - \partial^* u_2 \rangle,
 \end{aligned}$$

$$\begin{aligned}
 (5.4) \quad \omega((0, 0, 0, w_0), (u_2, u_1, u'_1, u_0)) &= C_{41}(w_0, u_2) \\
 &= u_2(w_0).
 \end{aligned}$$

From (5.1)–(5.4) it is immediate that $\text{Ann}(U_0) \oplus U_1^\perp \oplus U_1^\perp \oplus \text{Ann}(U_2) \subseteq L_2^\omega$.

Conversely, if $(w_2, w_1, w'_1, w_0) \in V_2$ is in L_2^ω , then from (5.1)–(5.4) we have

$$0 = \omega((w_2, w_1, w'_1, w_0), (0, 0, 0, u_0)) = -w_2(u_0)$$

for all $u_0 \in U_0$, so it follows that w_2 is in $\text{Ann}(U_0)$. Similarly,

$$0 = \omega((w_2, w_1, w'_1, w_0), (0, 0, u'_1, 0)) = w_2(\partial u'_1) - \langle w_1, u'_1 \rangle = -\langle w_1, u'_1 \rangle$$

and

$$0 = \omega((w_2, w_1, w'_1, w_0), (0, u_1, 0, 0)) = w_2(\partial u_1) + \langle w'_1, u_1 \rangle = \langle w'_1, u_1 \rangle$$

for all $u_1, u'_1 \in U_1$, so it follows that w_1 and w'_1 are in U_1^\perp . Finally,

$$0 = \omega((w_2, w_1, w'_1, w_0), (u_2, 0, 0, 0)) = -\langle w_1 + w'_1, \partial^* u_2 \rangle + u_2(w_0) = u_2(w_0)$$

for all $u_2 \in U_2$, so it follows that w_0 is in $\text{Ann}(U_2)$. \square

The following is an immediate consequence of Lemma 5.2.

Corollary 5.3. (1) L_2 is isotropic if and only if $U_0 \subseteq W_0$ and $U_2 \subseteq W_0^*$ pair to zero and $U_1 \subseteq W_1$ is isotropic.
 (2) L_2 is coisotropic if and only if the annihilator of U_2 is contained in U_0 , the annihilator of U_0 is contained in U_2 , and $U_1^\perp \subseteq U_1$.
 (3) L_2 is Lagrangian if and only if $U_2 = \text{Ann}(U_0)$ and $U_1^\perp = U_1$.

We can now obtain a description of linear Lagrangian sub-2-groupoids in terms of the data of Theorem 3.9.

Theorem 5.4. *Suppose that (V_\bullet, ω) is the symmetric constant symplectic 2-groupoid corresponding to the data $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial)$. There is a one-to-one correspondence between linear Lagrangian sub-2-groupoids $L_\bullet \subseteq V_\bullet$ and pairs (U_1, U_0) , where $U_i \subseteq W_i$ for $i = 0, 1$ are such that $U_1^\perp = U_1$ and $\partial(U_1) \subseteq U_0$.*

Proof. It is immediate from Proposition 5.1 and Corollary 5.3 that, if L_\bullet is Lagrangian, then the corresponding subspaces $U_i \subseteq W_i$ satisfy $U_1^\perp = U_1$ and $\partial(U_1) \subseteq U_0$. In the other direction, we set $U_2 = \text{Ann}(U_0)$. For any $u_2 \in U_2$ and $u_1 \in U_1$, we have

$$\langle \partial^* u_2, u_1 \rangle = u_2(\partial u_1) = 0$$

for all $u_1 \in U_1$, so $\partial^* u_2$ is in $U_1^\perp = U_1$. Therefore, U_\bullet is a subcomplex, and by Corollary 5.3 the corresponding L_2 is Lagrangian. \square

5.3. Constant Dirac structures. A Dirac structure in a Courant algebroid $E \rightarrow M$ is a subbundle $D \rightarrow M$ such that $D^\perp = D$ and $\Gamma(D)$ is closed under the Courant bracket³. The restriction of the Courant bracket $\llbracket \cdot, \cdot \rrbracket$ to any Dirac structure D is a Lie bracket, making $D \rightarrow M$ into a Lie algebroid.

Let $W_1 \times W_0 \rightarrow W_0$ be a constant Courant algebroid. We will restrict our attention to *constant Dirac structures*, i.e. Dirac structures of the form $U_1 \times W_0$, where U_1 is a subspace of W_1 .

Lemma 5.5. *Let U_1 be a subspace of W_1 . Then $U_1 \times W_0$ is a constant Dirac structure if and only if $U_1^\perp = U_1$.*

Proof. Since the pairing on $W_1 \times W_0 \rightarrow W_0$ is constant, it is immediate that $U_1^\perp = U_1$ if and only if $(U_1 \times W_0)^\perp = U_1 \times W_0$. The nontrivial part of the lemma is the observation that, in this case, closure under the Courant bracket is a consequence of the isotropic property. To see this, suppose that $U_1 \subseteq W_1$ is isotropic. Then, from (4.3), we have

$$\llbracket fu, gu' \rrbracket = f\rho(u)(g)u' - g\rho(u')(f)u + g\langle u, u' \rangle \mathcal{D}f$$

for $f, g \in C^\infty(W_0)$ and $u, u' \in U_1$. The last term vanishes, and the other two terms are clearly in $\Gamma(U_1 \times W_0)$. \square

Given $U_1 \subseteq W_1$ such that $U_1^\perp = U_1$, we may apply Theorem 5.4 to the pair (U_1, W_0) to obtain a linear Lagrangian sub-2-groupoid $L_\bullet \subseteq V_\bullet$ that is *wide*, in the sense that $L_0 = V_0$. It is clear from the correspondence of Theorem 5.4 that every wide linear Lagrangian sub-2-groupoid arises in this manner. Using Lemma 5.5, we then obtain the following result.

Theorem 5.6. *Suppose that (V_\bullet, ω) is the symmetric constant symplectic 2-groupoid corresponding to the data $(W_1, W_0, \langle \cdot, \cdot \rangle, \partial)$, and let $W_1 \times W_0 \rightarrow W_0$ be the corresponding constant Courant algebroid. There is a one-to-one correspondence between constant Dirac structures $U_1 \times W_0 \subseteq W_1 \times W_0$ and wide linear Lagrangian sub-2-groupoids $L_\bullet \subseteq V_\bullet$.*

³We note that we are using the condition $D^\perp = D$ in place of the usual requirement that D be maximally isotropic. In most cases of interest, E has signature (n, n) , and the two conditions are equivalent. However, if E does not have signature (n, n) , then, according to the definition we are using, there do not exist any Dirac structures in E .

- Remarks 5.7.* (1) From part (3) of Corollary 5.3, we can see that a linear Lagrangian sub-2-groupoid corresponding to the subcomplex $U_{\bullet} \subseteq (W_{\bullet}, \partial)$ is wide if and only if $U_2 = \{0\}$. Therefore, a wide linear Lagrangian sub-2-groupoid is actually a 1-groupoid $U_1 \oplus W_0 \rightrightarrows W_0$. It is straightforward to check that this is the Lie groupoid that integrates the Lie algebroid $U_1 \times W_0 \rightarrow W_0$.
- (2) We stress that there are many complications involved in extending the result of Theorem 5.6 to the nonlinear situation. In particular, we expect that a weaker definition of Lagrangian sub-2-groupoid, using some of the ideas of derived symplectic geometry [13], will be required.

REFERENCES

1. A. Coste, P. Dazord, and A. Weinstein, *Groupoïdes symplectiques*, Publications du Département de Mathématiques. Nouvelle Série. A, Vol. 2, Publ. Dép. Math. Nouvelle Sér. A, vol. 87, Univ. Claude-Bernard, Lyon, 1987, pp. i–ii, 1–62.
2. T. Courant and A. Weinstein, *Beyond Poisson structures*, Action hamiltoniennes de groupes. Troisième théorème de Lie (Lyon, 1986), Travaux en Cours, vol. 27, Hermann, Paris, 1988, pp. 39–49.
3. T. J. Courant, *Dirac manifolds*, Trans. Amer. Math. Soc. **319** (1990), no. 2, 631–661.
4. I. Dorfman, *Dirac structures and integrability of nonlinear evolution equations*, Nonlinear Science: Theory and Applications, John Wiley & Sons, Ltd., Chichester, 1993.
5. J. Duskin, *Higher-dimensional torsors and the cohomology of topoi: the abelian theory*, Applications of sheaves (Proc. Res. Sympos. Appl. Sheaf Theory to Logic, Algebra and Anal., Univ. Durham, 1977), Lecture Notes in Math., vol. 753, Springer, Berlin, 1979, pp. 255–279.
6. E. Getzler, *Differential forms on stacks*, 2014, Slides from minicourse at Winter School in Mathematical Physics, Les Diablerets.
7. Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Birkhäuser, Verlag, Basel, 2009.
8. A. Henriques, *Integrating L_{∞} -algebras*, Compos. Math. **144** (2008), no. 4, 1017–1045.
9. D. Li-Bland and P. Ševera, *Integration of exact Courant algebroids*, Electron. Res. Announc. Math. Sci. **19** (2012), 58–76.
10. Z. Liu, A. Weinstein, and P. Xu, *Manin triples for Lie bialgebroids*, J. Differential Geom. **45** (1997), no. 3, 547–574.
11. R. A. Mehta and X. Tang, *Symplectic structures on the integration of exact courant algebroids*, arXiv: 1310.6587.
12. ———, *From double Lie groupoids to local Lie 2-groupoids*, Bull. Braz. Math. Soc. (N.S.) **42** (2011), no. 4, 651–681.
13. T. Pantev, M. Toën, B. Vaquié, and G. Vezzosi, *Shifted symplectic structures*, Publ. Math. Inst. Hautes Études Sci. **117** (2013), 271–328.
14. D. Roytenberg, *On the structure of graded symplectic supermanifolds and Courant algebroids*, Quantization, Poisson brackets and beyond (Manchester, 2001), Contemp. Math., vol. 315, Amer. Math. Soc., Providence, RI, 2002, pp. 169–185.
15. P. Ševera, *Some title containing the words “homotopy” and “symplectic”, e.g. this one*, Travaux mathématiques. Fasc. XVI, Trav. Math., XVI, Univ. Luxemb., Luxembourg, 2005, pp. 121–137.
16. Y. Sheng and C. Zhu, *Higher extensions of Lie algebroids and application to Courant algebroids*, 2011, arXiv:1103.5920.
17. P. Ševera and M. Širaň, *Integration of differential graded manifolds*, 2015, arXiv: 1506.04898.
18. P. Xu, *Momentum maps and Morita equivalence*, J. Differential Geom. **67** (2004), no. 2, 289–333.
19. C. Zhu, *Kan replacement of simplicial manifolds*, Lett. Math. Phys. **90** (2009), no. 1-3, 383–405.

DEPARTMENT OF MATHEMATICS & STATISTICS, SMITH COLLEGE, 44 COLLEGE LANE, NORTHAMPTON, MA 01063

E-mail address: `rmehta@smith.edu`

DEPARTMENT OF MATHEMATICS, WASHINGTON UNIVERSITY IN SAINT LOUIS, ONE BROOKINGS DRIVE, SAINT LOUIS, MISSOURI, USA 63130

E-mail address: `xtang@math.wustl.edu`