

4-2001

Computational Geometry Column 41

Joseph O'Rourke

Smith College, jorourke@smith.edu

Follow this and additional works at: https://scholarworks.smith.edu/csc_facpubs

Part of the [Computer Sciences Commons](#), and the [Geometry and Topology Commons](#)

Recommended Citation

O'Rourke, Joseph, "Computational Geometry Column 41" (2001). Computer Science: Faculty Publications, Smith College, Northampton, MA.

https://scholarworks.smith.edu/csc_facpubs/66

This Article has been accepted for inclusion in Computer Science: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact scholarworks@smith.edu

Computational Geometry Column 41

Joseph O'Rourke*

Abstract

The recent result that n congruent balls in \mathbb{R}^d have at most 4 distinct geometric permutations is described.

A line ℓ *stabs* a set S of geometric objects in \mathbb{R}^d if ℓ intersects every member of S . Such a stabber is often called a *line traversal* of S . When the objects are pairwise disjoint convex bodies, then a line traversal induces a *geometric permutation* of the objects: either of the two piercing orders of the objects (one the reverse of the other). Basic bounds on the number of geometric permutations were established about a decade ago: the number is in $\Omega(n^{d-1})$ [KLL92] and $O(n^{2d-2})$ [Wen90], with a tight bound of $2n - 2$ known for $d = 2$ [ES90]. The study of geometric permutations was revitalized by a focus specifically on balls by Smorodinsky, Mitchell, and Sharir [SMS00]. They closed the gap in this special case to $\Theta(n^{d-1})$, a result which was quickly followed by an extension to the same bound for “fat” convex objects [KV99].




Figure 1: (a) Two line traversals with permutations $(1, 2, 3, 4, 5)$ and $(1, 2, 4, 3, 5)$; (b) Three permutations achievable.

One fascinating line of investigation opened in [SMS00], and independently by Asinowski and Katchalski [Asi98], is the even more special case of congruent balls. They proved that sufficiently many congruent disks in \mathbb{R}^2 admit only a constant number of geometric permutations—in fact, just 2. A little experimentation (Fig. 1a) quickly reveals the operable constraint: one line traversal necessarily strings out the disks roughly along a line, and any other stabbing line cannot deviate too much if it is to pierce all the disks. The only example that admits more than 2 permutations is a triangular arrangement of three disks (Fig. 1b); the proof in [Asi98] shows that the bound of 2 holds for all $n \geq 4$. If the disks are not congruent, then $\Omega(n)$ geometric permutations are possible.

Smorodinsky et al. left open the question of bounds for congruent balls in $d \geq 3$ dimensions, but conjectured that again only $O(1)$ geometric permutations are achievable. This conjecture was settled by Zhou and Suri, who established an upper bound of 16 for all d , for a sufficiently large number n of balls. Quickly they [ZS01], and independently Huang, Xu, and Chen [HXC01], improved this bound to 4, where it stands at this writing. Both sets of researchers have also extended their results to noncongruent balls of bounded radius ratio, delimiting when the n^{d-1} behavior kicks in.

A sense for the congruent-balls problem can be obtained by attempting to stab four spheres stacked as cannonballs (displaced slightly to ensure pairwise disjointness). See Fig. 2a. There is no line traversal for this

*Dept. of Computer Science, Smith College, Northampton, MA 01063, USA. orourke@cs.smith.edu. Supported by NSF Grant CCR-9731804.

set of balls, although I do not know a simple proof of this claim. Again, the existence of a line traversal ℓ requires the balls to be strung out along ℓ , a requirement incompatible with the clumping in this stack.




Figure 2: (a) A stabbing of three out of four congruent balls; (b) The centroids of the two pairs of unit-radius balls are separated by $\sqrt{2}$ (dashed segment).

I now sketch the proof of Zhou and Suri for a set S of unit-radius balls in \mathbb{R}^d , relying on [Zho00]. They call a pair of balls in the set S a *switched pair* if, like disks 3 and 4 in Fig. 1a, there are line traversals for S that meet them in two different orders. They first prove that the balls in any switched pair must be close to one another, and next, that a stabber ℓ of such a switched pair must be nearly perpendicular to the line through the centers of the two balls, and pass very nearly through their common center of gravity (centroid). (These constraints are evident in Fig. 1a.)

From this they prove that each ball of S can participate in at most one switched pair, which in turn leads to the conclusion that the two balls in a switched pair must appear consecutively in every geometric permutation of S .

The theorem is established by computing upper and lower bounds on the “separation” between two distinct switched pairs. They prove the distance between the centers of gravity of two switched pairs has a lower bound of $\sqrt{2} - \epsilon(n)$, where $\epsilon(n) > 0$ and $\lim_{n \rightarrow \infty} \epsilon(n) = 0$. This $\sqrt{2}$ term can be seen to derive (roughly) from the distance between the centers of gravity of the two pairs of balls in the cannonball stack of Fig. 2b: this is the distance between the midpoints of opposite edges of the tetrahedron formed by the ball centers.

So switched pairs cannot be too close together. Neither can they be too far apart, a claim we will leave to the intuition that the need for stabbers to pass close to the centers of gravity of the switched pairs limits their separation. Combining the explicit bounds leads to the conclusion that there can be at most two switched pairs, which in turn establishes a bound of 4 on the number of geometric permutations for sufficiently large n .

It remains possible that the correct bound is 2, again for sufficiently large n . For small values of n , examples might exist that determine more than 4 permutations. But as far as I know, no one has found a collection of $n \geq 4$ congruent balls in \mathbb{R}^3 that have more than one switched pair, or admit more than 2 geometric permutations.

Although the pursuit of geometric permutations has been largely driven by theoretical interests, there is an important application in \mathbb{R}^3 to computer graphics: lines of sight between mutually visible objects represent stabbers. Bounds on the number of stabbers can translate into bounds on visibility complexity. This makes especially interesting the result in [ZS01] that a collection of pairwise disjoint, axis-aligned boxes (e.g., the “bounding boxes” so useful in graphics) admit only 2^{d-1} geometric permutations, i.e., 4 in \mathbb{R}^3 . This bound is tight.

References

- [Asi98] A. Asinowski. Common transversals and geometric permutations. Master's thesis, Technion IIT, Haifa, Israel, 1998.
- [ES90] H. Edelsbrunner and M. Sharir. The maximum number of ways to stab n convex non-intersecting sets in the plane is $2n - 2$. *Discrete Comput. Geom.*, 5:35–42, 1990.
- [HXC01] Y. Huang, J. Xu, and D. Z. Chen. Geometric permutations of high dimensional spheres. In *Proc. 12th ACM-SIAM Sympos. Discrete Algorithms*, pages 244–245, 2001.
- [KLL92] M. Katchalski, T. Lewis, and A. Liu. The different ways of stabbing disjoint convex sets. *Discrete Comput. Geom.*, 7:197–206, 1992.
- [KV99] M.J. Katz and K. R. Varadarajan. A tight bound on the number of geometric permutations of convex fat objects in \mathbb{R}^d . Dept. Comput. Sci., Ben-Gurion University of the Negev, Israel, 1999.
- [SMS00] S. Smorodinsky, J.S.B. Mitchell, and M. Sharir. Sharp bounds on geometric permutations for pairwise disjoint balls in \mathbb{R}^d . *Discrete Comput. Geom.*, 23:247–259, 2000.
- [Wen90] R. Wenger. Upper bounds on geometric permutations for convex sets. *Discrete Comput. Geom.*, 5:27–33, 1990.
- [Zho00] Y. Zhou. *Shape Sensitive Geometric Complexity*. PhD thesis, Washington University, 2000. St. Louis, MO, USA.
- [ZS01] Y. Zhou and S. Suri. Shape sensitive geometric permutations. In *Proc. 12th ACM-SIAM Sympos. Discrete Algorithms*, pages 234–243, 2001.