
5-17-2019

Ni-Catalyzed Cross-Electrophile Coupling for the Synthesis of Skipped Polyenes

Catherine P. McGeough
Massachusetts Institute of Technology

Alexandra E. Strom
Stanford University, astrom@smith.edu

Timothy F. Jamison
Massachusetts Institute of Technology

Follow this and additional works at: https://scholarworks.smith.edu/chm_facpubs

 Part of the [Chemistry Commons](#)

Recommended Citation

McGeough, Catherine P.; Strom, Alexandra E.; and Jamison, Timothy F., "Ni-Catalyzed Cross-Electrophile Coupling for the Synthesis of Skipped Polyenes" (2019). Chemistry: Faculty Publications, Smith College, Northampton, MA.
https://scholarworks.smith.edu/chm_facpubs/79

This Article has been accepted for inclusion in Chemistry: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact scholarworks@smith.edu



Published in final edited form as:

Org Lett. 2019 May 17; 21(10): 3606–3609. doi:10.1021/acs.orglett.9b01019.

Ni-Catalyzed Cross-Electrophile Coupling for the Synthesis of Skipped Polyenes


Catherine P. McGeough[‡], Alexandra E. Strom^{†,‡}, Timothy F. Jamison^{*}

Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

Abstract

Skipped polyenes featuring high (*E*)-selectivity and varying methyl substitution patterns are synthesized using a nickel-catalyzed cross-coupling reaction between allyl trifluoroacetates and vinyl bromides. The utility of this cross-electrophile coupling is showcased in part, by the synthesis of the RST fragment of the marine ladder polyether, maitotoxin. Construction of this fragment is particularly challenging due to the alternating methyl substitution pattern.

Graphical Abstract



Skipped polyenes featuring various methyl substitution patterns are commonly found in biologically active natural products with anticancer, antibacterial, and antiviral properties.¹ These molecules can also provide efficient entry to the fused cyclic ether skeletons of marine ladder polyether natural products by way of epoxide ring opening cascade reactions.^{2a}

Given their significance, researchers have developed a number of individualized syntheses² of skipped polyenes as well as more generalized methods, but which are limited by particular substitution and unsaturation patterns. The Trost group, for example, recently reported the ruthenium catalyzed synthesis of 1,3,6-trienes via the coupling of propene with two alkyl substituted alkynes (Figure 1a).^{3a} The Micalizio lab alternatively, demonstrated a titanium mediated coupling reaction of vinylcyclopropanes and alkynes to generate 1,4-dienes and 1,4,7-trienes (Figure 1b).^{3b}

Herein we describe a more flexible approach to polyene synthesis that proceeds via the Ni-catalyzed cross-electrophile coupling of activated allyl alcohol **3** with vinyl bromide **4** (Figure 1c). Linear polyenes, including those that display a variety of methyl substitution

^{*}Corresponding Author: tfj@mit.edu.

[†]Present Addresses: Department of Chemistry, Smith College, 100 Green St. Northampton, MA 01063

[‡]These authors contributed equally.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and spectral data for polyenes **10**, **21–28**, **35–40**, as well as compounds **42–51**, **7**, **6**, and **1** (PDF).

patterns, are readily accessed with high (*E*)-selectivity while tolerating a free alcohol group. Among the polyenes synthesized using this strategy is the all linear (*E*)-pentaene **7**, which we use to prepare the RST fragment of the ladder polyether natural product, maitotoxin via an epoxide-opening cascade reaction (Figure 1d).^{2,4}

Few examples of Ni-catalyzed cross electrophile reductive couplings between allyl and sp² electrophiles are described in the literature and those which are, provide mixtures of branched, linear (*E*)-, and linear (*Z*)-products.⁵ To achieve high linear (*E*)-selectivity, our aim was to limit alkene isomerization by choosing coupling partners that react quickly under mild conditions and that have excellent cross-selectivity. While there are other metals that catalyze allyl vinyl couplings,^{6a} we were inspired by the Gong group^{6b} who reported the Ni-catalyzed coupling of aryl halides and allyl acetates (Figure 2a). We thus began our investigations by applying similar reaction conditions to the coupling of vinyl bromide **9** and compound **8** featuring allyl acetate (**8a**), allyl pivalate (**8b**) and allyl trifluoroacetate (**8c**) activating groups (Figure 2b). Nickel (II) bromide (10 mol %) and 2,2-bipyridine (10 mol %) served as the catalyst and ligand respectively, while zinc acted as the metal reductant. The addition of magnesium chloride and pyridine to the reaction mixture was found to be essential as similarly noted by the Gong group.^{6b} While each of the three Ni-catalyzed reactions examined provided only the desired linear coupled products **10/11** (i.e. no branch product), the highly reactive **8c** provided the best (*E*)-selectivity (**10:11**, *E:Z*, 95:5). Examination of the ¹H NMR spectrum of the crude reaction mixture revealed complete consumption of **8c**, but the formation of homo-coupled and hydrolyzed side-products, thereby accounting for the low yield of **10/11** (20%).⁷

Accordingly, we next conducted optimization studies to further improve the reaction outcome. Since we observed a slight increase in the yield of **10/11** (23%) upon doubling the equivalents of nickel (II) bromide and 2,2-bipyridine (i.e. 10 mol % to 20 mol %) under otherwise identical conditions shown in Figure 2b, we maintained the use of this higher catalyst and ligand loading throughout our optimization studies. As shown in Entries 1 – 4 of Table 1, decreasing the reaction time from 6 h to 1 h and lowering the reaction temperature from 60 °C to 40 °C significantly enhanced the product (*E*-selectivity (>99:1) and yield (40%). Utilizing manganese, rather than zinc as the reductant (Entries 5–6) and increasing the equivalents of vinyl bromide **9** from 2.0 to 3.0 (Entries 6 vs 7) provided additional improvement in product yield (*ca.* 63%) without affecting the (*E*)-selectivity. Notably, decreasing the equivalents of manganese from 4.0 to 2.0 had little impact (Table 1, Entries 7 vs 8). Although we observed near exclusive (*E*)-selectivity, the yield of this transformation was not improved beyond 63% until we noted the dramatic impact of the reaction stir rate as a result of the heterogeneity of the manganese reductant. Reaction stir rates of ~ 600 rpm for example, consistently provided product yields of *ca.* 50% – 60%, whereas a stir rate of 945 rpm led to 86% product yield (Table 1, Entry 9). Increasing the stir rate beyond 945 rpm had no additional effect.

As shown in Table 2, we then examined the scope of the vinyl halide coupling partners using the optimized conditions established in Table 1, Entry 9. We found that vinyl bromides substituted at the α-position (Entries 1–2, 6–8) resulted in moderate to good yields of the corresponding products (48% – 87%). Vinyl halides with α,β-substitution (Entries 3–4), β-

substitution (Entry 5), as well as β,β -substitution (Entry 9) provided a similar range in product yield (48% – 72%).

Next, we explored a range of allyl trifluoroacetate coupling partners (Table 3). Both di- and tri-substituted allyl trifluoroacetates (Entries 1–3) were coupled with vinyl bromide **16** without observation of isomerization in low to moderate yields (12% – 46%). In addition to desired product, we observed both homocoupling and hydrolysis of the trifluoroacetate starting materials, which accounted for the lower yields in some of these reactions. Tri-substituted allyl trifluoroacetates **32** and **29** coupled with vinyl bromide **9** in good yields (Entries 4–5, 52% – 61%). Interestingly, nerol **32** isomerized when coupled with vinyl bromide **5** to give linear (*E*) product **10** (Entry 4). Due to the incomplete isomerization of **33** and **34** during the coupling to **9**, we observed *E/Z* product mixtures as shown in Entries 6 and 7.

To further demonstrate the utility of this cross electrophile coupling reaction, we implemented it in the construction of pentaene **7**. This intermediate was then used to synthesize the RST fragment of maitotoxin via an epoxide-opening cascade reaction (Figure 1d). Maitotoxin is a member of the polyether marine biotoxin class of natural products and has been reported to exhibit anticancer activity.⁸ The synthesis of the RST fragment is particularly challenging due to its alternating methyl substitution pattern and 6-7-6 ring structure. While a previous synthesis of the QRSTU ring fragment of maitotoxin was reported by the Nicolaou group in 2010, they found the installation of the alternating methyl substituents to be cumbersome and challenging.⁹

As presented in Scheme 1, we commenced the synthesis of pentaene **7** by activating known compound **41** with trifluoroacetic anhydride. After aqueous workup, **42** was immediately coupled with vinyl bromide **43** to furnish the desired product **44** in 46% yield. Notably, the use of 4,4'-dimethoxy-2,2'-bipyridine in this coupling reaction minimized hydrolysis of the trifluoroacetate electrophile. Alcohol **44** was then transformed to alkyl bromide **45** and coupled to vinyl bromide **46** using a Negishi cross-coupling reaction.¹⁰ Although we attempted to synthesize **48** from alcohol **44** using nickel-catalyzed cross-electrophile coupling methodology, we could only obtain inseparable *E/Z* mixtures of the desired product. Following the formation of **47**, the primary alcohol was then deprotected and activated to generate **49**. A final coupling reaction under our developed nickel-catalyzed conditions provided the desired pentene **7**.

As illustrated in Scheme 2, a Sharpless epoxidation followed by Shi epoxidation was next implemented to chemoselectively and enantioselectively oxidize four of the five alkenes of pentane **7**. The free alcohol of the resulting product **51** was subsequently activated by forming the corresponding *tert*-butyl carbonate **6**. The final tetraepoxide opening cascade reaction was then initiated in the presence of the $\text{BF}_3 \cdot \text{OEt}_2$ catalyst at -40°C .⁴ Following acetylation of the secondary alcohol, the RST fragment **1** of maitotoxin was obtained. To the best of our knowledge, this is the first tetraepoxide opening cascade that has been performed to synthesize a natural product fragment containing both 6- and 7-membered rings.

In conclusion, we have demonstrated the Ni-catalyzed cross-electrophile coupling of activated allyl alcohols with vinyl bromides as a general strategy to construct skipped polyenes displaying high (*E*)-selectivity. This method has been applied to the synthesis of the all-(*E*), all-linear pentaene **7**, which we have further shown is a viable precursor for the synthesis of the challenging RST fragment of maitotoxin **1**.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


ACKNOWLEDGMENT

The authors thank NIGMS for financial support (GM72566 and a fellowship to A.E.S. GM117710). We also thank Liam Kelly [MIT] and Dr. Bruce Adams [MIT] for their assistance in obtaining HRMS data and NMR data, respectively. Lastly, we are very grateful to Dr. Rachel L. Beingsner [MIT] for her help with the preparation of this manuscript.


REFERENCES

- (1). (a) Yang F; Newsome J; Curran DJ *Am. Chem. Soc* 2006, 128, 14200–14205. (b) Glaus F; Dedi D; Tare P; Nagaraja V; Rodrigues L; Ainsa J; Kunze J; Schneider G; Hartkoorn R; Cole S; Altmann KJ *Org. Chem* 2018, 83, 7150–7172. (c) Singh P; Liu W; Gougoutas J; Malley M; Porubcan M; Trejo W; Wells J; Sykes RJ *Antibiot.* 1988, 41, 446–453. (d) Smith T; Kuo W; Bock V; Roizen J; Balskus E; Theberge A *Org. Lett* 2007, 9, 1153–1155. [PubMed: 17316014] (e) Surup F; Shojaei H; von Zezschwitz P; Kunze B; Grond S *Bioorg. Med. Chem* 2008, 16, 1738–1746. [PubMed: 18054490] (f) Zeng Z; Chen C; Zhang Y *Org. Chem. Front* 2018, 5, 838–840. (g) Kaysser L; Bernhardt P; Nam S; Loesgen S; Ruby J; Skewes-Cox P; Jensen P; Fenical W; Moore BJ *Am. Chem. Soc.* 2012, 134, 11988–11991. (h) Ronzitti G; Callegari F; Malaguit C.; Rossini G. *Br. J. Cancer* 2004, 90, 1100–1107. [PubMed: 14997213]
- (2). (a) Vilotjevic I; Jamison TF *Science* 2007, 317, 1189–1192. [PubMed: 17761875] (b) Morten C; Jamison T. F. J. *Am. Chem. Soc* 2009, 131, 6678–6679. (c) Katcher M; Jamison T. *Tetrahedron* 2018, 74, 1111–1122. [PubMed: 29479119]
- (3). (a) Trost B; Gholami HJ *Am. Chem. Soc* 2018, 140, 11623–11626. (b) Macklin T; Micalizio G. *Nat. Chem* 2010, 2, 638–643. [PubMed: 20651725]
- (4). (a) McDonald FE; Bravo F; Wang X; Wei X; Toganoh M; Rodriguez JR; Do B; Neiwert WA; Hardcastle KJ *Org. Chem* 2002, 67, 2515–2523. (b) Bravo F; McDonald F; Neiwert W; Hardcastle K. *Org. Lett* 2004, 6, 4487–4489. [PubMed: 15548057] (c) Nakanishi K. *Toxicol* 1985, 23, 473–479. [PubMed: 3895583] (d) Nicolaou K; *Angew. Chem., Int. Ed* 1996, 35, 589–607.
- (5). (a) Anka-Lufford LL; Prinsell M; Weix DJ *Org. Chem* 2012, 77, 9989–10000. (b) Cui X; Wang S; Zhang Y; Wei D; Qian Q; Gong H. *Org. Biomol. Chem* 2013, 11, 3094–3097. [PubMed: 23576012] (c) Gomes P; Gosmini C; Perichon J. *Org. Lett* 2003, 5, 1043–1045. [PubMed: 12659569]
- (6). (a) Seomoon D; Lee K; Kim H; Lee P *Chem. Eur. J* 2007, 13, 5197–5206. [PubMed: 17367099] (b) Cui X; Wang S; Zhang Y; Wei D; Qian Q; Gong H. *Org. Biomol. Chem* 2013, 11, 3094–3097. [PubMed: 23576012]
- (7). The use of dried versus non-dried DMA did not affect the amount of hydrolysis observed, so we suspect that the hydrolysis of the starting material is catalyzed by reagents in the reaction.
- (8). (a) Nicolaou K; Aversa R *Isr. J. Chem* 2011, 51, 359–177. [PubMed: 21709816] (b) Nicolaou KC; Heretsch P; Nakamura T; Rudo A; Murata M; Konoki K. *J. Am. Chem. Soc* 2014, 136, 16444–16451. [PubMed: 25374117]
- (9). Nicolaou KC; Gelin CF; Seo JH; Huang Z; Umezawa TJ *Am. Chem. Soc* 2010, 132, 9900–9907.
- (10). Li J; Ballmer S; Gillis E; Fuji S; Schmidt M; Palazzolo A; Lehmann J; Morehouse G; Burke M *Science* 2015, 347, 1221–1226. [PubMed: 25766227]

Previous Skipped Polyene Syntheses



General coupling scheme (this work)



Proposed maitotoxin RST fragment retrosynthesis (this work)





Figure 1.

Skipped polyene syntheses by the (a) Trost and (b) Micalizio groups. (c) Proposed polyene synthesis described herein. (d) Synthetic plan for the RST fragment of maitotoxin via pentaene 7.

a) Previous cross electrophile reductive coupling conditions



b) Our initial coupling conditions







Figure 2. (a) Allyl-aryl coupling conditions developed by the Gong group. (b) Effect of the activating groups of 8 on product (*E*)-selectivity.



Scheme 1.
Synthesis of pentaene 7.

**Scheme 2.**

Epoxidation of pentaene 7 followed by a tetra-epoxide opening cascade for the formation of the RST fragment of maitotoxin 1.

Table 1.Optimization of the coupling between 8c and 9.^a


entry	9 (equiv)	reductant (equiv)	<i>t</i> (°C)	<i>t</i> (h)	stirring (rpm)	yield (%) (10:11)
1	2.0	Zn _(s) (2.0)	60	6	600	23% (96:4)
2	2.0	Zn _(s) (2.0)	60	3	600	32% (97:3)
3	2.0	Zn _(s) (2.0)	60	1	600	30% (97:3)
4	2.0	Zn _(s) (2.0)	40	1	600	40% (>99:1)
5	2.0	Zn _(s) (4.0)	40	1	600	41% (>99:1)
6	2.0	Mn _(s) (4.0)	40	1	600	52% (>99:1)
7	3.0	Mn _(s) (4.0)	40	1	600	63% (>99:1)
8	3.0	Mn _(s) (2.0)	40	1	600	62% (>99:1)
9	3.0	Mn _(s) (2.0)	40	1	945	86% (>99:1)

^aYields are determined via NMR using 1,3,5-trimethoxybenzene as an internal standard.

Table 2.

Coupling of **8c** with vinyl halides **12** - **20** under optimized conditions.^a

entry	vinyl halide	product	yield
1			77%
2			66%
3			60%
4			72%
5			48%
6			87%
7			76%
8			56% ^b
9			54%

^aYields are determined via NMR using 1,3,5-trimethoxybenzene as an internal standard.^bReaction time of 30 minutes.

Table 3.

Coupling of vinyl bromide 16 or 9 with allyl electrophiles 29 - 34 under optimized conditions.^a

entry	allyl electrophile	vinyl bromide	product(s)	yield (E:Z)
1		16		40%
2		16		12%
3		16		46%
4		9		52%
5		9		61%
6		9		51% (0.4:0.6)
7		9		37% (2:1)

^aYields are determined via NMR using 1,3,5-trimethoxybenzene as an internal standard.