
3-1-2023

Aggregate Effects of Proliferating Low-Earth-Orbit Objects and Implications for Astronomical Data Lost in the Noise

John C. Barentine
LLC

Aparna Venkatesan
University of San Francisco

Jessica Heim
University of Southern Queensland

James Lowenthal
Smith College, jlowenth@smith.edu

Miroslav Kocifaj
Institute of Construction and Architecture Slovak Academy of Sciences

See next page for additional authors

Follow this and additional works at: https://scholarworks.smith.edu/ast_facpubs



Part of the [Astrophysics and Astronomy Commons](#)

Recommended Citation

Barentine, John C.; Venkatesan, Aparna; Heim, Jessica; Lowenthal, James; Kocifaj, Miroslav; and Bará, Salvador, "Aggregate Effects of Proliferating Low-Earth-Orbit Objects and Implications for Astronomical Data Lost in the Noise" (2023). Astronomy: Faculty Publications, Smith College, Northampton, MA.
https://scholarworks.smith.edu/ast_facpubs/82

This Article has been accepted for inclusion in Astronomy: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact scholarworks@smith.edu

Authors

John C. Barentine, Aparna Venkatesan, Jessica Heim, James Lowenthal, Miroslav Kocifaj, and Salvador Bará

Aggregate effects of proliferating low-Earth-orbit objects and implications for astronomical data lost in the noise

Received: 31 August 2022

Accepted: 30 January 2023

Published online: 20 March 2023

 Check for updates

John C. Barentine^{1,2}✉, Aparna Venkatesan³, Jessica Heim⁴, James Lowenthal⁵, Miroslav Kocifaj^{6,7} & Salvador Bará⁸

The rising population of artificial satellites and associated debris in low-altitude orbits is increasing the overall brightness of the night sky, threatening ground-based astronomy as well as a diversity of stakeholders and ecosystems reliant on dark skies. We present calculations of the potentially large rise in global sky brightness from space objects in low Earth orbit, including qualitative and quantitative assessments of how professional astronomy may be affected. Debris proliferation is of special concern: we calculate that all log-decades in debris size contribute approximately the same amount of night sky radiance, so debris-generating events are expected to lead to a rapid rise in night sky brightness along with serious collision risks for satellites from centimetre-sized objects. This increase in low-Earth-orbit traffic will lead to loss of astronomical data and diminish opportunities for ground-based discoveries as faint astrophysical signals become increasingly lost in the noise. Lastly, we discuss the broader consequences of brighter skies for a range of sky constituencies, equity/inclusion and accessibility for Earth- and space-based science, and cultural sky traditions. Space and dark skies represent an intangible heritage that deserves intentional preservation and safeguarding for future generations.

Orbital space near the Earth has been transformed radically since the launch of the first artificial satellite in 1957. The number of functional satellites in low Earth orbit (LEO) has more than doubled since early 2019 due to the advent of large groups of satellites informally known as megaconstellations. As LEO becomes an increasingly congested space, the risk of collisions between and among objects increases exponentially, as well as the likelihood of an uncontrolled chain reaction of debris-generating events.

In only three years, satellite megaconstellations have become an increasingly serious threat to astronomy. We are witnessing a dramatic,

fundamental and perhaps semi-permanent transformation of the night sky without historical precedent and with limited oversight. The number of satellites planned for launch in the 2020s and beyond is enormous¹, driven primarily by private companies motivated by profit. Understanding their potential effects on astronomy requires more basic data on the varying brightnesses of satellites in orbit. This is beginning to be studied through simulations, data processing solutions and calculations of impacts^{2–5}; see also the Trailblazer open data repository⁶. Nevertheless, we fear that faint astrophysical signals will become increasingly lost in the noise due to satellite megaconstellations.

¹Dark Sky Consulting, LLC, Tucson, AZ, USA. ²Consortium for Dark Sky Studies, University of Utah, Salt Lake City, UT, USA. ³University of San Francisco, San Francisco, CA, USA. ⁴University of Southern Queensland Centre for Astrophysics, Toowoomba, Queensland, Australia. ⁵Smith College, Northampton, MA, USA. ⁶ICA, Slovak Academy of Sciences, Bratislava, Slovakia. ⁷Department of Experimental Physics, FMPI, Comenius University, Bratislava, Slovakia. ⁸Agrupación Astronómica Coruñesa Ío, A Coruña, Spain. ✉e-mail: john@darkskyconsulting.com

LEO crowding is changing the nature of the space environment

Direct illumination by sunlight of functional satellites, failed satellites, leftover launch hardware and debris fragments (collectively ‘space objects’) makes them visible as streaks or trails in astronomical optical and infrared images, which can compromise scientific data^{4,7–11}. Myriad smaller objects contribute to elevating the diffuse brightness of the night sky. Kocifaj et al. estimated that, even before the megaconstellation era began, space objects already contributed additional light at the zenith amounting to as much as 10% of an assumed natural background level¹².

Large objects such as intact satellites make a small, but non-negligible, contribution to diffuse night sky brightness (NSB)¹³. The generation of small debris is a greater concern: as of mid-2022, the number of objects larger than 10 cm in size orbiting the Earth was estimated to be around 36,500 (ref. ¹⁴). Below that size scale there is little publicly available information on the population of debris. The number of centimetre-sized objects that could seriously damage satellites in collisions is probably around 1,000,000.

These small objects also disproportionately contribute to rising diffuse NSB. The cumulative numbers $N(>D)$ of space objects with a lower size limit (D) ranging from 1 μm to roughly 5 m in diameter adopted by Kocifaj et al.¹²—representing the latest and most complete data available to civilian scholars—imply that all log-decades in object size contribute approximately the same amount of night sky radiance. Therefore, a rapid rise in NSB is probable if space debris proliferates considerably. That remains true even if the rate of new launches slows or stops altogether, as the population of objects with smaller sizes will probably increase as a result of newly generated debris from existing satellites. There are no known effective mitigation options for the problem of elevated NSB other than drastic reductions in satellite launches or satellite brightness.

Satellites also pose a threat to astronomy outside the visual and near-infrared bands⁷. Radio astronomy is vulnerable to the direct and indirect emissions of radio energy from satellites, particularly out-of-band transmissions and sidelobes. Direct illumination of radio telescopes by satellites transmitting to the ground could damage or destroy sensitive radio detectors³.

Some scholars identify an environmental continuum between Earth and space that calls for a reconsideration of space ‘sustainability’ and define near-Earth space as part of the human environment¹⁵. A new ‘space environmentalism’ framework has been suggested to manage outer space sustainably and equitably^{1,4}.

Despite these efforts, many areas of astronomical research will be increasingly affected in a future in which LEO is ever more crowded with satellites. It is unrealistic and economically infeasible to move astronomy exclusively to space, and space telescopes can also be affected by orbital debris and satellite trails in images¹⁶. Some of the effects on astronomical images can be mitigated with software^{17,18}; however, this approach is expensive and imperfect. Not all effects, such as detector crosstalk during the passage of very bright objects through the field of view, can be removed. These destructive events risk opportunity loss that may impact scientific productivity of facilities and impede discovery. An overall increase in diffuse NSB requires longer exposure times to reach particular detection thresholds, which in turn increases the likelihood that streaks from resolved objects will affect images, and it increases the cost of data collection as we detail below. Unintended radio interference from satellites can hamper or prevent radio astronomy data collection.

These outcomes imply a diminution of future opportunities for discovery from both the ground and LEO. We next consider their associated costs.

The tangible costs of rising global NSB

For over a century, new ground-based astronomical observatories have been established in increasingly remote places, in part to evade

the influence of terrestrial skyglow. Even when such sites are identified and developed, the threat of skyglow remains ever present^{19,20}. Increased NSB resulting from proliferating space objects is a fundamentally new challenge to astronomy. In choosing new observatory sites, one cannot simply look for more distant locations because increasing global NSB will be experienced planet-wide. Here we attempt to quantify what this means for the future of terrestrial and LEO-based astronomy.

Loss of data and discovery for professional astronomy

As NSB rises, the exposure time required to reach any particular signal-to-noise ratio (S/N) rises concomitantly. In cases where exposure times are held fixed, such as in areal sky surveys, a brighter night sky corresponds to a brighter detection limit. As a result, fainter objects will be missed, which will directly diminish the pace and impact of astronomical discovery. It is impossible to obtain a reliable monetary cost estimate of the loss of opportunity, particularly if we miss rare astrophysical phenomena because satellites interfered with observations. An example with distinct and potentially severe social consequences is the detection of near-Earth objects that represent a high risk of colliding with our planet. For example, the Chelyabinsk bolide—the largest known natural object to have entered Earth’s atmosphere since the 1908 Tunguska event—was undiscovered at the time of its entry into the Earth’s atmosphere in 2013 in part due to its position on the sky, near the Sun, in the days and hours before impact²¹. Hazardous near-Earth objects that sky surveys may fail to detect often first appear in our skies in the twilight hours around sunrise and sunset, times when satellites and space debris are most likely to interfere with observations².

This could have profound consequences for high-profile terrestrial facilities in the coming decades. For example, the Vera Rubin Observatory estimates that if the SpaceX Starlink constellation achieved its full design buildout of 42,000 satellites, as many as 30% of all Legacy Survey of Space and Time (LSST) images would contain at least one satellite trail²². The Vera Rubin Observatory expects that software mitigations will not effectively deal with all systematic effects and the resulting spurious event triggers, especially at low brightness. Similarly, for a full buildout of Starlink and OneWeb’s proposed ~48,000 satellites, every 30 s exposure on the Large Magellanic Cloud during the Southern Hemisphere summer months is expected to contain at least one satellite trail².

The impact of brief satellite glints, however, is largely unknown. Such glints are poorly characterized and will almost certainly impact astronomical studies, including some of the fastest-growing research areas, such as time-domain astronomy. One recent example is the discovery of a faint, apparently transient object thought to represent an exotic astrophysical phenomenon²³, later suggested to have been caused by satellite interference with the observation^{24,25}. These instances could soon become commonplace.

Monetary cost of longer integration times with elevated diffuse NSB

If, as models predict, the night sky becomes brighter as a result of the proliferation of space objects, then progressively longer integration times will be required to reach any given S/N threshold. For any science programme with a defined S/N requirement, rising NSB inevitably imposes a loss of efficiency that can be interpreted as an increased financial burden.

As an example, we consider the potential efficiency loss for the LSST. The survey has specified a number of quality metrics tied to S/N. Following the LSST 5σ detection depth quoted by Ivezić et al.²⁶, we examined the effect of changing NSB on the LSST point-source efficiency near the single-visit detection threshold for a fixed exposure time. Figure 1 shows the expected S/N as a function of the brightness of the sky adjacent to a science target.




Fig. 1 | S/N as a function of background NSB for point sources in the Vera Rubin Observatory's LSST. The S/N as a function of background NSB is shown around the single-visit detection limit (magnitude $r = +24.5$ mag) and an exposure time of 20 s from predictions based on the model of Ivezić et al.²⁶. Four sources are plotted with varying r as indicated. Note that the x-axis values are reversed such that the night sky becomes brighter moving left to right along the axis.

Given the projection by Kocifaj et al. that zenith luminance due to space objects may approach a maximum value of $25 \mu\text{cd m}^{-2}$ for solar depression angles of $\sim 23^\circ$ by 2030¹², the zenith might by then be $\sim 12\%$ brighter than an assumed pristine night sky ($\sim 22.0 m_V \text{ arcsec}^{-2}$, or $\sim 2 \times 10^{-4} \text{ cd m}^{-2}$). For a sky-dominated observation near the point-source detection threshold, this 12% increase in NSB directly translates to a 12% increase in exposure time required to reach the same S/N as under pristine conditions. For brighter objects, this translation becomes less linear and the value of this factor decreases until it reaches unity (that is, until the signal is object-dominated, rather than sky-dominated).

Ivezić et al. note that for galaxies with a magnitude i around +25, LSST photometry is expected to achieve a root-mean-square accuracy $\sigma/(1+z)$ of 2% over the range $0.3 < z < 3.0$ at a S/N of 20 (ref. ²⁶). To reach that S/N in a single observation would take about 12% more exposure time for LEO space objects contributing $25 \mu\text{cd m}^{-2}$ of NSB versus a scenario in which they contributed no NSB. It is important to note that this is a small effect compared with the contribution of the natural night airglow, which is typically three times brighter under quiescent conditions²⁷. To the extent that observations of faint objects are almost always limited by the total NSB, any increase above the natural background set by airglow and other phenomena necessarily requires a longer exposure time to achieve a given S/N. The contribution of space objects to the diffuse NSB limiting such observations is non-negligible.

For sky-limited photometric measurements of point sources, the exposure time (t_{exp}) required to reach a particular S/N is proportional to the square of that ratio:

$$t_{\text{exp}} \simeq \left(\frac{S/N}{F_{\text{obj}}} \right)^2 n_{\text{pix}} F_{\text{sky}},$$

in which F_{obj} is the object flux (in photons s^{-1}), n_{pix} is the number of detector pixels over which the point spread function is sampled and F_{sky} is the sky background flux (in photons $\text{s}^{-1} \text{ pixel}^{-1}$). For any particular combination of S/N, F_{obj} and n_{pix} , the ratio of t_{exp} at any two different values of B_{sky} is constant (Fig. 2):

$$\frac{t_{\text{exp}}(B_2)}{t_{\text{exp}}(B_1)} = 10^{-0.4(B_1 - B_2)},$$

where B_1 and B_2 are arbitrary NSB values (in mag arcsec^{-2}). Furthermore, that ratio is very nearly identical to the ratio of B_{sky} to an assumed 'pristine' value at the zenith. In other words, if the NSB increases by a factor of M over pristine conditions, then the exposure time required to reach an arbitrary S/N at that NSB also increases directly by about a factor of M . We acknowledge that these estimates apply only to single observations and are not representative of the cumulative result of many observations. Nevertheless, given that telescope time is represented by a constant cost per unit at most facilities, we draw attention to the sobering reality that the financial cost of acquiring data under brightening night sky conditions will scale with increasing NSB. Moreover, as annual allocations of observing time are usually a fixed quantity, if each target observing programme requires more time with increasing NSB, fewer scientific programmes can be completed.

Estimating the financial cost of this effect is difficult given that the relationship between survey benchmarks and operations costs is complex. The base capital cost of LSST is estimated by the US National Science Foundation to be US\$473 million²⁸. The estimated annual operation costs add up to US\$290 million over the 10 yr lifetime of the baseline survey. Assuming a 1:1 correspondence between survey duration and financial cost, an increase in the duration of 12% (1.2 yr) would correspond to an additional US\$34.8 million in total project costs.

Not all objects contribute to the diffuse brightness of the sky when viewed through a telescope. Bright objects appear in images as discernible streaks. To appear as a streak, the irradiance of an object image on the detector plane must be sufficient to provide a detectable signal above the background and noise levels. The image on the detector is the two-dimensional convolution of the geometrical image of the object (whose typical linear dimension is proportional to the object size and inversely proportional to the object distance) and the overall point spread function of the optical system (determined by the optical response of the telescope and the relevant atmospheric effects averaged within the effective exposure time). The signal of the object on a pixel is proportional to the irradiance of its image, the pixel area and the effective exposure time (the time during which the moving image of the object illuminates the pixel's active surface). The radiance contribution of the objects detectable as individual streaks should be subtracted from the total diffuse radiance mentioned above. For facilities such as the Rubin Observatory, whose telescope and camera should detect streaks from objects in orbit as small as ~ 5 cm, this would remove about three-eighths of reflected sunlight from the maximum $25 \mu\text{cd m}^{-2}$ that would otherwise elevate the natural diffuse sky background. In practical terms this means that in these conditions the diffuse background will be $(5/8) \times 25 \mu\text{cd m}^{-2} = \sim 15 \mu\text{cd m}^{-2}$; that is, a 7.5% increase over the assumed natural reference level. Considering this effect alone, the LSST project would therefore require 7.5% more time, and at equal yearly cost this would be an overcharge of $0.075 \times \text{US\$290 million} = \text{US\$21.8 million}$.

Shortcomings of the current modelling regime

We do not have a way to easily predict or calculate the time evolution of the number of space objects as a function of their sizes and orbit altitude distribution. Instead, at present we have to live with the aftermath of unplanned events or debris cascades. As Kocifaj et al. point out, the most complete picture of the cumulative number of objects in LEO larger than a given size dates to the mid-1990s¹². The international scientific community is restricted to publicly available, civilian data typically derived from limited radar measurements²⁹ or space environment exposure experiments³⁰; we speculate that militaries and private companies may have access to more detailed information. Given the




Fig. 2 | Exposure time required to reach a S/N of 10. Exposure time required to reach S/N = 10, plotted on a log₁₀ scale, as a function of background NSB for point sources in the LSST. The range of target magnitudes shown is the same as in Fig. 1.

high relative velocities involved, the still largely unknown population of millimetre-sized and smaller objects poses the greatest threat to space hardware. This hazard is expected to be consequential for spacecraft in large satellite constellations³¹.

Furthermore, there is great variety—and therefore uncertainty—in the number and type of satellites proposed for near-term orbital deployment. Recently launched objects are observed to vary substantially in terms of their optical properties, both between individual objects and for a single object over time^{32–34}. Individual debris fragments are even less well understood, although their properties in aggregate can be estimated reasonably well. But the single greatest uncertain factor in the modelling realm, and most challenging to predict, is what exactly will dominate the changing NSB. We do not yet know whether it will be the over 400,000 satellites already planned for launch in the coming decade³⁵, a cascade of fragments from accidental collisions, debris from reckless acts undertaken in LEO by state actors or some combination of these.

The broader landscape of consequences

Loss of dark night skies

Brightening night skies impact not only astronomy but also the human experience of viewing the night sky. We attempt to quantify this by comparing viewing experiences under various NSB conditions. There are a number of ways of characterizing the brightness of the night sky using both qualitative and quantitative metrics³⁶. In Table 1 we compare several key visual night sky quality indicators for a series of levels of NSB above the pristine brightness assumed above, assuming a dark-adapted observer viewing the night sky under typical clear-sky conditions at sea level. Brightening night skies also make the Milky Way markedly harder to see from anywhere on Earth and diminish views of night sky phenomena such as night airglow, weak aurorae and faint meteors. Doubling the zenith brightness relative to an unpolluted night sky reduces the number of visible stars by roughly 30% and reduces the number of visually observable meteors by a factor of up to one-half.

Impacts on human heritage and culture

The night sky transcends science or utility; it is equally a source of inspiration, connection to nature and recreation. For some cultures, sky traditions are a prominent aspect of their social customs, cultural traditions and religious beliefs. As such, it represents a form of intangible

Table 1 | Visual night sky quality indicators as a function of rising zenith brightness

Zenith brightness (m_v arcsec ⁻²)	Increase over pristine night sky (%)	Limiting visual magnitude	Bortle scale	Number of visible stars per night
22.00	0	+6.3	1	3,500
21.99	1	+6.3	1	3,500
21.97	3	+6.2	1	3,150
21.95	5	+6.2	1	3,150
21.90	10	+6.2	1	3,150
21.76	25	+6.2	2	3,150
21.56	50	+6.1	3	2,800
21.25	100	+6.0	4	2,540

Brightness values vary from pristine conditions to those at which the zenith is twice as bright in comparison. The limiting visual magnitude was calculated using equation (53) in ref. ⁶⁴ for a typical observer (field factor $F=2$). The Bortle scale is a qualitative scale that ranks night skies from 1 (pristine) to 9 (most light-polluted) on the basis of a series of visual criteria⁶⁵.

human heritage that deserves intentional preservation and safeguarding for future generations^{37,38}.

Our calculations indicate that the brighter stars and constellations often utilized in navigational aspects of cultural sky traditions, including wayfinding, will remain visible even for the more extreme scenarios we consider here. However, the anticipated rise in NSB adds to the contribution of terrestrial skyglow and will wash out fainter stars and the Milky Way. This tends to diminish the visibility of the dark clouds seen in silhouette against the Milky Way that play an important role in many sky cultural traditions in the Southern Hemisphere³⁹. Fainter objects, such as nebulae, star clusters and dimmer groups of stars, are also often key elements of teachings in various Indigenous communities, as are observations of the heliacal rising of various celestial objects⁴⁰. These are all likely to be impacted by rising NSB. In addition, satellites visible as moving points of light alter the appearance of the night sky; for some communities, this is seen to ‘interrupt’ their ‘relationship with the stars and ceremonial ways of connecting with them’⁴¹. While various mitigation techniques may help address some satellite constellation impacts on professional astronomy, real-time observations—and thus living sky traditions—will be adversely impacted by visible satellites and rising NSB.

Implications for equity, inclusion and accessibility in astronomy

The roiling waves of the COVID-19 pandemic, climate change and global economic turbulence in recent years have resonantly combined in unpredictable ways to jeopardize the lives and livelihoods of the world’s most vulnerable populations. Serious adverse health effects also arise from disproportionate light and noise pollution in these communities⁴². Rising NSB has a documented impact on the health of humans and the well-being of broader ecological systems⁴¹; this is especially concerning given its inescapable, planet-wide nature. Reduced visibility of, for example, the Milky Way impacts the migratory patterns of many creatures^{41,43–45}; Lawrence et al. point out that most circadian rhythms are apparently controlled by diffuse ambient light and not by moving point sources¹.

For professional astronomy, in this time of shrinking budgets and fewer grant dollars in a zero-sum game, the competition for observing time on ground-based telescopes and facilities will become even more highly contested than it is at present, especially if longer exposures are needed for sources in a sky with greater radiance from space objects. Observing time, grants and awards, like all privilege, tend to accumulate in select academic lineages and institutional classes; longer integration times are likely to even further concentrate this privilege

within a shrinking circle of institutions. In such a professional environment, the recruitment, retention and promotion of under-represented and marginalized groups in astronomy faces increasing challenges, at a time when our field's future workforce is already confronting a radically altered landscape of professional and research opportunities³⁸.

The COVID-19 pandemic has revealed many pre-existing systemic conditions that have widened socioeconomic and learning gaps, including access to affordable global broadband⁴⁶. Very few would argue against the dire need for expanded access to broadband; our capacity to conduct teaching and research in astronomy, as well as a competitive future workforce in astronomy and other fields, depends on this. While the commercial space industry has argued for the benefits of providing broadband Internet from space, we note that, so far, no company has ever demonstrated a satellite broadband business model that is both profitable and sustainable. The latter is often mentioned prominently, but the driving factor is the former: profit. Rawls et al. questioned the motive often stated by commercial space companies that have proposed launching megaconstellations, which is to provide broadband to underserved populations globally; instead, they found that “the Internet service offered by these satellites will primarily target populations where it is unaffordable, not needed, or both”⁴⁷.

Furthermore, the effective consolidation of control of LEO space for communications by a handful of private companies, or by a small number of privileged industrialized states, risks diminishing the equity, inclusion and accessibility to broadband communications. We note that there are technological alternatives to broadband from orbit, such as fibre-optic transmission and the latest generation of terrestrial wireless data networks, that could achieve the same result without the scientific and business risks attendant to the launch and operation of satellite megaconstellations. In addition, distributed broadband ground communication networks are somewhat more difficult to oligopolize.

Potential gains and risks of mitigation

Mitigation of the threats to astronomy described above largely falls into two categories: (1) modifying satellite and satellite constellation designs, and (2) back-correction or restoration of astronomical data impacted by satellites and space debris^{4,48}. The former approach was taken by SpaceX in 2020 after astronomers first raised the prospect of harm to astronomy research resulting from its Starlink satellites. Its DarkSat and VisorSat designs were intended to reduce the brightness of Starlink objects. Some observers measured decreases in on-station brightnesses among objects of these designs by up to 1.5 mag compared with the original Starlink design^{33,49–52}, while others saw essentially no change⁵³ or even brightness increases⁵⁴. However, both mitigations were abandoned for engineering reasons, and neither achieved the goal of reducing the brightness of first-generation Starlink objects at station to below the threshold of unaided-eye observability. Although this problem remains unsolved, SpaceX and its competitors are planning to launch new satellites of unknown brightness.

Design solutions may reduce streak brightnesses but are not expected to have any meaningful effect on rising diffuse NSB as most of this increase in NSB is due to small debris. Design solutions will also have little or no effect on the potential for debris-generating collisions. In addition, mitigating solutions at one bandpass may be a problem for other bandpasses—for example, optically darker objects often radiate more brightly in infrared and submillimetre wavelengths, which creates interference with ground-based observations at those wavelengths. It also remains a problem that many operators are not forthcoming with details of their satellite designs, including materials, albedos, bidirectional reflectance distribution functions and other parameters, because they protect them as intellectual property and governments generally do not require their public disclosure. That often leaves astronomers to invest their taxpayer-funded time and resources into inferring these properties through observations and modelling. We note that some operators have been more collaborative

with astronomers and with data sharing; for example, SpaceX recently released a report on their attempts to decrease satellite reflectivity⁵⁵. Notable concerns remain about the brightness of next-generation Starlink satellites, as the extraordinarily bright AST SpaceMobile Blue-Walker 3 prototype satellite proved quite recently³⁴.

Data treatments can improve the quality of images impacted by trails and glints and recover some science pixels. But saturated pixels are lost, and they can induce crosstalk between detector amplifiers. In their simulations of LSST data, Tyson et al. found that they could effectively remove the influence of nonlinear image artefacts induced by satellite trails at a maintained brightness of magnitude +7 or fainter, although they expected that “systematic errors that may impact data analysis and limit some science” would remain after correction¹⁷.

Perhaps the mitigation option least palatable to the commercial space industry, but one that governments may certainly impose, is to simply launch fewer satellites into near-Earth space. It is the only solution that simultaneously tackles all the problems we describe here. With any such plan for satellite launch reduction should come the intent to responsibly de-orbit those already launched at end-of-mission to minimize ongoing collision risks. Satellite operators should also be held to broader standards of responsible reduction, and ideally elimination, of debris associated with all stages of satellite operation including launch and disposal at end-of-mission⁵⁶. While US policy was recently amended to reduce the disposal time from within 25 years to within 5 years of end-of-mission as a condition of licensing spacecraft operations⁵⁷, not all jurisdictions make the same demand for objects launched from their territories. On-orbit debris removal has been proposed^{58,59}, but so far no one has demonstrated this successfully in practice. The same is true for perfect collision avoidance through various space situational awareness schemes⁶⁰. It is a far more reliable course of action to avoid debris cascades in the first place, which necessarily entails launching fewer objects and reducing the number of objects already in orbit⁶¹.

This brewing crisis in LEO has powerful lessons for our shared future in near-Earth space. There are still opportunities to get ahead of the problem elsewhere. For example, now is the time to consider a future in which astronomical observations performed from the lunar surface may be similarly affected by a growing swarm of space objects orbiting the Moon⁶². There are fewer legal restrictions on the use of cislunar space, and the race to occupy that space is already on⁶³.

Looking ahead

The lack of coordinated and effective global policy, regulation and oversight among spacefaring nations and space actors has led to the prospect of hundreds of thousands of satellites planned for launch in the coming decade, without attendant conditions on effective mitigation strategies or environmental self-assessment as a condition of licensing. Despite a narrative of democratizing space and delivering affordable global broadband, it is a model that prioritizes urgency, privatized benefits and short-term goals over real sustainability and the public interest. This also ignores our shared ancestry and heritage in space.

In this Perspective, we have emphasized the potential consequences of a global increase in NSB given the rising numbers of space objects in LEO. Unlike satellite streaks and glints that hold some options for back-corrections, this is an inescapable, planet-wide phenomenon that will affect professional and casual observations of the skies, as well as myriad biological systems. We have attempted some estimates of the loss of professional observing time and unaided-eye observations of astronomical phenomena and pointed to the work of other teams on space environmentalism and the risks from space debris.

We also recognize that the broader loss of dark skies for humanity is essentially incalculable, given the many ways we have connected to the night sky for millennia. The night sky needs preserving and defending for future generations who may not know what we have today, akin to the analogous situation of disappearing rainforests and glaciers. But

unlike these examples of the toll of climate change and human activity, we are still in the early stages of changing the night sky and the environment of space for future generations. Many of the consequences of this remain unknown.

To paraphrase the Senegalese conservationist Baba Dioum, we will conserve only what we value, value only what we know and know only what we are taught. We conclude by offering the hope that we can still deliver on the promise of human activity and development in and from space by proceeding sustainably for all stakeholders and preserving space as a resource for future generations. We must become proactive now or risk irreparable harm to astronomy resulting from the current rush to capitalize on the exploitation of LEO and other space resources.

References

- Lawrence, A. et al. The case for space environmentalism. *Nat. Astron.* **6**, 428–435 (2022).
- Walker, C. et al. *Impact of Satellite Constellations on Optical Astronomy and Recommendations Toward Mitigations* (NOIRLab, NSF, 2020); <https://aas.org/sites/default/files/2020-08/SATCON1-Report.pdf>
- Walker, C. et al. *Dark and Quiet Skies for Science and Society* (International Astronomical Union, 2020); <https://www.iau.org/static/publications/dqskies-book-29-12-20.pdf>
- Hall, J. et al. *Report of the SATCON2 Workshop 12-16 July 2021: Executive Summary* (NOIRLab, NSF, 2021); <https://noirlab.edu/public/media/archives/techdocs/pdf/techdoc031.pdf>
- Walker, C. & Benvenuti, P. (eds) *Dark and Quiet Skies for Science and Society II: Working Group Reports* (NOIRLab, NSF, 2022); <https://noirlab.edu/public/media/archives/techdocs/pdf/techdoc021.pdf>
- Rawls, M. & Bektešević, D. *Trailblazer* (International Astronomical Union Centre for the Protection of the Dark and Quiet Sky from Satellite Constellation Interference, 2022); <https://trailblazer.dirac.dev/>
- McDowell, J. The low earth orbit satellite population and impacts of the SpaceX Starlink constellation. *Astrophys. J.* **892**, L36 (2020).
- Hainaut, O. & Williams, A. Impact of satellite constellations on astronomical observations with ESO telescopes in the visible and infrared domains. *Astron. Astrophys.* **636**, A121 (2020).
- Ragazzoni, R. The surface brightness of megaconstellation satellite trails on large telescopes. *Publ. Astron. Soc. Pac.* **132**, 114502 (2020).
- Mróz, P. et al. Impact of the SpaceX Starlink satellites on the Zwicky Transient Facility Survey observations. *Astrophys. J. Lett.* **924**, L30 (2022).
- Lawler, S., Boley, A. & Rein, H. Visibility predictions for near-future satellite megaconstellations: latitudes near 50° will experience the worst light pollution. *Astron. J.* **163**, 21 (2021).
- Kocifaj, M., Kundracik, F., Barentine, J. & Bará, S. The proliferation of space objects is a rapidly increasing source of artificial night sky brightness. *Mon. Not. R. Astron. Soc. Lett.* **504**, L40–L44 (2021).
- Bassa, C., Hainaut, O. & Galadí-Enríquez, D. Analytical simulations of the effect of satellite constellations on optical and near-infrared observations. *Astron. Astrophys.* **657**, A75 (2022).
- Space Environment Report* (European Space Agency, 2022); <https://sdup.esoc.esa.int/discosweb/statistics/>
- Yap, X.-S. & Truffer, B. Contouring ‘earth-space sustainability’. *Environ. Innov. Soc. Transit.* **44**, 185–193 (2022).
- Kruk, S. et al. The impact of satellite trails on Hubble Space Telescope observations. In *From Measurements to Understanding: MASTER Modelling Workshop2–4 March 2021* (eds Oikonomidou, X. et al.) (European Space Agency, 2021); https://indico.esa.int/event/370/contributions/5925/attachments/4238/6337/Sandor_Kruk_The_impact_of_satellite_trails_on_Hubble_observations_compressed.pdf
- Tyson, J. A. et al. Mitigation of LEO satellite brightness and trail effects on the Rubin Observatory LSST. *Astron. J.* **160**, 226 (2020).
- Hasan, I., Tyson, J. A., Saunders, C., & Xin, B. Mitigating satellite trails: A study of residual light after masking. *Astron. Comput.* **39**, 100584 (2022).
- Green, R., Luginbuhl, C., Wainscoat, R. & Duriscoe, D. The growing threat of light pollution to ground-based observatories. *Astron. Astrophys. Rev.* **30**, 1 (2022).
- Falchi, F. et al. Light pollution indicators for all the major astronomical observatories. *Mon. Not. R. Astron. Soc.* **519**, 26–33 (2022).
- Borovička, J. et al. The trajectory, structure and origin of the Chelyabinsk asteroidal impactor. *Nature* **503**, 235–237 (2013).
- Impact of Satellite Constellations* (Vera Rubin Observatory, 2022); <https://www.lsst.org/content/lsst-statement-regarding-increased-deployment-satellite-constellations>
- Jiang, L. et al. A possible bright ultraviolet flash from a galaxy at redshift $z \approx 11$. *Nat. Astron.* **5**, 262–267 (2020).
- Nir, G., Ofek, E. & Gal-Yam, A. The GN-z11-flash event can be a satellite glint. *Res. Not. Am. Astron. Soc.* **5**, 27 (2021).
- Michałowski, M., Kamiński, K., Kamińska, M. & Wnuk, E. GN-z11-flash from a man-made satellite not a gamma-ray burst at redshift 11. *Nat. Astron.* **5**, 995–997 (2021).
- Ivezić, Ž. et al. LSST: from science drivers to reference design and anticipated data products. *Astrophys. J.* **873**, 111 (2019).
- Sternberg, J. & Ingham, M. Observations of the airglow continuum. *Mon. Not. R. Astron. Soc.* **159**, 1–20 (1972).
- NSF FY 2021 Budget Request to Congress* (NSF, 2020); https://nsf.gov/about/budget/fy2021/pdf/34g_fy2021.pdf
- Muntoni, G., Montisci, G., Pisanu, T., Andronico, P. & Valente, G. Crowded space: a review on radar measurements for space debris monitoring and tracking. *Appl. Sci.* **11**, 1364 (2021).
- Mandeville, J. & Berthoud, L. From LDEF to EURECA: orbital debris and meteoroids in low earth orbit. *Adv. Space Res.* **16**, 67–72 (1995).
- Le May, S., Gehly, S., Carter, B. & Flegel, S. Space debris collision probability analysis for proposed global broadband constellations. *Acta Astronaut.* **151**, 445–455 (2018).
- Mallama, A. A bidirectional reflectance distribution function for VisorSat calibrated with 10,628 magnitudes from the MMT-9 database. Preprint at <https://arxiv.org/abs/2109.07345> (2021).
- Tregloan-Reed, J. et al. Optical-to-NIR magnitude measurements of the Starlink LEO darksat satellite and effectiveness of the darkening treatment. *Astron. Astrophys.* **647**, A54 (2021).
- Mallama, A., Cole, R., Harrington, S. & Maley, P. Visual magnitude of the BlueWalker 3 satellite. Preprint at <https://arxiv.org/abs/2211.09811> (2022).
- McDowell, J. Enormous (‘mega’) satellite constellations. *Jonathan’s Space Report* <https://planet4589.org/space/con/conlist.html> (2023).
- Barentine, J. Night sky brightness measurement, quality assessment and monitoring. *Nat. Astron.* **6**, 1120–1132 (2022).
- Ruggles, C. Astronomy and world heritage. *Proc. Int. Astron. Union* **6**, 12–17 (2010).
- Venkatesan, A., Lowenthal, J., Prem, P. & Vidaurre, M. The impact of satellite constellations on space as an ancestral global commons. *Nat. Astron.* **4**, 1043–1048 (2020).
- Gullberg, S. et al. A cultural comparison of the ‘dark constellations’ in the Milky Way. *J. Astron. Hist. Herit.* **23**, 390–404 (2020).
- Lee, A. et al. Indigenous astronomy: best practices and protocols for including Indigenous astronomy in the planetarium setting. In *Proceedings of the 25th International Planetarium Society Conference 3–7 August 2020* (ed. Smith, D. W.) 69–77. (International Planetarium Society, 2020) https://cdn.ymaws.com/www.ips-planetarium.org/resource/resmgr/vcon2020/papers/IPS_2020.pdf

41. Venkatesan, A. et al. in *SATCON2 Working Group Reports* (eds Hall, J. & Walker, C) 102 (NOIRLab, 2021).
42. Nadybal, S., Collins, T. & Grineski, S. Light pollution inequities in the continental United States: a distributive environmental justice analysis. *Environ. Res.* **189**, 109959 (2020).
43. Foster, J., Smolka, J., Nilsson, D.-E. & Dacke, M. How animals follow the stars. *Proc. R. Soc. B* **285**, 20172322 (2018).
44. Stone, E., Harris, S. & Jones, G. Impacts of artificial lighting on bats: a review of challenges and solutions. *Mamm. Biol.* **80**, 213–219 (2015).
45. Pakhomov, A., Anashina, A. & Chernetsov, N. Further evidence of a time-independent stellar compass in a night-migrating songbird. *Behav. Ecol. Sociobiol.* **71**, 48 (2017).
46. Patrick, S., Grissom, J., Woods, S. & Newsome, U. Broadband access, district policy, and student opportunities for remote learning during COVID-19 school closures. *AERA Open* **7**, 233285842110642 (2021).
47. Rawls, M. et al. Satellite constellation internet affordability and need. *Res. Notes Am. Astron. Soc.* **4**, 189 (2020).
48. Massey, R., Lucatello, S. & Benvenuti, P. The challenge of satellite megaconstellations. *Nat. Astron.* **4**, 1022–1023 (2020).
49. Lalbakhsh, A. et al. Darkening low-Earth orbit satellite constellations: a review. *IEEE Access* **10**, 24383–24394 (2022).
50. Cole, R. Measurement of the brightness of the Starlink spacecraft named ‘DARKSAT’. *Res. Notes Am. Astron. Soc.* **4**, 42 (2020).
51. Tregloan-Reed, J. et al. First observations and magnitude measurement of Starlink’s Darksat. *Astron. Astrophys.* **637**, L1 (2020).
52. Mallama, A. Starlink satellite brightness—characterized from 100,000 visible light magnitudes. Preprint at <https://arxiv.org/abs/2111.09735> (2021).
53. Cole, R. A sky brightness model for the Starlink ‘Visorsat’ spacecraft. *Res. Notes Am. Astron. Soc.* **4**, 182 (2020).
54. Horiuchi, T., Hanayama, H. & Ohishi, M. Simultaneous multicolor observations of Starlink’s Darksat by the Murikabushi Telescope with MITSuME. *Astrophys. J.* **905**, 3 (2020).
55. *Brightness Mitigation Best Practices for Satellite Operators* (SpaceX, 2022); <https://api.starlink.com/public-files/BrightnessMitigationBestPracticesSatelliteOperators.pdf>
56. Byers, M., Wright, E., Boley, A. & Byers, C. Unnecessary risks created by uncontrolled rocket reentries. *Nat. Astron.* **6**, 1093–1097 (2022).
57. *Space Innovation IB Docket No. 22-271 Mitigation of Orbital Debris in the New Space Age IB Docket No. 18-313* (Federal Communications Commission, 2022); <https://www.fcc.gov/document/fcc-adopts-new-5-year-rule-deorbiting-satellites-0>
58. Shan, M., Guo, J. & Gill, E. Review and comparison of active space debris capturing and removal methods. *Prog. Aerosp. Sci.* **80**, 18–32 (2016).
59. Mark, C. & Kamath, S. Review of active space debris removal methods. *Space Policy* **47**, 194–206 (2019).
60. Rybus, T. Obstacle avoidance in space robotics: review of major challenges and proposed solutions. *Prog. Aerosp. Sci.* **101**, 31–48 (2018).
61. Lewis, H. Understanding long-term orbital debris population dynamics. *J. Space Saf. Eng.* **7**, 164–170 (2020).
62. Silk, J., Crawford, I., Elvis, M. & Zarnecki, J. Astronomy from the Moon: the next decades. *Phil. Trans. R. Soc. A* **379**, 20190560 (2020).
63. Spudis, P. in *Toward a Theory of Spacepower* (eds Lutes, C. D. & Hays, P. L.) 241–251 (Institute For National Strategic Studies, National Defense Univ. Press, 2011).
64. Crumey, A. Human contrast threshold and astronomical visibility. *Mon. Not. R. Astron. Soc.* **442**, 2600–2619 (2014).
65. Bortle, J. Introducing the Bortle Dark-Sky Scale. *Sky Telescope* **101**, 126–129 (2001).

Acknowledgements

M.K. acknowledges support from the Slovak Research and Development Agency under contract number APVV-18-0014. A.V. gratefully acknowledges support from the University of San Francisco Faculty Development Fund. The views expressed in this Perspective do not necessarily represent the positions of either the American Astronomical Society or the International Astronomical Union.

Competing interests

J.C.B., A.V., J.H. and J.L. are unpaid members of committees of the American Astronomical Society and International Astronomical Union, whose scopes of concern include the topics covered by this Perspective.

Additional information

Correspondence should be addressed to John C. Barentine.

Peer review information *Nature Astronomy* thanks Piero Benvenuti and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2023