

---

4-5-2019

## Mercury Chemostratigraphy Across the Cambrian Series 2 – Series 3 Boundary: Evidence for Increased Volcanic Activity Coincident with Extinction?

L. E. Faggetter  
*University of Leeds*

P. B. Wignall  
*University of Leeds*

S. B. Pruss  
*Smith College, spruss@smith.edu*

D. S. Jones  
*Amherst College*

S. Grasby  
*Geological Survey of Canada Calgary*

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.smith.edu/geo\\_facpubs](https://scholarworks.smith.edu/geo_facpubs)



Part of the [Geology Commons](#)

---

### Recommended Citation

Faggetter, L. E.; Wignall, P. B.; Pruss, S. B.; Jones, D. S.; Grasby, S.; Widdowson, M.; and Newton, R. J., "Mercury Chemostratigraphy Across the Cambrian Series 2 – Series 3 Boundary: Evidence for Increased Volcanic Activity Coincident with Extinction?" (2019). Geosciences: Faculty Publications, Smith College, Northampton, MA.  
[https://scholarworks.smith.edu/geo\\_facpubs/117](https://scholarworks.smith.edu/geo_facpubs/117)

This Article has been accepted for inclusion in Geosciences: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact [scholarworks@smith.edu](mailto:scholarworks@smith.edu)

---

**Authors**

L. E. Faggetter, P. B. Wignall, S. B. Pruss, D. S. Jones, S. Grasby, M. Widdowson, and R. J. Newton

1 **Mercury chemostratigraphy across the Cambrian Series 2 – Series 3 boundary:**  
2 **evidence for increased volcanic activity coincident with extinction?**

3 Faggetter, L.E., Wignall, P.B., Pruss, S.B., Jones, D.S., Grasby, S., Widdowson, M.  
4 Newton, R.J.

5  
6 **ABSTRACT**

7 Flood basalt volcanism represented by the Kalkarindji Province (Australia) is  
8 temporally associated with a trilobite mass extinction at the Cambrian Series 2 –  
9 Series 3 boundary, providing one of the oldest potential links between volcanism and  
10 biotic crisis in the Phanerozoic. However, the relative timing of flood basalt volcanism  
11 (Kalkarindji Province, Australia) and the trilobite extinctions, first recorded in North  
12 America, is not known. Mercury (Hg) enrichment in the sedimentary record provides  
13 a potential proxy for volcanism which may facilitate improved chronologies of  
14 eruption and extinction. Here we report mercury records for three sections from mid-  
15 shelf strata of the Great Basin (western USA) that straddle the Series 2 – Series 3  
16 boundary. One section (Oak Springs Summit, NV) features a Hg enrichment at the  
17 start of the extinction interval, but mercury anomalies are also present at lower  
18 levels. These older anomalies may record either earlier phases of Kalkarindji  
19 volcanism, eruptions in other locations, or may be the result of sedimentary and/or  
20 diagenetic processes affecting the Hg record. In the Carrara Formation at Emigrant  
21 Pass, CA, the precise extinction horizon is not well defined, but a carbon isotope  
22 anomaly (the Redlichiid-Olenellid Extinction Carbon isotope Event; ROECE)  
23 provides a stratigraphic tie point to the Oak Springs Summit section. At Emigrant  
24 Pass, Hg enrichments precede the ROECE interval and are absent in the inferred  
25 extinction zone. The Pioche Formation at Ruin Wash, NV, lacks Hg enrichment at  
26 the extinction horizon but contains older enrichments. The inconsistent Hg records  
27 between the three sections demonstrate that factors controlling Hg accumulation and  
28 preservation in marine sedimentary environments are not yet fully understood. The  
29 effects of redox fluctuations may complicate one-to-one association of sedimentary  
30 Hg enrichments and massive volcanism at the Cambrian Series 2 – Series 3  
31 boundary and elsewhere in the geologic record.

32  
33 **1. INTRODUCTION**

34           The temporal connection between large igneous province (LIP) eruptions and  
35 Phanerozoic mass extinction events is well established, and the associated cause  
36 and effect linkage compelling (e.g. Wignall, 2001, 2015; Courtillot and Renne, 2003;  
37 Kravchinsky, 2012; Ernst & Youbi, 2017). Biotic change and carbon isotope  
38 excursions are an established feature of the Cambrian, but their relationship to LIP  
39 volcanism is poorly understood. In this context, the favoured candidate Cambrian LIP  
40 is the Kalkarindji Province (including the Antrim lavas) of northern and western  
41 Australia; this has been dated by a single zircon (Milliwindi dyke) to yield a U-Pb age  
42 of  $510.7 \pm 0.6$  Ma (Jourdan et al., 2014), demonstrating a close temporal relationship  
43 to the previously reported  $^{40}\text{Ar}/^{39}\text{Ar}$  date of  $507.5 \pm 1.6$  Ma determined for its  
44 extrusive portion (i.e. Antrim Lavas; Glass and Phillips, 2006). This LIP currently has  
45 a surface exposure of c. 425 000 km<sup>2</sup> in northern and central Australia (Veevers,  
46 2001), but was likely erupted over a much larger area; its scattered remnants could  
47 indicate a possible original extent of >2 million km<sup>2</sup> (e.g. Glass and Phillips, 2006,  
48 Jourdan et al., 2014). Accordingly, the province may thus be dimensionally  
49 comparable with other significant Phanerozoic LIPs (e.g. Columbia River Basalts),  
50 especially since it may also correlate with volcanics of similar age and/or  
51 composition preserved in the Tarim Block in NW China and the North China Block (Li  
52 et al., 1996; 2008), and the Sibumasu terrane preserved in current day Thailand and  
53 Myanmar (Zhu et al., 2012; Cocks and Torsvik, 2013).

54           Dating of the Kalkarindji province indicates emplacement close to the  
55 Cambrian Series 2 – Series 3 boundary (traditionally Lower – Middle), and thus  
56 potentially contemporaneous with the extinction of the redlichiid and olenellid  
57 trilobites (Palmer, 1998; Jourdan et al., 2014; Zhang et al., 2015). The extinction of  
58 the olenellids has been well studied in detailed sections located in the western USA,  
59 but locating the horizon that coincides with the Kalkarindji eruptions has hitherto  
60 been difficult due to a lack of an eruption proxy within ancient marine sediments.  
61 Temporal correlation between the Kalkarindji LIP emplacement and the Series 2 –  
62 Series 3 boundary is based upon a U-Pb zircon date of  $510.7 \pm 0.6$ Ma (Jourdan et  
63 al., 2014), and the provisional age of the Series 2 – Series 3 boundary (~509 Ma,  
64 Ogg et al., 2016). The association of this LIP with trilobite extinction at the Series 2 –  
65 Series 3 boundary is therefore inferred on the basis of this temporal correlation  
66 (Glass and Phillips, 2006; Hough et al., 2006; Jourdan et al., 2014).

67           Testing a causal link between the Kalkarindji and the Series 2 – Series 3  
68 trilobite extinction requires improved correlations between the volcanic event(s) and  
69 the trilobite extinction horizons. Recently, mercury (Hg) concentrations in the  
70 sedimentary record have provided a proxy for both regional and global volcanic  
71 activity (e.g. Schuster et al., 2002, albeit in ice-core, not sediments) and thus offers  
72 the potential to correlate the interval of extinction with evidence for volcanism in the  
73 same stratigraphic sections. The value of this technique has been demonstrated at  
74 several key mass extinction and oceanic anoxic events including those at the  
75 Ordovician–Silurian; latest Permian; Permian-Triassic; end-Triassic; early Jurassic  
76 and end-Cretaceous events (Sanei et al., 2012; Sial et al., 2013; 2014; Percival et  
77 al., 2015; Thibodeau et al., 2016; Font et al., 2016; Grasby et al., 2013; 2016;  
78 Bergquist, 2017; Gong et al., 2017; Jones et al., 2017; Percival et al., 2018). Here  
79 we present sedimentary Hg records for strata in which the trilobite extinction  
80 horizons have been constrained and evaluate the potential of the Kalkarindji LIP to  
81 produce this sedimentary geochemical record.

82           Volcanism represents a primary source of gaseous  $\text{Hg}^0$  to Earth's surface;  
83 unlike other volcanic trace metals, its long atmospheric residence time (0.5-2 years)  
84 permits hemispheric circulation and establishes its potential as a tracer for volcanism  
85 (Pyle and Mather, 2003; Percival et al., 2015, 2017). Atmospheric oxidation of  $\text{Hg}^0$   
86 by halogens, ozone and radicals forms reactive  $\text{Hg}^{2+}$ , a soluble ion which is  
87 deposited during precipitation (wet deposition). Horowitz et al. (2017) found that,  
88 during its residence in the troposphere, Hg is most effectively oxidised by bromine  
89 (Br), forming atmospheric HgBr complexes. As the largest source of atmospheric Br  
90 is organobromines - which are produced as a by-product of phytoplankton  
91 photosynthesis - the most effective oxidation and wet deposition of  $\text{Hg}^{2+}$  occurs  
92 above and, subsequently to, the marine realm (Horowitz et al., 2017). In modern  
93 oceans ~49% of marine Hg deposition occurs in tropical oceans due to the greater  
94 availability of productivity-driven organobromines and other oxidising radicals at  
95 these latitudes (Horowitz et al., 2017). Once in the marine realm  $\text{Hg}^{2+}$  forms  
96 complexes with clay minerals (Kongchum et al., 2011), organic matter (Benoit et al.,  
97 2001), and, in anoxic/euxinic conditions, it can be scavenged from seawater by  
98 sulphide complexes (Benoit et al., 1999). The effective oxidation of  $\text{Hg}^0$  by marine-  
99 derived organobromines and the complexing of Hg by organics and sulphides in the  
100 oceans establishes marine sediments as an efficient sink of atmospheric Hg (Benoit

101 et al., 1999; Emili et al., 2011; Horowitz et al., 2017). Therefore, marine siliciclastic  
102 and carbonate rocks can act as an important repository of Hg during times of  
103 heightened environmental loading (Percival et al., 2015; Grasby et al., 2016). It is  
104 also worth noting that once in the atmosphere particulate Hg can also be removed  
105 via “dry deposition”, a portion of which may make its way into the terrestrial realm  
106 (see Munthe et al., 2009 for discussion).

107         The Cambrian Period is marked by large oscillations of the inorganic carbon  
108 isotope record which, at times, coincided with intensified extinction rates (Brasier et  
109 al., 1994; Montañez et al., 2000; Zhu et al., 2006). At the Cambrian Series 2 – Series  
110 3 boundary, a negative  $\delta^{13}\text{C}$  excursion referred to as the Redlichiid – Olenellid  
111 Extinction Carbon isotope Excursion (ROECE) has been documented from Laurentia  
112 (Montañez, 2000; Faggetter et al., 2017), Gondwana (Schmid, 2017) and China (e.g.  
113 Fan et al., 2011; Wang et al., 2011; Chang et al., 2017; Ren et al., 2017). This  
114 coincides with major trilobite losses in both Gondwana and Laurentia (Montañez et  
115 al., 2000; Zhu et al., 2004; 2006; Faggetter et al., 2017; Ren et al., 2017). The Series  
116 2 – Series 3 boundary age (~509 Ma, Ogg et al., 2016) approximately coincides with  
117 the  $510.7 \pm 0.6$  Ma age of the Kalkarindji Province (Jourdan et al., 2014) but detailed  
118 correlation is lacking. We attempt to resolve this issue by examining sedimentary Hg  
119 concentrations in Cambrian Series 2 – Series 3 sections of the western USA. We  
120 have analysed Hg concentrations and Hg/ total organic carbon content (TOC) from  
121 two formations and three sections in the western Great Basin: Carrara Formation,  
122 Emigrant Pass (Death Valley, CA) and the Pioche Formation at Oak Springs Summit  
123 and Ruin Wash (Lincoln County, NV). These sections have an established  
124 biostratigraphic framework, and also a record of trilobite extinction at Oak Springs  
125 Summit and Ruin Wash (Webster et al., 2008; Moore and Lieberman, 2009).

126

## 127         **2. STUDY AREA**

128         Cambrian successions of the western Great Basin (USA) constitute the  
129 primary field locations of this study. Following the breakup of the supercontinent  
130 Rodinia in the late Neoproterozoic, a broad, equatorial clastic shelf developed on the  
131 rapidly subsiding Laurentian margin (Prave, 1991; Howley et al., 2006). During  
132 Cambrian Series 2, deposition in the Great Basin was on a broad shelf located on  
133 the north-western margin of Laurentia (Fig. 1). Clastic deposition was waning and, by  
134 Series 3, it had been replaced by carbonate production, resulting in the formation of

135 an extensive carbonate shelf (Fig. 1; Howley et al., 2006; Landing, 2012). We  
136 present data from two formations spanning the Cambrian Series 2 – Series 3  
137 boundary in the western Great Basin. The first section records the Carrara Formation  
138 of Death Valley, exposed at Emigrant Pass, California (Fig. 2). The second section is  
139 the Pioche Formation of eastern Nevada exposed at Oak Springs Summit (Fig. 3),  
140 and a third section at Ruin Wash, also recording the Pioche Formation (Fig. 4). Oak  
141 Springs Summit and Ruin Wash are close to each other (~20kms apart), whilst  
142 Emigrant Pass is ~225km south-west of these two locations (Fig. 1). Both formations  
143 comprise alternating siliciclastic- and limestone-dominated units (Merriam and  
144 Palmer, 1964; Palmer and Halley, 1979; Faggetter et al., 2017). At Oak Springs  
145 Summit and Ruin Wash, the Pioche Formation records the abrupt extinction of the  
146 olenellid trilobites, making these two sections candidates for paired Hg-  
147 biostratigraphic studies. At Emigrant Pass there is a paucity of trilobite fossils, but  
148  $\delta^{13}\text{C}$  correlation, based on the record of the ROECE (e.g., Zhu et al., 2004), allows  
149 an extinction interval to be inferred at the boundary between the *Olenellus* and  
150 *Eokochaspis nodosa* biozones in the mid Pyramid Shale (Fig. 2; Faggetter et al.,  
151 2017). As a further stratigraphic tie point between the two formations, the extinction  
152 horizon within the Pioche Formation represents the top of the *Olenellus* biozone  
153 (Palmer, 1998; Sundberg and McCollum, 2000).

154

### 155 **3. METHODS**

156 We analysed total Hg content in whole-rock powders from 93 samples taken  
157 from the three Series 2 – Series 3 boundary sections in the western US (Figs. 2, 3,  
158 4). The samples from Emigrant Pass and Oak Springs Summit were run at the  
159 Geological Survey of Canada with a LECO® AMA254 mercury analyser (10%  
160 precision, 5% relative standard deviation (RSD), Hall and Pelchat, 1997). Whole rock  
161 powders from Ruin Wash, as well as a duplicate sample set from Oak Springs  
162 Summit, were analysed at Amherst College (Massachusetts, USA) using a Teledyne  
163 Leeman Labs Hydra IIc mercury analyser (RSD <10%). Duplicate samples returned  
164 a correlation coefficient of 0.99, indicating a robust positive correlation between the  
165 results from the two laboratories.

166 Whole-rock powders were decarbonated using hydrochloric acid, and their  
167 carbonate content was calculated by mass loss following acid digestion. With the  
168 exception of Ruin Wash samples, TOC was measured from insoluble residues at the

169 University of Leeds using a LECO® SC-144DR Dual Range carbon and sulphur  
170 analyser. The carbon content of insoluble residues from Ruin Wash was measured  
171 with a Costech ECS 4010 elemental analyser at Amherst College in order to  
172 generate TOC measurements, with RSD <5%.

173 Inorganic carbon isotope values from the Carrara Formation at Emigrant Pass  
174 and Pioche Formation, Oak Springs Summit are reproduced from Faggetter et al.  
175 (2017), and new inorganic carbon isotope values from Ruin Wash are presented  
176 here. Whole-rock powders were analysed at the GeoZentrum Nordbayern, FAU  
177 Erlangen-Nuremberg, Germany, where carbon dioxide was prepared via reaction  
178 with phosphoric acid at 70°C using a Gasbench II preparation system; carbon  
179 isotope ratios were measured by a ThermoFisher Delta V plus mass spectrometer in  
180 continuous flow mode. Isotope ratios are reported relative to the V-PDB standard,  
181 with a reproducibility of  $\pm 0.06\text{‰}$  for  $\delta^{13}\text{C}$  and  $0.05\text{‰}$  for  $\delta^{18}\text{O}$ .

182 Inferred redox conditions (Figs. 2, 3, 4) are based upon pyrite framboid size  
183 distribution as reported in Faggetter et al. (2017); samples were assessed using a  
184 scanning electron microscope (FEI Quanta 650 FEG-ESEM) in backscatter mode  
185 following Bond and Wignall (2010).

186

## 187 **4. RESULTS**

### 188 **4.1. TOC concentrations**

189 Throughout all three sections TOC content is generally very low and exhibits  
190 correlation with facies (Fig. 5). The lowest TOC values (<0.15 wt% TOC) of all three  
191 sections are found in marl facies that coincide with extremely low/carbonate free  
192 intervals such as the C-Shale Member of the Pioche Formation (Fig. 5). In the  
193 Pioche Formation, higher TOC values are preserved in limestone of the Combined  
194 Metals Member, at Oak Springs Summit where levels reach 0.48 – 2.69 wt% within  
195 an oncoidal limestone of the Combined Metals Member (Fig. 5). The majority of the  
196 Carrara Formation is composed of marl with very low (<0.15 wt%) TOC content, the  
197 exception being horizons within the Echo Shale and Gold Ace members where  
198 values span the greatest range of all three sections (<0.15 – 5.17 wt% TOC) (Fig. 5).

199

### 200 **4.2. The ROECE and trilobite extinction**

201 The inorganic carbon isotope record from the Carrara Formation at Emigrant  
202 Pass and the Pioche Formation at Oak Springs Summit are discussed in Faggetter



203 et al. (2017) in which a C isotope excursion of  $\sim -3.8$  ‰ is interpreted to be the  
204 ROECE. Within the Pioche Formation at Oak Springs Summit, the most negative  
205 inorganic carbon isotope values coincide with the extinction horizon of the olenellid  
206 trilobites. ROECE is also expected to occur at Ruin Wash (Palmer 1998; Faggetter  
207 et al., 2017), but the extremely low carbonate content (below detection limits) in the  
208 shale of the C-Shale Member at this location does not allow measurement of a  
209 continuous inorganic carbon isotope curve. The ROECE is also observed within the  
210 Pyramid Shale Member of the Carrara Formation, but unlike the Pioche Formation,  
211 there is no trilobite fauna to delineate a clear extinction horizon at Emigrant Pass  
212 (Fig. 2). An inferred extinction interval is therefore proposed at Emigrant Pass within  
213 the Pyramid Shale Member (Fig. 2), based on the biostratigraphic boundary between  
214 the Olenellus Zone and the Eokochaspis nodosa Zone (Fig. 2; see Palmer and  
215 Halley, 1979; Faggetter et al, 2017).

216

### 217 **4.3. Hg concentrations, Hg/TOC ratios and extinction**

218 The Carrara and Pioche formations contain enrichments in Hg concentrations  
219 (ppb) and excursions in Hg/TOC (ppb/wt% TOC) ratios (Figs. 2, 3, 4). Hg/TOC ratios  
220 from samples with extremely low TOC of  $<0.01$  wt% C are not considered robust  
221 enough to record a primary Hg signal and are not plotted in figures but are included  
222 in Table 1.

223

#### 224 **4.3.1. Carrara Formation, Emigrant Pass**

225 The Carrara Formation at Emigrant Pass exhibits background Hg  
226 concentrations of  $<50$  ppb throughout the section (Fig. 2). High Hg and Hg/TOC  
227 values are seen in samples with both high and low TOC wt% values (Fig. 6). A  
228 number of prominent enrichments occur in the lowest 90 m, with the most enriched  
229 sample containing 270 ppb Hg found in the basal Eagle Mountain Shale Member.  
230 Smaller enrichments recorded by one or two data points each occur in the Echo  
231 Shale, Gold Ace and Pyramid Shale members. Within the ROECE interval Hg values  
232 are elevated at the beginning of the  $\delta^{13}\text{C}$  excursion before concentrations return to  
233 low levels for the remainder of the section.

234 The Hg/TOC enrichments in the Carrara Formation occur in two distinct  
235 pulses where Hg and Hg/TOC peaks correlate; an initial, multi-peak enrichment in  
236 the basal 30m of the Eagle Mountain Shale Member and another during early

237 ROECE at the base of the Pyramid Shale Member (Fig. 2). It is noteworthy that  
238 across the inferred extinction horizon, there are no abrupt Hg or Hg/TOC peaks and  
239 values remain stable in this interval.

240

#### 241 4.3.2. Pioche Formation, Oak Springs Summit

242 Background Hg concentrations in the Pioche Formation at Oak Springs  
243 Summit are <10 ppb (Fig. 3). Highest Hg values coincide with samples with low (<0.1  
244 wt%) TOC (Figs. 5 and 6). A single elevated value (46 ppb Hg) occurs in the  
245 Combined Metals Member, just below the base of the ROECE, and a smaller  
246 enrichment (32 ppb Hg) coincides with the most negative ROECE value and the  
247 olenellid extinction horizon at the base of the C-Shale Member. Finally, there is a  
248 small increase in Hg concentration (17 ppb) at ~35m in the section, which, like the  
249 other enrichments, correspond to three prominent Hg/TOC spikes at Oak Springs  
250 Summit (Fig. 3).

251

252

#### 253 4.3.3. Pioche Formation, Ruin Wash

254 High Hg and Hg/TOC values are recorded in samples with both high and low  
255 TOC wt% values (Fig. 6) at Ruin Wash. The Hg values are highest (up to 500 ppb) at  
256 the base of the measured section in the basal ~5m of the Combined Metals Member  
257 (Fig. 4). Peaks are around an order of magnitude higher than the maximum values  
258 seen at Oak Springs Summit. Above the level of elevated values at Ruin Wash, Hg  
259 concentrations are consistently <5 ppb, and there is no increase at the olenellid  
260 extinction horizon.

261

## 262 5. DISCUSSION

263 The three sections show an inconsistent correspondence between the  
264 olenellid trilobite extinction, ROECE and sedimentary Hg enrichments (Fig. 7). We  
265 review possible points for correlation between sections (e.g. ROECE interval and the  
266 trilobite extinction) and discuss processes which may account for the variable Hg  
267 signal.

268

### 269 5.1. ROECE, Hg and Hg/TOC correlation

270 To assess any correlation between the timing of Hg enrichment and ROECE,  
271 we delineate the base of the excursion based on the following two criteria. Firstly, the  
272 onset of ROECE should be present within the Pyramid Shale Member of the Carrara  
273 Formation and the upper Combined Metals Member of the Pioche Formation; this is  
274 based on an abrupt negative carbon isotope excursion within the *Olenellus* trilobite  
275 biozone, immediately preceding the olenellid extinction (Montañez et al., 2000).  
276 Secondly, given this stratigraphic constraint, we mark the onset of the excursion as  
277 the stratigraphic base of the negative inflexion interpreted within these members, i.e.  
278 ~60 m at Emigrant Pass and ~13 m at Oak Springs Summit. Between the Carrara  
279 and Pioche formations, our data show no clear relationship between the onset of  
280 ROECE and Hg or Hg/TOC enrichments. At Emigrant Pass an enrichment occurs  
281 around 10 m above the base of ROECE and at Oak Springs Summit an enrichment  
282 occurs ~1 m below the base of the excursion. It is clear from enrichments in the  
283 Eagle Mountain Shale Member of the Carrara Formation and in the Combined  
284 Metals Member of the Pioche Formation that the majority of Hg and Hg/TOC  
285 excursions occur before the ROECE interval.

286

## 287 **5.2. Extinction, Hg and Hg/TOC**

288 In the Pioche Formation at Oak Springs Summit, a small Hg (32 ppb,  
289 compared to a background of <5ppb for this section) and Hg/TOC excursion  
290 corresponds closely with the extinction horizon and the top of the *Olenellus* biozone  
291 (Fig. 7). However, the relationship between olenellid extinction and Hg or Hg/TOC  
292 enrichment is inconsistent among the other studied sections (Figs. 3 and 7). There is  
293 no enrichment coincident with the top of the *Olenellus* biozone at Emigrant Pass, nor  
294 at the extinction horizon at Ruin Wash (Fig. 7).

295 The high levels of Hg enrichment are generally recorded low in the study  
296 sections. In the Combined Metals Member at Ruin Wash, this is seen ~15 and ~10m  
297 below the olenellid extinction level at Ruin Wash (Fig. 4), and even lower below this  
298 level at Emigrant Pass (Fig. 2). Based on the lithostratigraphic correlation of Palmer  
299 (1998), it is unlikely that the levels of Hg enrichment can be correlated with each  
300 other (Fig. 7).

301 The inconsistency of the relationship between Hg, ROECE and extinction  
302 across all three sections challenges the conventional use of Hg as a tracer for global  
303 environmental Hg loading in this case. Previous Hg chemostratigraphic profiles

304 invoked to trace global LIP eruptions predict, and exhibit, synchronous Hg signals  
305 across regional and global sites (e.g. Percival et al., 2017). The lack of a  
306 reproducible Hg signal across our sections precludes a straightforward interpretation  
307 of the Hg chemostratigraphy. Our data show that only at Emigrant Pass does the  
308 base of ROECE coincide with Hg enrichment, but not the duration of the isotopic  
309 excursion. There is no Hg enrichment across the extinction interval.

310 The observed heterogeneity of enrichments in Hg and Hg/TOC values could  
311 be caused by several factors related to environmental and diagenetic processes. Our  
312 data demonstrate that volcanic loading and binding to organic matter cannot be the  
313 sole drivers of the Hg record in these strata. We therefore consider the possible roles  
314 that redox variations and binding to clays (and possibly sulphides) may have played  
315 in controlling Hg levels.

316

### 317 **5.3. Hg enrichments, redox variation and TOC**

318 Previous studies have exhibited limited correlation between redox conditions,  
319 organic matter deposition and Hg drawdown (e.g. Grasby et al., 2013; Percival et al.,  
320 2015). However, studies from the modern suggest the redox state of the sediment  
321 and water column can play an important role in mobilising or re-mobilising Hg  
322 species and enhancing or diminishing Hg exchange between the sediment and water  
323 column (Mason et al., 2006; Emili et al., 2011). Mercury and methyl mercury (MeHg)  
324 are scavenged in oxic conditions and sequestered in the sediments; however, the  
325 upward migration of the redox boundary causes Hg and MeHg to be released into  
326 the water column (Emili et al., 2011; Yin et al., 2017). Emili et al. (2011) modelled Hg  
327 cycling between the sediment and water column under anoxic conditions and  
328 showed a strong redox control on the mobility of Hg. They found that benthic Hg flux  
329 from the sediment to water column is highest during anoxic conditions and is also  
330 accentuated during sulfate reduction in euxinic conditions.

331 Assessment of redox states within the Carrara and Pioche formations found  
332 intermittent and locally variable periods of dysoxia during the olenellid extinction  
333 interval (Webster et al., 2008; Faggetter et al., 2017). Pyrite petrography shows  
334 dominantly oxygenated conditions (i.e. no framboids, scarce pyrite crystals) at  
335 Emigrant Pass (Fig. 2), variably oxygenated-dysoxic-euxinic conditions at Oak  
336 Springs Summit close to the extinction horizon (Fig. 3) and, euxinic to oxygenated  
337 conditions across the extinction horizon at Ruin Wash (Fig. 4) (Faggetter et al.

338 2017). These fluctuating redox conditions could have altered drawdown of Hg from  
339 the water column to the sediments (Horowitz et al., 2017) during times of heightened  
340 Hg loading. The euxinic pyrite framboid size data from two samples at the Ruin  
341 Wash trilobite extinction level do not correspond with Hg enrichment. Thus, the  
342 varying behaviour of Hg under different redox conditions could underlie the differing  
343 relationship between Hg and extinction in the Pioche Formation.

344 It is important to note that previously published studies reporting Hg  
345 sedimentary trends across multiple sections similarly reveal variable Hg and Hg/TOC  
346 ranges between localities (e.g. Grasby et al., 2016; Jones et al., 2017). Such  
347 discrepancies may be an inherent component of volcanically-derived deposition and  
348 fixation in marine sediments. For instance, a variable record of Hg enrichment during  
349 anoxia-related extinction is reported during the Toarcian (Early Jurassic) extinction  
350 (Percival et al., 2015). This event coincides with the organic-rich shales of the Jet  
351 Rock in northern England, but these sediments lack the Hg enrichment that  
352 otherwise might be expected given contemporaneous eruption of the Karoo-Ferrar  
353 flood basalt province. Percival et al. (2015) argue that efficient organo-Hg  
354 scavenging in organic-rich euxinic settings may have caused over-printing of the  
355 Hg/TOC anomaly by excess organic matter deposition. However, such a mechanism  
356 is unlikely in our reported Cambrian examples because TOC values in the C-Shale  
357 at Ruin Wash are low (<0.3 wt %; Fig. 5). The Carrara and Pioche formations are  
358 similarly characterised by low TOC and a range of Hg concentrations and Hg/TOC  
359 values; we interpret this signature as showing that even in organic-poor rocks it is  
360 possible to record elevated Hg concentrations during times of heightened  
361 environmental loading (Figs. 5 and 6), a conclusion drawn from other organic-lean  
362 records (e.g. Font et al., 2016; Percival et al., 2017). At Oak Springs Summit Hg and  
363 Hg/TOC excursions occur in samples containing low TOC (<0.15 wt% C), whilst at  
364 Emigrant Pass and Ruin Wash enrichments occur across a range of wt% TOC  
365 values (Fig. 6). The absence of a strong correlation between Hg and TOC (Fig. 5) at  
366 all three sections therefore indicates that Hg enrichment is not a function of variable  
367 TOC; we therefore posit that these anomalies are not solely a function of low TOC.

368

#### 369 **5.4. Hg and mineralogy**

370 Hg accumulation may also be controlled in part by binding to phases other  
371 than organic matter. The samples analysed here exhibit very low TOC (wt%) values

372 and commonly comprise marl facies. Because clay minerals can act as an efficient  
373 Hg binding medium in the absence of a larger organic matter reservoir, the  
374 sediment/rock composition at the time of Hg deposition may partially control Hg  
375 concentration (Zhong and Wang, 2008; Kongchum et al., 2011). High surface area  
376 reactivity for clay minerals make them effective regulators of Hg in aquatic sediments  
377 and up to an order of magnitude higher Hg concentrations have been found in  
378 secondary minerals such as clay versus primary minerals such as quartz and  
379 feldspar (Tessier et al., 1982). Higher proportions of primary silicate minerals  
380 therefore have the capability to “dilute” the amount of Hg binding during Hg loading  
381 (Tessier et al., 1982), and a variation in sediment composition across our study  
382 location could be a contributing factor to the observed inconsistent Hg records.

383

#### 384 **5.5. Hg as an indicator of volcanism**

385 The pre-ROECE levels of Hg enrichment observed at Emigrant Pass and  
386 Ruin Wash are enigmatic, and it is currently unclear if they record an unknown local  
387 volcanic source, an early eruptive pulse of the Kalkarindji LIP, or are instead a  
388 response to sedimentary/diagenetic redox variations. To resolve this uncertainty,  
389 further work is required to determine how widespread these perturbations are, both  
390 within Laurentia and globally.

391 Challenges to linking Hg excursions with LIP volcanism are not restricted to  
392 our Cambrian successions. Percival et al. (2018) present a comprehensive  
393 comparison of Hg records from the Mesozoic, focussing on the coincidence of LIP  
394 emplacement and ocean anoxic events, and report variable (both in Hg and  
395 Hg/TOC) concentrations across sections and lithologies and also contrasting  
396 evidence of Hg enrichment during periods of LIP volcanism (e.g. an absence of a  
397 broad global Hg excursion during Deccan volcanism contrasting with osmium-  
398 isotope records). The apparent inconsistency between these proxies is likely due to  
399 the various environments, lithologies and depositional processes.

400 Although the lack of consistent Hg records between the analysed sections  
401 prevents arguing definitively for a link between Kalkarindji and the extinction event at  
402 this stage, it remains likely that some of the recorded Hg excursions do record an  
403 expression of these LIP eruptions for the following reasons:

- 404 • The Kalkarindji LIP lacks a robust body of radiometric dates, Accordingly,  
405 additional dating efforts within the Kalkarindji province itself may also help

406 clarify its emplacement history, and whether initial volcanic pulses occurred  
407 considerably earlier than current geochronological constraints suggest.  
408 Current age dating does suggest that the onset of eruption may straddle, or  
409 possibly predates the Series 2 – Series 3 boundary (Marshall et al., 2018),  
410 thus raising the possibility that the Hg record from the Great Basin records  
411 other volcanic eruptions. If this proves to be the case, then Hg anomalies may  
412 be a useful marker for independently implicating specific volcanic events, but  
413 offers limited resolution when attempting to discriminate between multiple  
414 contemporaneous sources.

- 415 • Many LIPs, whilst erupted rapidly on a geological timescale, are iterative in  
416 their eruptive behaviour and being characterised by short periods of very  
417 intense activity (Chenet et al., 2008; Vye-Brown et al., 2013). For instance,  
418 high-resolution geochronological studies have revealed that eruptions of other  
419 large igneous provinces occur in a pulsed nature, particularly the Permo-  
420 Triassic Siberian Traps (Burgess et al., 2017), the end-Triassic Central  
421 Atlantic Magmatic Province (Davies et al., 2017) and the end-Cretaceous  
422 Deccan Traps (Schoene et al., 2015). Given the evidence for pulsed LIP  
423 emplacement throughout Earth history, it is plausible that multiple eruption  
424 episodes characterise the emplacement of the Kalkarindji LIP.
- 425 • The broad-scale architecture of the province, as evidenced by the several  
426 geochemically related sub-provinces (Glass and Phillips, 2006), may indicate  
427 more than one eruptive focus during its eruptive lifetime.
- 428 • Marshall et al. (2016) report that many Kalkarindji flows were effectively  
429 degassed: such near-complete degassing could have occurred either at the  
430 vent source (Guilbaud et al., 2007) or during propagation across the evolving  
431 lava fields. Importantly, degassing occurs either during fissure eruption and  
432 associated fire-fountaining similar to that observed from Laki eruptions. Such  
433 effusions can transport volcanogenic volatiles high into the troposphere, and  
434 possibly into the stratosphere, since fire fountains, augmented by heat  
435 released from nascent flows drive thermal uplift generating buoyant ash and  
436 gas plumes (Thordarson and Self, 1998; Glaze et al., 2017). Accordingly,  
437 volatiles may be lofted high into the atmosphere (Stothers et al., 1986;  
438 Woods, 1993), and this available for distribution certainly at local and regional

439 scales or, under favourable conditions, more globally. In this context, it is  
440 important to note that the Kalkarindji LIP was erupted near the equator (Fig. 1;  
441 Cocks and Torsvik, 2013; Lawver et al., 2015) where the tropopause would  
442 have been at a greater altitude thus mitigating against wider distribution, but  
443 that atmospheric circulation patterns an equatorial location would have  
444 otherwise aided in allowing aerosol distribution to both hemispheres

- 445 • In addition, Marshall et al., (2016) also argue for a fundamental change in  
446 eruptive style; the main succession being typified by thick, inflated pahoehoe  
447 flow and the overlying Blackfella Rockhole Member (BRM), which is instead  
448 characterised by huge rubble-topped flows. This change in eruptive style,  
449 together the occurrence of stromatolite reefs and aeolian(?) sand horizons  
450 preserved within these upper eruptive units indicates that later eruptive  
451 episodes occurred into a complex paleoenvironment affected by both  
452 terrestrial and shallow marine conditions.

453

454 To summarise, the degree and timing of Hg release is likely to have varied  
455 significantly during construction of the Kalkarindji LIP and, together with outlined  
456 factors controlling lofting and circulation of volatiles, may thus help explain the  
457 iterative and/or incomplete record of Hg here reported in the mid-Cambrian marine  
458 sediments.

459  
460

## 461 **6. CONCLUSION**

462 We report sedimentary Hg and Hg/TOC enrichments from both the Carrara  
463 and Pioche formations of the western Great Basin, USA. These successions are  
464 constrained within a biostratigraphic and chemostratigraphic framework to record the  
465 Cambrian Series 2 – Series 3 boundary, and the extinction of the Laurentian  
466 olenellid trilobites is observable within the Pioche Formation. The ROECE is present  
467 at two of the three studied sections (Emigrant Pass, Carrara Formation and Oak  
468 Springs Summit, Pioche Formation). In the Carrara Formation, the majority of Hg  
469 enrichments predate ROECE, with a single enrichment occurring just above the base  
470 of the excursion. At Oak Springs Summit ROECE is preceded by Hg enrichment.

471 Within the Pioche Formation at Oak Springs Summit the extinction horizon of  
472 the olenellid trilobites is marked by positive Hg and Hg/TOC excursions; however, a



473 similar excursion is not apparent at the equivalent horizon from Ruin Wash. The  
474 failure to locate Hg enrichment in the euxinic Ruin Wash section suggest that the  
475 redox conditions were unfavourable. Our data supports the hypothesis that Hg and  
476 Hg/TOC enrichments within the Carrara and Pioche formations are not solely derived  
477 from enhanced TOC preservation, but that inconsistent Hg trends may have resulted  
478 from variable environmental and diagenetic processes at the different sites. Given  
479 the strong control anoxia exerts on Hg flux, speciation and accumulation in modern  
480 settings, the role of redox states in deep time is clearly important when assessing the  
481 record of Hg in rocks.

482 The timing, volume and palaeo-position of the Kalkarindji LIP makes it a key  
483 candidate as the source of environmental Hg loading and subsequent enrichments in  
484 the Carrara and Pioche formations. Thus, the Kalkarindji potentially contributed to  
485 the Hg concentrations at the Cambrian Series 2 – Series 3 boundary, and that the  
486 occurrence of precursor levels of Hg enrichment may point to hitherto unrecognised  
487 phases of volcanism during construction of the Kalkarindji LIP, or else the possibility  
488 of other, as yet, unknown major volcanic episodes.

489

## 490 **ACKNOWLEDGMENTS**

491 LF was funded by a Natural Environment Research Council doctoral training  
492 grant at Leeds University. We thank Michael Joachimski and Yadong Sun for the use  
493 of their stable isotope lab at the Friedrich-Alexander Univesität, Erlangen-Nürnberg,  
494 and Maria Kopicki for laboratory assistance at Amherst College. We also thank A.N.  
495 Sial and one anonymous reviewer for their helpful feedback during the peer-review  
496 process.

497

## 498 **REFERENCES CITED**

- 499 Chenet, A. L., Courtillot, V., Fluteau, F., Gérard, M., Quidelleur, X., Khadri, S. F. R.,  
500 Thordarson, T. (2009). Determination of rapid Deccan eruptions across the  
501 Cretaceous-Tertiary boundary using paleomagnetic secular variation: 2.  
502 Constraints from analysis of eight new sections and synthesis for a 3500-m-  
503 thick composite section. *Journal of Geophysical Research: Solid*  
504 *Earth*, 114(B6).
- 505 Benoit, J.M., Gilmour, C.C., Mason, R.P., Heyes, R., (1999). Sulfide controls on  
506 mercury speciation and bioavailability to methylating bacteria in sediment pore

507 waters. *Environ. Sci. Technol.* 33, 951–957.

508 Benoit, J.M., Mason, R.P., Gilmour, C.C., Aiken, G.R., (2001). Constants for mercury  
509 binding by dissolved organic matter isolates from the Florida Everglades.  
510 *Geochim. Cosmochim. Acta* 65, 4445–4451.

511 Bergquist, B. A. (2017). Mercury, volcanism, and mass extinctions. *Proceedings of*  
512 *the National Academy of Sciences*, 114(33), 8675-8677.

513 Blackburn, T. J., Olsen, P. E., Bowring, S. A., McLean, N. M., Kent, D. V., Puffer, J.,  
514 Et-Touhami, M. (2013). Zircon U-Pb geochronology links the end-Triassic  
515 extinction with the Central Atlantic Magmatic Province. *Science*, 340(6135),  
516 941-945.

517 Bond, D. P., Wignall, P. B. (2010). Pyrite framboid study of marine Permian–Triassic  
518 boundary sections: a complex anoxic event and its relationship to  
519 contemporaneous mass extinction. *Geological Society of America*  
520 *Bulletin*, 122(7-8), 1265-1279.

521 Brasier, M. D., Corfield, R. M., Derry, L. A., Rozanov, A. Y., Zhuravlev, A. Y. (1994).  
522 Multiple  $\delta^{13}\text{C}$  excursions spanning the Cambrian explosion to the Botomian  
523 crisis in Siberia. *Geology*, 22(5), 455-458.

524 Burgess, S. D., Muirhead, J. D., Bowring, S. A. (2017). Initial pulse of Siberian Traps  
525 sills as the trigger of the end-Permian mass extinction. *Nature*  
526 *communications*, 8(1), 164.

527 Chang, C., Hu, W., Wang, X., Yu, H., Yang, A., Cao, J., Yao, S. (2017). Carbon  
528 isotope stratigraphy of the lower to middle Cambrian on the eastern Yangtze  
529 Platform, South China. *Palaeogeography, Palaeoclimatology,*  
530 *Palaeoecology*, 479, 90-101.

531 Cocks, L. R. M. and Torsvik, T.M. (2013). The dynamic evolution of the Palaeozoic  
532 geography of eastern Asia. *Earth-Science Reviews* 117: 40-79.

533 Courtillot, V. E., & Renne, P. R. (2003). On the ages of flood basalt events. *Comptes*  
534 *Rendus Geoscience*, 335(1), 113-140.

535 Dal Corso, J., Marzoli, A., Tateo, F., Jenkyns, H. C., Bertrand, H., Youbi, N., Cirilli, S.  
536 (2014). The dawn of CAMP volcanism and its bearing on the end-Triassic  
537 carbon cycle disruption. *Journal of the Geological Society*, 171(2), 153-164.

538 Davies, J. H. F. L., Marzoli, A., Bertrand, H., Youbi, N., Ernesto, M., Schaltegger, U.  
539 (2017). End-Triassic mass extinction started by intrusive CAMP  
540 activity. *Nature communications*, 8, 15596.

541 Emili, A., Koron, N., Covelli, S., Faganeli, J., Acquavita, A., Predonzani, S., De Vittor,  
542 C. (2011). Does anoxia affect mercury cycling at the sediment–water interface  
543 in the Gulf of Trieste (northern Adriatic Sea)? Incubation experiments using  
544 benthic flux chambers. *Applied geochemistry*, 26(2), 194-204.

545 Ernst, R.E. and Youbi, N. (2017). How large igneous provinces affect global climate,  
546 sometimes cause mass extinction, and represent natural markers in the  
547 geological record. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 478,  
548 30-52

549 Faggetter, L.E., Wignall, P.B., Pruss, S.B., Newton, R.J., Sun, Y.D. and Crowley, S.  
550 (2017). Trilobite extinctions, facies changes and the ROECE carbon isotope  
551 excursion at the Cambrian Series 2-3 boundary, Great Basin, western USA.  
552 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 478, 53-66

553 Fan, R., Deng, S., Zhang, X. (2011). Significant carbon isotope excursions in the  
554 Cambrian and their implications for global correlations. *Science China Earth  
555 Sciences*, 54(11), 1686-1695.

556 Fitzgerald, W.F., Lamborg, C.H., Hammerschmidt, C.R. (2007). Marine  
557 biogeochemical cycling of mercury. *Chemical Reviews*. 107, 641–662

558 Font, E., Adatte, T., Sial, A. N., de Lacerda, L. D., Keller, G., Punekar, J. (2016).  
559 Mercury anomaly, Deccan volcanism, and the end-Cretaceous mass  
560 extinction. *Geology*, 44(2), 171-174.

561 Glass, L. M., Phillips, D. (2006). The Kalkarindji continental flood basalt province: A  
562 new Cambrian large igneous province in Australia with possible links to faunal  
563 extinctions. *Geology*, 34(6), 461-464.

564 Glaze, L. S., Self, S., Schmidt, A., Hunter, S. J. (2017). Assessing eruption column  
565 height in ancient flood basalt eruptions. *Earth and Planetary Science  
566 Letters*, 457, 263-270.

567 Gong, Q., Wang, X., Zhao, L., Grasby, S. E., Chen, Z. Q., Zhang, L., Li, Z. (2017).  
568 Mercury spikes suggest volcanic driver of the Ordovician-Silurian mass  
569 extinction. *Scientific Reports*, 7.

570 Grasby, S.E., Sanei, H., Beauchamp, B., Chen, Z. (2013). Mercury deposition  
571 through the Permo–Triassic biotic crisis. *Chemical Geology*, 351, 209-216.

572 Grasby, S.E., Beauchamp, B., Bond, D.P.G., Wignall, P.B., Sanei, H. (2016).  
573 Mercury anomalies associated with three extinction events (Capitanian crisis,  
574 latest Permian extinction and the Smithian/Spathian extinction) in NW

575 Pangea. Geological Magazine 153, 285-297.  
576 doi:10.1017/S0016756815000436

577 Hall, G. E., Pelchat, P. (1997). Evaluation of a direct solid sampling atomic  
578 absorption spectrometer for the trace determination of mercury in geological  
579 samples. Analyst, 122(9), 921-924.

580 Horowitz, H. M., Jacob, D. J., Zhang, Y., Dibble, T. S., Slemr, F., Amos, H. M.,  
581 Sunderland, E. M. (2017). A new mechanism for atmospheric mercury redox  
582 chemistry: Implications for the global mercury budget. Atmospheric Chemistry  
583 and Physics, 17(10), 6353-6371.

584 Hough, M.L., Shields, G.A., Evins, L.Z., Strauss, H., Henderson, R.A., Mackenzie, S.  
585 (2006). A major Sulphur isotope event at c. 510 Ma: a possible anoxia-  
586 extinction-volcanism connection during the Early-Middle Cambrian transition?  
587 Terra Nova 18, 257-263. doi: 10.1111/j.1365-3121.2006.00687.x

588 Howley, R.A., Rees, M. N., Jiang, G. (2006). Significance of Middle Cambrian mixed  
589 carbonate-siliciclastic units for global correlation: southern Nevada,  
590 USA. Palaeoworld, 15(3), 360-366.

591 Jones, D. S., Martini, A. M., Fike, D. A. Kaiho, K. (2017). A volcanic trigger for the  
592 Late Ordovician mass extinction? Mercury data from south China and  
593 Laurentia. Geology, 45(7), 631-634.

594 Jourdan, F., Hodges, K., Sell, B., Schaltegger, U., Wingate, M.T. D., Evins, L.Z.,  
595 Blenkinsop, T. (2014). High-precision dating of the Kalkarindji large igneous  
596 province, Australia, and synchrony with the Early–Middle Cambrian (Stage 4–  
597 5) extinction. Geology, 42(6), 543-546.

598 Kongchum, M., Hudnall, W. H., Delaune, R. D. (2011). Relationship between  
599 sediment clay minerals and total mercury. Journal of Environmental Science  
600 and Health, Part A, 46(5), 534-539.

601 Kravchinsky, V.A. (2012). Paleozoic large igneous provinces of Northern Eurasia:  
602 Correlation with mass extinction events. Glob. Planet. Change 86–87, 31–36.  
603 doi:10.1016/j.gloplacha.2012.01.007

604 Landing, E. (2012). The great American carbonate bank in eastern Laurentia: Its  
605 births, deaths, and linkage to paleoceanic oxygenation (Early Cambrian – Late  
606 Ordovician), in J. R. Derby, R. D. Fritz, S. A. Longacre, W. A. Morgan, and C. A.  
607 Sternbach, eds., The great American carbonate bank: The geology and

608 economic resources of the Cambrian – Ordovician Sauk megasequence of  
609 Laurentia: AAPG Memoir 98, p. 451 – 492.

610 Lawver, L.A., Dalziel, I.W.D., Norton, I.O., Gahagan, L.M., Davis, J., (2015). The  
611 PLATES 2014 Atlas of Plate Reconstructions (550 Ma to Present Day), PLATES  
612 Progress Report No. 374-0215, University of Texas Technical Report No. 201,  
613 220p.

614 Li, Z. X., Zhang, L., Powell, C. M. (1996). Positions of the East Asian cratons in the  
615 Neoproterozoic supercontinent Rodinia. *Australian Journal of Earth  
616 Sciences*, 43(6), 593-604.

617 Li, Z. X., Bogdanova, S. V., Collins, A. S., Davidson, A., De Waele, B., Ernst, R. E.,  
618 Karlstrom, K. E. (2008). Assembly, configuration, and break-up history of  
619 Rodinia: a synthesis. *Precambrian Research*, 160(1), 179-210.

620 Marshall, P. E., Widdowson, M., Murphy, D. T. (2016). The Giant Lavas of  
621 Kalkarindji: rubbly pāhoehoe lava in an ancient continental flood basalt  
622 province. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 441, 22-37.

623 Marshall, P. E., Halton, A. M., Kelley, S. P., Widdowson, M., Sherlock, S. C. (2018).  
624 New <sup>40</sup>Ar/<sup>39</sup>Ar dating of the Antrim Plateau Volcanics, Australia: clarifying an  
625 age for the eruptive phase of the Kalkarindji continental flood basalt  
626 province. *Journal of the Geological Society*, 2018-035.

627 Mason, R. P., Fitzgerald, W. F., Hurley, J., Hanson, A. K., Donaghay, P. L., Sieburth,  
628 J. M. (1993). Mercury biogeochemical cycling in a stratified estuary. *Limnology  
629 and Oceanography*, 38(6), 1227-1241.

630 Mason, R. P., Kim, E. H., Cornwell, J., Heyes, D. (2006). An examination of the  
631 factors influencing the flux of mercury, methylmercury and other constituents  
632 from estuarine sediment. *Marine Chemistry*, 102(1-2), 96-110.

633 Merriam, C. W., Palmer, A. R. (1964). Cambrian rocks of the Pioche mining district,  
634 Nevada, with a section on Pioche shale faunules (No. 469). United States  
635 Geological Survey Professional Paper, 569, 59

636 Monperrus, M., Tessier, E., Amouroux, D., Leynaert, A., Huonnic, P., Donard, O. F.  
637 X. (2007). Mercury methylation, demethylation and reduction rates in coastal

638 and marine surface waters of the Mediterranean Sea. *Marine Chemistry*, 107(1),  
639 49-63.

640 Montañez, I. P., Osleger, D. A., Banner, J. L., Mack, L. E., Musgrove, M. (2000).  
641 Evolution of the Sr and C isotope composition of Cambrian oceans. *GSA*  
642 *Today*, 10(5), 1-7.

643 Moore, R. A., Lieberman, B. S. (2009). Preservation of early and Middle Cambrian  
644 soft-bodied arthropods from the Pioche Shale, Nevada, USA. *Palaeogeography,*  
645 *Palaeoclimatology, Palaeoecology*, 277(1), 57-62.

646 Munthe, J., Wängberg, I., & Shang, L. (2009). The origin and fate of mercury species  
647 in the natural environment. *Eur. Chlor. Sci. Doss*, 14.

648 Ogg, J. G., Ogg, G., Gradstein, F. M. (2016). *A Concise Geologic Time Scale: 2016.*  
649 Elsevier.

650 Palmer, A.R. (1998). Terminal early Cambrian extinction of the Olenellina:  
651 documentation from the Pioche Formation, Nevada. *Journal of Paleontology*,  
652 72 (04), 650-672.

653 Palmer, A. R., Halley, R. B. (1979). Physical stratigraphy and trilobite biostratigraphy  
654 of the Carrara Formation (Lower and Middle Cambrian) in the southern Great  
655 Basin (No. 1047). *United States Geological Survey Professional Paper*, 1047,  
656 113p

657 Payne, J.L., Lehrmann, D.J., Wei, J., Orchard, M.J., Schrag, D.P., Knoll, A.H. (2004).  
658 Large perturbations of the carbon cycle during recovery from the end-Permian  
659 extinction. *Science* 305 (5683), 506–509.

660 Percival, L.M.E., Witt, M.L.I., Mather, T.A., Hermoso, M., Jenkyns, H.C., Hesselbo,  
661 S.P., Al-Suwaidi, A.H., Storm, M.S., Xu, W., Ruhl, M. (2015). Globally  
662 enhanced mercury deposition during the end-Pliensbachian extinction and  
663 Toarcian OAE: A link to the Karoo-Ferrar Large Igneous Province. *Earth ad*  
664 *Planetary Science Letters* 428, 267-280.

665 Percival, L. M., Ruhl, M., Hesselbo, S. P., Jenkyns, H. C., Mather, T. A., Whiteside,  
666 J. H. (2017). Mercury evidence for pulsed volcanism during the end-Triassic  
667 mass extinction. *Proceedings of the National Academy of Sciences*, 114(30),  
668 7929-7934.

669 Percival, L. M., Jenkyns, H. C., Mather, T. A., Dickson, A. J., Batenburg, S. J., Ruhl,  
670 M., Woelders, L. (2018). Does large igneous province volcanism always perturb  
671 the mercury cycle? Comparing the records of Oceanic Anoxic Event 2 and the  
672 end-Cretaceous to other Mesozoic events. *American Journal of Science*, 318(8),  
673 799-860.

674 Prave, A.R. (1991). Depositional and sequence stratigraphic framework of the Lower  
675 Cambrian Zabriskie Quartzite: implications for regional correlations and the Early  
676 Cambrian paleogeography of the Death Valley region of California and Nevada.  
677 *Geological Society of America Bulletin*, 104(5), 505-515.

678 Pyle, D.M., Mather, T.A. (2003). The importance of volcanic emissions for the global  
679 atmospheric mercury cycle: *Atmospheric Environment*, v. 37, p. 5115–5124,  
680 doi:10.1016/j.atmosenv.2003.07.011.

681 Ren, Y., Zhong, D., Gao, C., Liang, T., Sun, H., Wu, D., Zheng, X. (2017). High-  
682 resolution carbon isotope records and correlations of the lower Cambrian  
683 Longwangmiao formation (stage 4, Toyonian) in Chongqing, South  
684 China. *Palaeogeography, Palaeoclimatology, Palaeoecology*.

685 Sanei, H., Grasby, S., Beauchamp, B. (2012). Latest Permian mercury anomalies.  
686 *Geology* 40, 63–66. <http://dx.doi.org/10.1130/G32596.1>.

687 Schuster, P. F., Krabbenhoft, D. P., Naftz, D. L., Cecil, L. D., Olson, M. L., Dewild, J.  
688 F., Abbott, M. L. (2002). Atmospheric mercury deposition during the last 270  
689 years: a glacial ice core record of natural and anthropogenic  
690 sources. *Environmental Science & Technology*, 36(11), 2303-2310.

691 Sial, A. N., Lacerda, L. D., Ferreira, V. P., Frei, R., Marquillas, R. A., Barbosa, J. A.,  
692 Pereira, N. S. (2013). Mercury as a proxy for volcanic activity during extreme  
693 environmental turnover: The Cretaceous–Paleogene  
694 transition. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 387, 153-  
695 164.

696 Sial, A. N., Chen, J., Lacerda, L. D., Peralta, S., Gaucher, C., Frei, R., Pereira, N. S.  
697 (2014). High-resolution Hg chemostratigraphy: A contribution to the distinction  
698 of chemical fingerprints of the Deccan volcanism and Cretaceous–Paleogene  
699 Boundary impact event. *Palaeogeography, Palaeoclimatology,*  
700 *Palaeoecology*, 414, 98-115.

701 Schmid, S. (2017). Chemostratigraphy and palaeo-environmental characterisation of

702 the Cambrian stratigraphy in the Amadeus Basin, Australia. *Chemical*  
703 *Geology*, 451, 169-182.

704 Schoene, B., Samperton, K. M., Eddy, M. P., Keller, G., Adatte, T., Bowring, S. A.,  
705 Gertsch, B. (2015). U-Pb geochronology of the Deccan Traps and relation to  
706 the end-Cretaceous mass extinction. *Science*, 347(6218), 182-184.

707 Stothers, R. B., Wolff, J. A., Self, S., Rampino, M. R. (1986). Basaltic fissure  
708 eruptions, plume heights, and atmospheric aerosols. *Geophysical Research*  
709 *Letters*, 13(8), 725-728.

710 Sundberg, F. A., McCollum, L. B. (2000). Ptychopariid trilobites of the lower-middle  
711 Cambrian boundary interval, Pioche Shale, southeastern Nevada. *Journal of*  
712 *Paleontology*, 74(4), 604-630.

713 Tessier, A. Campbell, P. G., Bisson, M. (1982). Particulate trace metal speciation in  
714 stream sediments and relationships with grain size: implications for  
715 geochemical exploration. *Journal of Geochemical Exploration*, 16(2), 77-104.

716 Thibodeau, A.M., Ritterbush, K., Yager, J.A., West, A.J., Ibarra, Y., Bottjer, D.J.,  
717 Berelson, W.M., Bergquist, B.A., Corsetti, F.A. (2016). Mercury anomalies and  
718 the timing of biotic recovery following the end-Triassic mass extinction. *Nature*  
719 *Communications* 7:11147, doi: 19.1038/ncomms11147

720 Veevers, J.J. (2001). Atlas of billion-year old earth history of Australia and  
721 neighbours in Gondwanaland. GEMOC Press, Sydney, Australia.

722 Vye-Brown, C., Self, S., & Barry, T. L. (2013). Architecture and emplacement of flood  
723 basalt flow fields: case studies from the Columbia River Basalt Group, NW  
724 USA. *Bulletin of Volcanology*, 75(3), 697.

725 Wang, X., Hu, W., Yao, S., Chen, Q., Xie, X. (2011). Carbon and strontium isotopes  
726 and global correlation of Cambrian Series 2-Series 3 carbonate rocks in the  
727 Keping area of the northwestern Tarim Basin, NW China. *Marine and*  
728 *Petroleum Geology*, 28(5), 992-1002.

729 Webster, M., Gaines, R. R., Hughes, N. C. (2008). Microstratigraphy, trilobite  
730 biostratigraphy, and depositional environment of the "Lower Cambrian" Ruin  
731 Wash Lagerstätte, Pioche Formation, Nevada. *Palaeogeography,*  
732 *Palaeoclimatology, Palaeoecology*, 264(1), 100-122.

733 Wignall, P. B. (2001). Large igneous provinces and mass extinctions. *Earth-science*  
734 *Reviews*, 53(1), 1-33.



- 735 Wignall, P.B. (2015). *The Worst of Times: How Life on Earth Survived 80 Million*  
736 *Years of Extinction*. Princeton University Press.
- 737 Woods, A. W. (1993). A model of the plumes above basaltic fissure  
738 eruptions. *Geophysical Research Letters*, 20(12), 1115-1118.
- 739 Yin, R., Xu, L., Lehmann, B., Lepak, R. F., Hurley, J. P., Mao, J., Hu, R. (2017).  
740 Anomalous mercury enrichment in Early Cambrian black shales of South  
741 China: Mercury isotopes indicate a seawater source. *Chemical Geology*. 467,  
742 159-167.
- 743 Zhang, W.H., Shi, X.Y., Jiang, G.Q., Tang, D.J., Wang, X.Q. (2015). Mass-  
744 occurrence of oncoids at the Cambrian Series 2-Series 3 transition:  
745 Implications for microbial resurgence following an Early Cambrian extinction.  
746 *Gondwana Research* 28, 432-450.
- 747 Zhong, H., Wang, W. X. (2008). Effects of sediment composition on inorganic  
748 mercury partitioning, speciation and bioavailability in oxic surficial  
749 sediments. *Environmental pollution*, 151(1), 222-230.
- 750 Zhu, M. Y., Zhang, J. M., Li, G. X., Yang, A. H. (2004). Evolution of C isotopes in the  
751 Cambrian of China: implications for Cambrian subdivision and trilobite mass  
752 extinctions. *Geobios*, 37(2), 287-301.
- 753 Zhu, M. Y., Babcock, L. E., & Peng, S. C. (2006). Advances in Cambrian stratigraphy  
754 and paleontology: integrating correlation techniques, paleobiology, taphonomy  
755 and paleoenvironmental reconstruction. *Palaeoworld*, 15(3), 217-222.
- 756 Zhu, D. C., Zhao, Z. D., Niu, Y., Dilek, Y., Wang, Q., Ji, W. H., Mo, X. X. (2012).  
757 Cambrian bimodal volcanism in the Lhasa Terrane, southern Tibet: record of  
758 an early Paleozoic Andean-type magmatic arc in the Australian proto-Tethyan  
759 margin. *Chemical Geology*, 328, 290-308.

760

761

762

### 763 Figure captions

- 764 1. Cambrian global palaeogeography at 510 million years ago showing  
765 approximate palaeoposition of study sections, adapted from Lawver et al.  
766 (2014). Architecture of Kalkarindji LIP taken from Marshall (2016) and aligned  
767 to the palaeogeographic position of Australia at 510 Ma (green outline) as  
768 depicted in Lawver et al. (2014).

- 769 2. Geochemical data from Emigrant Pass, Carrara Formation: TOC wt % C,  
770  $\delta^{13}\text{C}_{\text{org}}$  (permil), Hg (ppb) and Hg/TOC (ppb/ wt % C). The position of the  
771 Olenellus - Eokochaspsis nodosa biozone boundary is from Palmer and  
772 Halley (1979) inferred redox conditions are based on framboid size  
773 distribution data (Faggetter et al. 2017).
- 774 3. Geochemical data from Oak Springs Summit, Pioche Formation: TOC wt % C,  
775  $\delta^{13}\text{C}_{\text{org}}$  (permil), Hg (ppb) and Hg/TOC (ppb/ wt % C). The position of the  
776 Olenellus - Eokochaspsis nodosa biozone boundary is from Palmer (1998),  
777 inferred redox conditions are based on framboid size distribution data from  
778 Faggetter et al. (2017). See Fig. 2 for key.
- 779 4. Geochemical data from Ruin Wash, Pioche Formation: TOC wt % C,  $\delta^{13}\text{C}_{\text{org}}$   
780 (permil), Hg (ppb) and Hg/TOC (ppb/ wt % C). The position of the Olenellus -  
781 Eokochaspsis nodosa biozone boundary is from Palmer (1998), inferred redox  
782 conditions are based on framboid size distribution data from Faggetter et al.  
783 (2017). See Fig. 2 for key.
- 784 5. Stratigraphic columns showing TOC wt % C through the Carrara and Pioche  
785 formations. See Fig. 2 for key.
- 786 6. Cross plots of TOC wt % C vs. Hg (ppb) for the Carrara and Pioche  
787 formations.
- 788 7. Summary of Hg (ppb) and Hg/TOC (ppb/ wt % C) from the Carrara and  
789 Pioche formations. The position of the Olenellus - Eokochaspsis nodosa  
790 biozone boundary is from Palmer (1998), inferred redox conditions are based  
791 on framboid size distribution data from Faggetter et al. (2017). The yellow  
792 highlight correlates the Gold Ace Member of the Carrara Formation with the  
793 Combined Metals Member of the Pioche, after Palmer (1998). See Fig. 2 for  
794 key.

795

796 Table 1. Table containing stratigraphic and geochemical data. Rows coloured in grey  
797 indicate samples excluded due to low TOC (< 0.01 wt% TOC). Rows coloured in  
798 orange indicate the extinction horizon at Oak Springs Summit and Ruin Wash.

799

800

801

510 Ma

60° N



30° N

0°


30° S

60° S






-  Kalkarindji LIP
-  Approximate location of western Great Basin study sections



# Emigrant Pass, CA.





## KEY



Oak Springs Summit, NV.



Ruin Wash, NV.



## Emigrant Pass, CA.






## TOC (wt% C)

## Oak Springs Summit, NV.


## TOC (wt% C)

## Ruin Wash, NV.


## TOC (wt% C)



Ruin Wash, NV.



Oak Springs Summit, NV.



Emigrant Pass, CA.

