

6-1992

## The Holocene Carbonate Eolianites of North Point and Some Nearby Marine Environments, San Salvador Island, Bahamas: A Field Trip Guide

Brian White  
*Smith College*

H. Allen Curran  
*Smith College*, [acurran@smith.edu](mailto:acurran@smith.edu)

Follow this and additional works at: [https://scholarworks.smith.edu/geo\\_facpubs](https://scholarworks.smith.edu/geo_facpubs)

 Part of the [Geology Commons](#)

---

### Recommended Citation

White, Brian and Curran, H. Allen, "The Holocene Carbonate Eolianites of North Point and Some Nearby Marine Environments, San Salvador Island, Bahamas: A Field Trip Guide" (1992). Geosciences: Faculty Publications, Smith College, Northampton, MA.  
[https://scholarworks.smith.edu/geo\\_facpubs/134](https://scholarworks.smith.edu/geo_facpubs/134)

This Conference Proceeding has been accepted for inclusion in Geosciences: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact [scholarworks@smith.edu](mailto:scholarworks@smith.edu)

# Field Trip Guidebook

The  
**6<sup>th</sup> Symposium**  
on the  
**Geology**  
of the  
**Bahamas**



**June 11 - 15, 1992**  
at the  
Bahamian Field Station  
San Salvador, Bahamas



# THE HOLOCENE CARBONATE EOLIANITES OF NORTH POINT AND SOME NEARBY MARINE ENVIRONMENTS, SAN SALVADOR ISLAND, BAHAMAS: A FIELD GUIDE

Brian White and H. Allen Curran  
Department of Geology  
Smith College  
Northampton, Massachusetts 01063, USA



REPRINTED FROM:

*FIELD TRIP GUIDEBOOK: The 6<sup>th</sup> Symposium on the Geology of the Bahamas*  
June 11–15, 1992  
Bahamian Field Station, San Salvador, Bahamas, p. 33-39.

# THE HOLOCENE CARBONATE EOLIANITES OF NORTH POINT AND SOME NEARBY MARINE ENVIRONMENTS, SAN SALVADOR ISLAND, BAHAMAS

Brian White and H. Allen Curran  
Department of Geology  
Smith College  
Northampton, MA 01063

## INTRODUCTION

The goals of this part of the field trip are to study the types of strata, sedimentary structures, trace fossils, and dune morphologies of the Holocene eolianites excellently exposed along the coasts of Rice Bay and North Point. Here the mechanisms of how eolian deposition of carbonate grains occurs and how dunes are initiated and evolve are well illustrated in the rocks. Toward the end of this excursion, a short snorkel dive can be conducted in the shallow, sheltered waters between North Point and Cut Cay to study grass beds, sandy substrates, and hard substrates and to compare their associated sediments, plants, and animals. The area to be visited and the locations of field stops are shown in Fig. 1.

## Previous Work

Adams (1980), in his pioneering study of the geology of San Salvador, briefly described the eolianites of North Point. He distinguished lobate dunes that have large-scale cross beds on their leeward faces and flanks and smaller, more varied, cross beds on their windward slopes. In their overview study of the petro-graphy of the eolianites of San Salvador, Hutto and Carew (1984) found that most are dominantly oolitic. However, this was not true for their North Point samples, which have a higher proportion of skeletal grains and pellets. The North Point eolianites were assigned to the North Point Member of the Rice Bay Formation by Carew and Mylroie (1985). Samples of North Point Member rocks from North Point gave radiocar-

bon dates of  $5,345 \pm 125$  and  $5,360 \pm 110$  years B.P. (Carew and Mylroie, 1987).

In her more detailed study of the eolianites of the Rice Bay area, Lawlor (unpublished data, 1985) found that these rocks are pelsparites with lesser proportions of ooids and skeletal grains. The rocks are dominantly aragonite with incomplete cementation by vadose low Mg calcite. Inverse graded bedding is a common feature of the rocks on a microscopic scale, with somewhat more complete cementation in the finer layers (White and Curran, 1988). More recently, White and White (1991) have shown that these eolianites do bear evidence of diagenesis from the ongoing episode of Holocene marine transgression.




Fig. 1. The North Point area with locations for described stops.

## The Eolianites of the Rice Bay to North Point Area

The Holocene dunes of this area are composed of carbonate sand which was blown landward from marine beaches by onshore winds. Initially, small dunes developed landward of the beach, a few around cores formed of the eroded remnants of earlier dunes. In some cases two small adjacent dunes were enveloped by later dune strata to form a compound dune. Elsewhere, dunes grew higher and migrated inland as individual lobes, with slip faces sloping not only downwind, but to the right and left on the flanks of each lobe. Eventually the lower parts of the flanks of adjacent dunes overlapped to cover the interdune areas and thereby form a long, hummocky dune ridge. This dune morphology is clearly reflected in the undulating topography of the west side of North Point as seen from the campus of the Bahamian Field Station.

Large-scale, steeply-dipping cross beds occur on many lee slopes and flanks of the lobate dunes. Windward dune slopes reveal a greater variety in scale and type of cross bedding, including examples of tabular-planar, wedge-planar, and trough sets. The latter are more numerous in the lower parts of the dunes. Wind ripples are visible on some bedding surfaces, but they are scarce. In some of the cross bedded exposures, it is possible to distinguish the different strata produced by climbing wind ripples, grainfall, and sandflow as described from modern coastal dunes by Hunter (1977).

Although fossil burrows previously have been thought to be rare in eolianites (McKee and Ward, 1983), there are several large and well preserved burrows and numerous smaller burrows here, all thought to have been formed by invertebrates in dunal sands that now are the eolianite exposures along Rice Bay (White and Curran, 1988). Rhizomorphs, trace fossils produced by plant roots, are common in these eolianites. An overview of the significance of this trace fossil assemblage was given by Curran and White (1991).

## DESCRIPTIONS OF THE FIELD STOPS

### Introduction

The beach in front of the former U.S. Coast Guard Station makes a convenient starting point for this trip. From here the view extends northeast over the sandy beach and across the waters of Rice Bay to Man Head Cay. To the northwest (left) some of the North Point eolianite exposures can be seen and the first of these are reached by walking about 90 m along the beach in a northwesterly direction.

Continuous exposures of Holocene dunes in sea cliffs and on narrow, rocky shore platforms along this reach of coast reveal numerous features of the eolianites. The stops described below were selected to demonstrate particularly good, and readily accessible, examples of trace fossils, sedimentary structures, and stratum types found in the eolianites, and of dune morphology. After walking about 1 km along the peninsula, the end of North Point is reached, where final observations of the dunes can be made. The small beach on the west side of the point is a good place to begin a snorkel dive to view the nearby hard substrates, *Thalassia* grass beds, and sandy substrates and to examine their associated sediments, floras, and faunas.

### Stop 1. Cluster Burrow Trace Fossil Type Locality

To reach this stop, walk northwest along Coast Guard Beach to the first rock exposures and then continue over outcrops and a small sandy bay for 60 m. Here sea cliffs 3 to 4 m high are cut into an 85 m wide fossil dune. Along much of this dune's width cross bed dip directions are rather variable, but generally towards the southwest, and dip angles range from almost flat-lying to 15°. On the northwest flank of the dune, dips are northwesterly and steepen to 30°, with some sandflow cross strata evident. At the southeast end of the dune, steepening cross strata dip to the south at angles of up to 20°. Most of the strata are in small wedge-planar sets, suggesting variable wind directions,

and climbing wind-ripple strata are evident in some places.

A prominent trace fossil consisting of a cluster of vertically oriented burrows is exposed in a small cliff face here, and in a large counterpart block that has fallen away from the cliff. The trace fossil cuts vertically across 1.4 m of small-scale, wedge-planar and trough cross strata sets, which are obscured in places by bioturbation. Fine, millimeter laminations are evident on much of the cliff face and weathered surfaces reveal some very thin laminations of slightly coarser calcarenite. In the rocks immediately overlying this trace fossil, rhizomorphs are prominently displayed on some bedding planes, where they weather out in relief.

In detail, this trace fossil consists of multiple, straight to gently curved, unlined shafts. Shaft diameters are 1 to 2 centimeters (average 1.2-1.4 cm), and shaft lengths can be as long as at least 1.4 meters. This is a minimum length for the shafts of this specimen because a break in the cliff face in which it is exposed terminates the lower part of the specimen. Shafts occasionally branch in the upward direction and some definite cross-overs also occur. Shaft diameters tend to contract somewhat toward their upward ends.

The apparently obligatory clustered nature of the shafts and their large number suggest that the structure records the brooding/hatching activity of an invertebrate organism, very possibly a species of burrowing (digger) wasp of the Family Sphecidae, with the shafts having been formed by the juvenile wasps burrowing up to the surface. This hypothesis was discussed in some detail by Curran and White (1987).

About 50 m northwest of the trace fossil just described, a similar one is revealed in horizontal cross-section. Here the circular nature of the cluster and the large number of individual burrows (about 800 shafts) that it contains are clearly revealed. Here, too, this trace fossil is within a sequence of small-scale, wedge-planar and trough cross strata sets, with scattered rhizomorphs and climbing-ripple laminations. Of additional interest is a small bedding surface at least 1 m below the cluster trace fossil and

within the present day intertidal zone. On this bedding plane there are ripple marks with their crests oriented perpendicular to the strike of the bedding plane, a feature believed to indicate an eolian origin (McKee and Ward, 1983). The ripples have a very low amplitude and ripple indices of 25 to 30, further evidence that they are wind-formed ripples (McKee, 1979; Tanner, 1967). Two interesting conclusions may be drawn from these observations. As wind-deposited strata are located in the present intertidal zone, this clearly means that sea level was lower at the time of formation of these beds than at present. The presence of wind-formed ripples beneath the trace fossil confirms that the burrowing took place in an eolian dune and not in a beach or nearshore environment.

## **Stop 2. Fossil Proto-dunes on Rocky Shore Platform**

This next locality is reached by dropping down the northwest flank of the dune at Stop 1 to a broad, rocky shore platform, some 80 m long and up to 20 m wide, backed on its landward side by low cliffs and extending seaward into the intertidal zone. Because of the extensive horizontal and vertical exposures, this is an excellent place to study sedimentary structures and the early stage of dune development.

In the landward cliffs several small dune cores are exposed. Some of these are better lithified and contain more abundant rhizomorphs than overlying strata and appear to be the eroded remnants of earlier dunes. Large-scale trough cross beds occur immediately adjacent to some of the dune cores, and these may have formed by windscouring around the dune remnants. Subsequent deposition of wind-blown sand buried the dune cores and the growing dune extended laterally and vertically to encompass them into a form of compound dune.

Other sedimentary structures well-displayed here are convex-upward cross strata sets, trough cross beds, and wedge-planar sets. Some of the latter have cross strata with acute angular relationships to the underlying set, whereas others show a tangential relationship. Several



bedding surfaces have ripple marks with ripple indices between 23 and 32, clearly wind ripples (McKee, 1979; Tanner, 1967). Again, some of these are within the present intertidal zone, confirming the conclusions drawn at Stop 1 about lower sea level at the time of eolianite deposition.

Beyond the rocky terrace of Stop 2, follow the coast as it takes a short jog to the west, then cross a 30 m wide bay with a sand and rock floor, to reach another promontory. Scramble down the northerly side of this headland and traverse the narrowing beach for about 40 m, until a small rock arch is reached. Progress along here will be blocked at some point, exactly where depending on wind conditions and the state of the tide. In any case, from the vicinity of the small arch, climb upward obliquely across the cliffs to a small gully, which constitutes the next stop.

### Stop 3. Eolianite Stratification Types

In this locality are exposed, in close proximity, three types of wind-deposited strata that were produced by mechanisms described by Hunter (1977, 1981) as grainfall, sandflow down lee slopes, and climbing wind-ripples.

The strata produced by climbing wind-ripples are millimeter laminations with even thickness and sharp contacts resulting from inverse size grading, although the latter detail is not always readily visible in the field. If net sedimentation is to occur by the migration of wind-ripples, then each successive ripple must climb relative to the stratum deposited by the previous one. Grain size segregation in wind-ripples concentrates relatively coarse sediment on the crests and relatively fine sediment in the troughs. As the crests and troughs migrate, they deposit a layer of relatively coarse grains overlying a layer of relatively fine ones; hence the upward size grading within each stratum produced during deposition by a migrating wind-ripple. Wind-ripples may climb up, down, or along both lee and stoss sides of dunes. Thus the dip angle and direction of the resulting strata are more a function of the geometry of the surface

over which they have migrated than the direction of the driving wind. The passage of many wind-ripples can lead to the accumulation of sets of ripple-formed strata. On lee slopes these strata may be preserved under grainfall sediments or sandflows, providing the latter are not erosive.

Grainfall occurs when moving air currents carry saltating and suspended sediment into a sheltered area, for example the zone of separation to the lee of a dune crest. The sediment settles like falling snow and accumulates on the lee slope of the dune, where it may be joined by grains that crept over the dune crest in response to collisions with saltating grains. Grainfall strata tend to be thin and indistinct, and, because they commonly form on lee slopes, they often have a high initial dip (Hunter, 1981). On small dunes, though, grainfall could occur as far forward as the toe of the dune, and the strata would lie at low angles. In wind tunnel experiments conducted by Fryberger and Schenk (1981), grainfall strata deposited on lee slopes consistently wedged thinner downslope, and this may be anticipated on natural dune lee slopes as well.

Sandflow strata form by resedimentation of sands that accumulate on the upper part of lee slopes, often by grainfall, until the slope oversteepens and becomes unstable. If the sands are dry, they will flow non-cohesively, but, if crusted or partially lithified in some way, they may founder as blocks subject to all kinds of jumbling and deformation. Sandflow strata are typically thicker than other wind-deposited strata, commonly exceeding 1 cm. They have sharp contacts, lie close to the angle of repose, and tend to pinch out towards the base of a foreset (Hunter, 1981). They have a distinctive lenticular shape when seen in strike cross-section or in horizontal exposures.

These types of wind-formed strata were recognized by studying modern coastal dunes (Hunter, 1977). Several attempts have been made to use these strata to identify and more closely characterize ancient siliciclastic rocks believed to be of eolian origin. A sampling includes: Pleistocene of Oregon (Hunter, 1980); Permian of Arran, Scotland (Clemmensen and

Abrahamsen, 1983); and various Paleozoic and Mesozoic formations of the western United States (Fryberger and Schenk, 1981; Hunter, 1981). Prior to this study, similar analyses do not seem to have been reported for carbonate rocks, and these various wind-deposited strata are not mentioned by McKee and Ward (1983) in their review of carbonate eolian environments.

Small, irregularly meandering burrows 3 to 4 mm in diameter and reaching greater than 20 cm in length occur within and upon grainfall and sandflow strata in the North Point Member eolianites (White and Curran, 1988). Examples of these trace fossils are common in the vicinity of Stop 3. We suggest that these burrows also probably were formed by insects or insect larvae, but a specific modern trace-maker analogue for this burrow type has not yet been identified. Recently we have found similar small, irregular burrows and cluster burrows in Holocene eolianites on Lee Stocking Island of the Exuma Cays.

To continue this field trip, stay at the top of the sea cliffs and walk around the small bays and headlands for about 200 m. Hereabouts a more prominent trail joins from the south, and the wreck of a misplaced tanker scars the coast to the northeast. Follow the winding trail northward along the spine of the narrowing peninsula. Along the way one will pass many exposures of eolianites, and one can enjoy fine views to the west (left) over Grahams Harbour and to the east over Rice Bay and Man Head Cay. After walking about 400 m beyond the wreck, the edge of a cliff is reached, overlooking a tidal inlet and an island to the north. This is the next field trip locality.

#### Stop 4. Cut Cay Overlook

Here, at the north end of North Point, the cliffs are formed by the north flank of a well-developed lobate dune, and the cross bedding dips north and steeply down into the sea. A 40 m wide inlet, The Cut, separates North Point from the nearby island of Cut Cay. The cliffs of the south end of Cut Cay are part of the south

flank of another dune, and the cross bedding dips down into the sea on that side too, but in a southerly direction. Evidently, the sea has driven through along a low interdune area and separated Cut Cay from the rest of the peninsula. According to legend, The Cut did not exist at the time of Columbus' visit in 1492.

From this location a good view of the seafloor to the west out into Grahams Harbour and to the northwest towards Cut Cay can be obtained. This perch provides an excellent overview of the three different substrates which are easily explored by snorkeling in this calm (usually) water. The dark green grassbeds are dominated by *Thalassia*, the pale green areas are sandy bottoms, and the tan areas are hard substrates. Calcareous green algae, including *Halimeda*, *Penicillus*, *Udotea*, and *Acetabularia*, grow in the grassy and sandy areas, their abundance and distribution varying from time to time, perhaps seasonally. A considerable variety of invertebrate animals lives among the various plants of these different environments and awaits careful and sharp-eyed explorers.

Following this preview, climb down the west side of North Point by taking the only obvious (and safe) route to the small beach. This is the location of the next field trip stop and the starting point for the snorkel dive.

#### Stop 5. Dune Morphology Revealed in Sea Cliffs

A number of well-developed lobate dunes are clearly exposed in the cliffs on the west side of North Point. Here the dunes have reached a more mature stage of development than some seen along Rice Bay. The opposing flanks of each dune dip steeply and in opposite directions. Along this part of the coast the relationships between adjacent dunes are revealed. In some cases, one dune flank overlaps the flank of the nearest dune, suggesting that the former was mobile and the latter stabilized, at least temporarily. In other situations, adjacent dune flanks interfinger and both dunes appear to have been mobile. This entire coastline is made up of a row of these coalesced lobate dunes that



is clearly visible from the vicinity of the Bahamian Field Station, especially when illuminated by the setting sun. Dune lobes that coalesce to form such a transverse dune ridge have been described from Pleistocene carbonate rocks of other parts of the Bahamas by Ball (1967) and from Bermuda by MacKenzie (1964a,b).

The fact that wind-deposited cross beds dip down into the sea here at North Point adds further evidence that these eolian dunes formed before sea level rose to its present position. Additionally, such evidence shows that the wind-blown sands were sufficiently lithified by the time sea level rose to resist simple reworking of the sand.

### Stop 6. Snorkeling Beach

The small beach here is an excellent place to begin a snorkel dive to explore various substrates and associated flora, fauna, and sediments between North Point and Cut Cay. From a short distance offshore an excellent view of several of the coalescing dune lobes may be obtained.

### ACKNOWLEDGMENTS

We thank the staff of the Bahamian Field Station for full logistical support of our field work on San Salvador. Valuable contributions to the North Point research project were made by Smith College geology majors: Liz Dole, Kate Japy, Karen Kurkky, Jean Lawlor, Kim Pirie, and Kathy White. Acknowledgment is made to the donors of the The Petroleum Research Fund, administered by the American Chemical Society, for partial support of our work through separate grants to Curran and White. Generous financial support also was provided by the Merck, Keck, Pew, and Shell foundations, and we extend our thanks to each of them.

### REFERENCES

- Adams, R.W., 1980, General guide to the geological features of San Salvador, *in* Field Guide to the Geology of San Salvador, edited by D.T. Gerace, CCFL Bahamian Field Station, San Salvador, Bahamas, pp. 1-66.
- Ball, M.M., 1967, Carbonate sand bodies of Florida and the Bahamas: *Journal of Sedimentary Petrology*, 37, 556-591.
- Carew, J.L. and J.E. Mylroie, 1985, The Pleistocene and Holocene stratigraphy of San Salvador Island, Bahamas, with reference to marine and terrestrial lithofacies in French Bay, *in* Pleistocene and Holocene Carbonate Environments on San Salvador Island, Bahamas - Guidebook for Geological Society of America, Orlando Annual Meeting field trip #2, edited by H.A. Curran, CCFL Bahamian Field Station, San Salvador, Bahamas, pp. 11-61.
- Carew, J.L. and J.E. Mylroie, 1987, A refined geochronology for San Salvador Island, Bahamas *in* Proceedings of the Third Symposium on the Geology of the Bahamas, edited by H.A. Curran, CCFL Bahamian Field Station, San Salvador, Bahamas, pp. 35-44.
- Curran, H.A. and B. White, 1987, Trace fossils in carbonate upper beach rocks and eolianites: Recognition of the backshore to dune transition, *in* Proceedings of the Third Symposium on the Geology of the Bahamas, edited by H.A. Curran, CCFL Bahamian Field Station, San Salvador, Bahamas, pp. 243-254.
- Curran, H.A. and B. White, 1991, Trace fossils of shallow subtidal to dunal ichnofacies in Bahamian Quaternary carbonates: *Palaos*, 6, 498-510.
- Clemmensen, L.B. and K. Abrahamsen, 1983, Aeolian stratification and facies association in desert sediments, Arran basin (Permian), Scotland: *Sedimentology*, 30, 311-339.

- Fryberger, S.G. and C. Schenk, 1981, Wind sedimentation tunnel experiments on the origins of aeolian strata: *Sedimentology*, 28, 805-821.
- Hunter, R.E., 1977, Basic types of stratification in small aeolian dunes: *Sedimentology*, 24, 361-387.
- Hunter, R.E., 1980, Depositional environments of some Pleistocene coastal terrace deposits, southwestern Oregon - Case history of a progradational beach and dune sequence: *Sedimentary Geology*, 27, 241-262.
- Hunter, R.E., 1981, Stratification styles in eolian sandstones: Some Pennsylvanian to Jurassic examples from the Western Interior U.S.A., *in* Recent and Ancient Nonmarine Depositional Environments: Models for Exploration, edited by F.A. Ethridge and R.M. Flores, Society of Economic Paleontologists and Mineralogists Special Publication 31, pp. 315-329.
- Hutto, T. and J.L. Carew, 1984, Petrology of eolian calcarenites, San Salvador, Bahamas, *in* Proceedings of the Second Symposium on the Geology of the Bahamas, edited by J.W. Teeter, CCFL Bahamian Field Station, San Salvador, Bahamas, pp. 197-207.
- MacKenzie, F.T., 1964a, Geometry of Bermuda calcareous dune cross-bedding: *Science*, 144, 1449-1450.
- MacKenzie, F.T., 1964b, Bermuda Pleistocene eolianites and paleowinds: *Sedimentology*, 3, 52-64.
- McKee, E.D., 1979, Ancient sandstones considered to be eolian, *in* A Study of Global Sand Seas, edited by E.D. McKee, United States Geological Survey Professional Paper 1052, pp. 187-238.
- McKee, E.D. and W.C. Ward, 1983, Eolian environments, *in* Carbonate Depositional Environments, edited by P.A. Scholle, D.G. Bebout, and C.H. Moore, American Association of Petroleum Geologists Memoir 33, pp. 131-170.
- Tanner, W.F., 1967, Ripple mark indices and their uses: *Sedimentology*, 9, 89-104.
- White, B. and H.A. Curran, 1988, Mesoscale physical sedimentary structures and trace fossils in Holocene carbonate eolianites from San Salvador Island, Bahamas: *Sedimentary Geology*, 55, 163-184.
- White, K.S. and B. White, 1991, The effects of Holocene sea-level rise on the diagenesis of Quarternary carbonate eolianites, San Salvador Island, Bahamas, *in* Proceedings of the Fifth Symposium on the Geology of the Bahamas, edited by R.J. Bain, Bahamian Field Station, San Salvador, Bahamas, pp. 235-247.