

---

1-1-2019

## Effects of Thermal Stress on Amount, Composition, and Antibacterial Properties of Coral Mucus

Rachel M. Wright

*Harvard Medical School, [rwright@smith.edu](mailto:rwright@smith.edu)*

Marie E. Strader

*The University of Texas at Austin*

Heather M. Genuise

*The University of Texas at Austin*

Mikhail Matz

*The University of Texas at Austin*

Follow this and additional works at: [https://scholarworks.smith.edu/bio\\_facpubs](https://scholarworks.smith.edu/bio_facpubs)



Part of the [Biology Commons](#)

---

### Recommended Citation

Wright, Rachel M.; Strader, Marie E.; Genuise, Heather M.; and Matz, Mikhail, "Effects of Thermal Stress on Amount, Composition, and Antibacterial Properties of Coral Mucus" (2019). Biological Sciences: Faculty Publications, Smith College, Northampton, MA.

[https://scholarworks.smith.edu/bio\\_facpubs/179](https://scholarworks.smith.edu/bio_facpubs/179)

This Article has been accepted for inclusion in Biological Sciences: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact [scholarworks@smith.edu](mailto:scholarworks@smith.edu)



# Effects of thermal stress on amount, composition, and antibacterial properties of coral mucus

Rachel M. Wright<sup>1,2</sup>, Marie E. Strader<sup>2,3</sup>, Heather M. Genuise<sup>2</sup> and Mikhail Matz<sup>2</sup>

<sup>1</sup>Department of Genetics, Harvard Medical School, Boston, MA, United States of America

<sup>2</sup>Department of Integrative Biology, University of Texas at Austin, Austin, TX, United States of America

<sup>3</sup>Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA, United States of America

## ABSTRACT

The surface mucus layer of reef-building corals supports feeding, sediment clearing, and protection from pathogenic invaders. As much as half of the fixed carbon supplied by the corals' photosynthetic symbionts is incorporated into expelled mucus. It is therefore reasonable to expect that coral bleaching (disruption of the coral–algal symbiosis) would affect mucus production. Since coral mucus serves as an important nutrient source for the entire reef community, this could have substantial ecosystem-wide consequences. In this study, we examined the effects of heat stress-induced coral bleaching on the composition and antibacterial properties of coral mucus. In a controlled laboratory thermal challenge, stressed corals produced mucus with higher protein ( $\beta = 2.1$ ,  $p < 0.001$ ) and lipid content ( $\beta = 15.7$ ,  $p = 0.02$ ) and increased antibacterial activity (likelihood ratio = 100,  $p < 0.001$ ) relative to clonal controls. These results are likely explained by the expelled symbionts in the mucus of bleached individuals. Our study suggests that coral bleaching could immediately impact the nutrient flux in the coral reef ecosystem via its effect on coral mucus.

Submitted 13 July 2018  
Accepted 26 March 2019  
Published 29 April 2019

Corresponding author  
Rachel M. Wright,  
rachelwright8@gmail.com,  
rachel\_wright@hms.harvard.edu

Academic editor  
Craig Nelson

Additional Information and  
Declarations can be found on  
page 11

DOI 10.7717/peerj.6849

© Copyright  
2019 Wright et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Ecology, Marine Biology, Molecular Biology, Climate Change Biology

**Keywords** Coral reef, Mucus, Climate change, Coral bleaching, *Acropora cervicornis*

## INTRODUCTION

Rising sea surface temperature has increased the global risk of coral bleaching, the breakdown of the symbiosis between a coral host and its algal symbiont, to alarming levels (Hughes *et al.*, 2018). The direct impacts of bleaching on the animal host and algal symbiont are well studied. For example, coral bleaching has been shown to downregulate the expression of genes related to host immunity (Pinzón *et al.*, 2015) and alter host metabolism (Kenkel, Meyer & Matz, 2013; Rodrigues & Grottoli, 2007). Symbionts (family Symbiodiniaceae) expelled during bleaching produce elevated amounts of reactive oxygen species but are otherwise physiologically similar to healthy endosymbionts (Nielsen, Petrou & Gates, 2018). Sustained coral bleaching often leads to host mortality and subsequent shifts in the reef community structure (Harriott, 1985). For example, coral bleaching and mortality causes shifts in reef-fish assemblage structure and alters recruitment success (Booth & Beretta, 2002; Richardson *et al.*, 2018).


Coral mucus is a complex mixture of proteins, lipids, and carbohydrates that is produced by mucocytes in the coral epidermal layer and secreted by coral surface tissues (reviewed in [Brown & Bythell, 2005](#)). Arabinose and glucose primarily compose the monosaccharide portion of coral mucus ([Wild, Woyt & Huettel, 2005](#)). Arabinose is likely a symbiont-specific contribution to coral mucus, as aposymbiotic, heterotrophic cold-water corals produce mucus nearly identical in composition to mucus from symbiotic warm-water corals, except for the presence of arabinose in the latter ([Wild et al., 2010](#)). Up to about half of the photosynthetically fixed carbon supplied by a coral's algal symbiont is expelled as mucus ([Crossland, 1987](#); [Crossland, Barnes & Borowitzka, 1980](#); [Davies, 1984](#)). This coral surface mucus layer acts as a defense against desiccation and pathogens ([Shnit-Orland & Kushmaro, 2009](#)), and is also released into the water column where it traps suspended particles and acts as an energy source for benthic communities ([Wild et al., 2004a](#)). Given the integral role of photosynthetically fixed carbon in mucus production of most reef-building corals, it is predicted that coral bleaching events will reduce mucus production and subsequently impact the flow of energy throughout the reef ecosystem ([Bythell & Wild, 2011](#)). However, one study has shown that acroporid corals exposed to an experimental heat stress released almost twice as much dissolved and particulate organic carbon relative to unchallenged controls ([Niggel et al., 2009](#)). One explanation for this finding is that increased mucus release in heat-stressed corals augments defense against pathogens when other immune defense mechanisms may be compromised by thermal challenge ([Palmer, Bythell & Willis, 2010](#)).

In the Florida Keys, annual mass bleaching events are predicted to begin by the mid-century ([Manzello, 2015](#)). Currently, multiple anthropogenic factors including thermal stress, increased storms, and disease outbreaks have led to a near 80% decline in reef cover in the Florida Keys since the 1980s ([Porter et al., 2001](#); [Williams & Miller, 2011](#); [Williams, Miller & Kramer, 2008](#)). In particular, corals in the genus *Acropora* have faced some of the most dramatic declines in this region ([Greenstein, Curran & Pandolfi, 1998](#)). The staghorn coral, *Acropora cervicornis*, has been selected as a focal species for multiple active restoration programs, such as the Coral Restoration Foundation, due to its relatively fast asexual growth through fragmentation ([Lirman et al., 2010](#)). As restoration efforts aim to replenish stands of *A. cervicornis*, it is critical to assess this species' greater role in coral reef ecosystem. This study aims to characterize coral mucus release, composition, and antibacterial activity during acute thermal stress in *A. cervicornis* as a reference to guide further research on the potential consequences of bleaching-mediated mucus changes on the larger coral reef ecosystem.

## MATERIALS & METHODS

### Corals

Fifteen *Acropora cervicornis* genets ( $n = 3$  fragments per genet) were shipped from the Coral Restoration Foundation (Key Largo, Florida USA, Project ID CRF-2016-021) on 7 September 2016 to the University of Texas at Austin. Upon arrival, corals were immediately tagged with colored zip ties to uniquely identify each genet and allowed to recover for 12



**Figure 1** Experimental design and representative coral image. Corals were maintained in either (A) control (26 °C) or (B) experimental (31 °C) conditions for four days. Paling was observed for fragments in the experimental treatment, but not under control conditions.

Full-size  DOI: [10.7717/peerj.6849/fig-1](https://doi.org/10.7717/peerj.6849/fig-1)

days in artificial seawater (ASW; 30–31 ppt) at 25 °C under 12000K LED lights on a 12 h/12 h day/night cycle. Corals were fed weekly with Ultimate Coral Food (Coral Frenzy, LLC).

### Experimental conditions

Coral fragments were partitioned into experimental and control tanks. One genet (U10) experienced mortality during the recovery period, so only one U10 fragment remained when the experiment began. For all other genets, one fragment was placed in a control tank (26 °C) and one fragment was placed in an experimental tank. Any remaining fragments from the shipment of  $n = 3$  per genet were retained in a holding tank, though many genets developed tissue loss or experienced damage on a single fragment during shipping. The single remaining U10 fragment was placed in the experimental tank. The temperature in the experimental tank was ramped from 26 °C to 31 °C over 33 h. High summer temperature in the Florida Keys often reaches 31 °C (Manzello, 2015). Therefore, a 31 °C heat treatment was chosen to represent an ecologically relevant stressor.

After corals had been exposed to experimental conditions for four days, corals appeared visibly pale relative to initial photographs and paired control fragments (Fig. 1). At this time, the temperature in the experimental tank was reduced to 26 °C over 6 h.

### Image analysis

Prior to the experiment, photographs of each fragment were taken using a Nikon D5100 camera. Images of the front and back of each fragment were taken using the same camera,

settings, and lighting each day of the experiment. Brightness values in images were measured for the front and back sides of each fragment using image analysis software (ImageJ, [Schneider, Rasband & Eliceiri, 2012](#)). Corals become brighter (paler) as their symbioses with pigmented algae break down. Therefore, changes in coral brightness reflect changes in algal densities ([Winters et al., 2009](#)). A standard curve of brightness values was constructed using standard Coral Health Charts that were included in each image. Briefly, brightness (white value intensity measured in unitless values ranging 0–255) was measured for a series of standard color cards (arbitrary values 1–6) in each photograph to construct a linear curve of brightness to compare fragments. Brightness values were standardized to color cards to normalize for any minor differences in lighting across days.

### **Mucus collection**

After the experiment, each coral fragment was placed within a pre-weighed 50 mL conical tube containing 5 mL ASW from the respective tank. Tubes were placed on their sides and secured to a gently rocking incubator plate (135 RPM, 28 °C) for 20 min, rotating the tubes every 5 min to ensure that all sides of the coral fragment were submerged in water. After rocking, fragments were inverted dry above the liquid in the conical for 20 min and lightly centrifuged (200 RPM) for 2 min to pull down mucus adhering to the surface of the coral, modified from ([Wild et al., 2004a](#)). The volume and mass of mucus from each fragment was measured and stored at –80 °C. Coral fragments were returned to their tanks. The mucus collection procedure was repeated six days later, exactly as described above. During mucus collection, algal cells were clearly visible in some samples. All mucus aliquots were briefly centrifuged at 3,500 g for 2 min to sediment the algal pellet and decant algal cell-free mucus before the experiments described below.

### **Mucus composition**

Total protein was measured following the Coomassie (Bradford) Protein Assay Kit (Thermo Scientific, Waltham, MA, USA). Total carbohydrate was measured using the Total Carbohydrate Quantification Assay Kit (Abcam, Cambridge, UK). Total lipids were extracted and the dry mass of each mucus sample were measured. A standard curve was prepared using reagent grade cholesterol in a 2:1 chloroform:methanol mixture and an aliquot of 2:1 chloroform:methanol was added to each sample tube. After mixing, the solvent was evaporated from all standard and sample tubes on a heat block at 90 °C. Concentrated sulfuric acid was added to each tube, then incubated at 90 °C for 20 min. Samples were cooled, then plated in triplicate into wells of a 96-well plate. Background absorbance was measured at 540 nm. After incubating each sample with 50 µL of vanillin-phosphoric acid for 10 min, absorbance was measured again at 540 nm. The concentrations of protein, carbohydrate, and lipid in the mucus were estimated using standard curves constructed by measuring absorbance for known concentrations of protein, carbohydrate and lipids, per manufacturer's recommendations. All measurements were normalized to the volume of mucus expelled and the surface area of the fragment.

### Antibacterial activity

Cultures of laboratory *E. coli* (K-12) were grown overnight in Luria broth (LB), then washed twice in sterile ASW to remove remaining culture media. Coral mucus (140  $\mu$ L) and washed *E. coli* culture (60  $\mu$ L) was added to triplicate wells in 96-well plates. The covered plates were incubated at 37 °C for 12 h. Every 30 min the plate was shaken and the absorbance at 600 nm was measured. The plates were completely dry by the end of the incubation, so only the first 5 h, the latest time at which we observed liquid in the wells, are included in these analyses. Control wells included bacteria in LB or coral mucus without added bacteria to demonstrate bacterial growth or quantify endogenous mucus bacteria, respectively.

### Coral surface area

Fragment surface area was estimated using a 3D scanner and accompanying ScanStudioPro software (NextEngine, Santa Monica, CA, USA). Each scan was completed using a 360 degree scan with 16 divisions and 10,000 points/inch<sup>2</sup>. Scans were then trimmed, polished to fill holes, fused and then surface area was estimated based on a size standard.


### Real-time quantitative PCR

The forward primer 5'-TCTGTACGCCAACACTGTGCTT-3' and reverse primer 5'-AGTGATGCCAAGATGGAGCCT-3' was used to amplify the *Acropora cervicornis* actin sequence as developed in (Winter, 2017). The forward primer 5'-GTGAATTGCAGAACTCCGTG-3' and reverse primer 5'-CCTCCGTTACTTATATGCTT-3' was used to amplify the Symbiodiniaceae ITS2 sequence. Primer pair specificity was verified by gel electrophoresis and melt curve analysis of the amplification product obtained with *A. cervicornis* holobiont DNA. Primer efficiencies were determined by amplifying a series of four-fold dilutions of *A. cervicornis* holobiont DNA and analyzing the results using *PrimEff* function in the *MCMC.qpcr* package (Matz, Wright & Scott, 2013) in R. Briefly,  $C_T$  (threshold cycle) results were plotted as CT vs.  $\log_2$ [DNA], and amplification efficiencies (amplification factor per cycle) of each primer pair were derived from the slope of the regression using formula: efficiency =  $2^{-(1/\text{slope})}$  (Pfaffl, 2001).

Mucus aliquots were centrifuged to remove any cell debris. A 14  $\mu$ L aliquot of coral mucus was combined with SYBR Green PCR Master Mix (Applied Biosystems), 1.5  $\mu$ M forward and reverse primers, and water. The Roche LightCycler 480 system was used to carry out the PCR protocol (95 °C for 40 s, then 40 cycles of 60 °C for 1 min and 72 °C for 1 min) and detect the fluorescence signal.

### Statistics

All statistical analyses were performed in R (3.4.0, R Core Team, 2016). Bayesian models implemented using the MCMCglmm package (Hadfield, 2010) was used to explain variation in coral color and mucus composition, with treatment as a fixed effect and genotype as a random effect under a Gaussian distribution for coral color and metabolic assays and a Poisson distribution for marker gene counts. A nonlinear mixed-effects model followed by ANOVA was implemented using the nlme package (Pinheiro et al., 2017) was used for time-series analysis of antibacterial activity, with time, treatment, and their interaction



**Figure 2** Mucus production. Mucus was collected immediately after paling was observed (“Initial”) and six days after the challenged corals were returned to control conditions (“Recovery”). The volume (A) and mass (B) of the recovered mucus was normalized to the surface area of the coral fragment. Red boxes represent fragments in control conditions; beige boxes represent heat-stressed fragments. Error bars represent standard error.

Full-size DOI: [10.7717/peerj.6849/fig-2](https://doi.org/10.7717/peerj.6849/fig-2)

as fixed effects and plate well as a random effect. Experimental data and R scripts are included as [Data S1](#) and [S2](#), respectively. Data and scripts can also be accessed on GitHub: [https://github.com/rachelwright8/bleached\\_coral\\_mucus](https://github.com/rachelwright8/bleached_coral_mucus).


## RESULTS

### Coral bleaching and mucus collection

Corals from both treatments produced similar amounts of mucus by volume ( $\beta = 0.01$ ,  $p = 0.25$ , [Fig. 2A](#)) and mass ( $\beta = 0.14$ ,  $p = 0.27$ , [Fig. 2B](#)). After a six-day recovery period, corals in both treatments produced significantly less mucus by volume ( $\beta = -0.14$ ,  $p < 0.001$ , [Fig. 2A](#)) and mass ( $\beta = -1.1$ ,  $p < 0.001$ , [Fig. 2B](#)) compared to the first time point. After four days in the experimental treatment at 31 °C, corals paled significantly compared to corals in the control condition ( $\beta = -1.16$ ,  $p < 0.001$ , [Fig. 3A](#)).

### Mucus biochemistry

The mucus produced by heat-stressed fragments contained significantly more total protein ( $\beta = 2.1$ ,  $p < 0.001$ , [Fig. 3B](#)) and total lipid ( $\beta = 15.7$ ,  $p = 0.02$ , [Fig. 3C](#)). There was also a slight increase in carbohydrate content in mucus from heat-stressed corals, though this difference was not significant ( $\beta = 0.64$ ,  $p = 0.10$ , [Fig. 3D](#)).




**Figure 3** Effects of heat stress. (A) Effect of treatment on coral color (decrease in color score indicates bleaching). (B–D) Effects on mucus composition: protein (B), in mg/cm<sup>2</sup> fragment surface area, and lipid (C) and carbohydrate (D), in µg/cm<sup>2</sup> fragment surface area. Red boxes represent fragments in control conditions; beige boxes represent heat-stressed fragments. Error bars represent standard error.

Full-size [DOI: 10.7717/peerj.6849/fig-3](https://doi.org/10.7717/peerj.6849/fig-3)

### Mucus antibacterial activity

Antibacterial activity increased in thermally stressed corals from the experimental treatment relative to healthy corals (likelihood ratio = 100,  $p < 0.001$ , Fig. 4). Optical density at 600 nm (OD<sub>600</sub>), an absorbance measure that reflects bacterial density, decreased throughout the incubation period in all mucus samples. However, bacterial density declined significantly faster in mucus samples from heat-stressed corals. Bacterial density remained low throughout the sampling time period in negative control samples, which contained mucus from control corals with no added bacteria, suggesting that decreases in observed bacterial density were due to elimination of the experimentally added *E. coli* (Fig. S1). Positive control wells that contained *E. coli* in growth media with no coral mucus increased in OD, showing that the bacteria were alive and capable of growth under the experimental conditions (Fig. S1).



**Figure 4** Antibacterial activity of coral mucus. Optical density (OD) at 600 nm indicates the density of inoculated bacteria in coral mucus samples. Red lines represent fragments in control conditions; beige lines represent heat-stressed fragments. Error bars represent standard error.

Full-size DOI: [10.7717/peerj.6849/fig-4](https://doi.org/10.7717/peerj.6849/fig-4)


### Presence of host and symbiont DNA in mucus

Real-time quantitative PCR (qPCR) was performed to determine the relative abundances of coral- and symbiont-derived marker gene copies present in the mucus released by healthy and heat-stressed corals. Primers were designed to target a coral-specific actin gene and the Symbiodiniaceae ITS2 region. Presumably, copies of the coral-specific actin gene would represent lysed coral cells, while copies of the ITS2 region would represent material released from Symbiodiniaceae cells in the mucus. We did not detect any ITS2 sequences in the mucus of unchallenged corals (Fig. 5). Mucus released by heat-stressed coral fragments contained significantly more marker gene copies of both coral-derived (95% credible interval (CI) = 1.6–4.6,  $p = 0.001$ ) and symbiont-derived sequences (95% CI [4.3–25.7],  $p < 0.001$ , Fig. 5).

## DISCUSSION

### Coral mucus stores take a long time to replenish

Mucus volumes released from both control and heat-stressed fragments during the initial collection ( $0.18 \pm 0.04$  mL/cm<sup>2</sup>, or about 1.8 L/m<sup>2</sup>) are consistent with daily mucus release values previously reported for submerged acroporid corals (1.7 L/m<sup>2</sup> in Wild *et al.*, 2004a). However, following a six-day recovery period, less than half of the original mucus volume was collected from both control and heat-stressed fragments ( $0.18 \pm 0.04$  mL/cm<sup>2</sup> initial



**Figure 5** Symbiodiniaceae and coral DNA in mucus. Real-time quantitative PCR detected *A. cervicornis* actin (A) or Symbiodiniaceae ITS2 (B) marker gene copies in the coral mucus of healthy and pale corals. Red boxes represent fragments in control conditions; beige boxes represent heat-stressed fragments. Error bars represent standard error.

Full-size [DOI: 10.7717/peerj.6849/fig-5](https://doi.org/10.7717/peerj.6849/fig-5)

collection vs.  $0.08 \pm 0.02$  mL/cm<sup>2</sup> post-recovery,  $p_{\text{MCMCglmm}} < 0.001$ , Fig. 2A), suggesting that mucus stores were not completely replenished to the original volume in this amount of time. The methods used in this study to extract mucus left the coral nubbins completely dry. Therefore, the results presented here represent the total volume of stored mucus at the time of collection, which may not correspond directly to daily mucus release in natural settings. These results emphasize the importance of measuring mucus release over time to confidently estimate daily release rates and predict daily energetic flow throughout the reef ecosystem.

### Stressed corals produce mucus high in protein and lipid

We found no difference in the quantity of mucus produced by stressed corals compared to healthy corals after four days of heat stress or after a six-day recovery period (Fig. 2), suggesting that the quantity of mucus produced by a coral is relatively unaffected by thermal stress and that mucus stores cannot be replenished within a week. Given the energetic cost of producing mucus (Riegl & Branch, 1995), it is reasonable to predict that corals with low densities of autotrophic symbionts would produce less, or lower nutritional quality, mucus than healthy corals. These findings of enhanced protein and lipid content in mucus from heat-stressed corals relative to mucus produced by healthy corals (Fig. 3) support a previous study that found increased organic carbon release by thermally stressed corals (Niggel et al., 2009). Symbiodiniaceae store reserve energy as lipid

droplets and starch granules that are translocated from the algal membrane to coral cells in a healthy coral–algae symbiotic relationship (Patton & Burris, 1983). We observed a pellet of algal cells in the mucus of thermally stressed corals, but not in mucus produced by healthy corals. Though algal cells were pelleted and removed from all mucus collections, extracellular lipid droplets would remain in the mucus and could represent a potential explanation for the increased abundance of lipids in mucus from stressed corals. Likewise, proteins and lipids released from damaged Symbiodiniaceae and host cells would also be present in the mucus of stressed corals. This particular possibility is supported by finding both coral and Symbiodiniaceae DNA in the mucus of stressed corals (Fig. 5). Future studies should measure long-term effects of bleaching to determine the duration of this observed enrichment in coral mucus quality following thermal stress.

This short-term enrichment in nutrition could invoke shifts in trophic interactions and density dependent foraging behavior in corallivorous fish and invertebrates. In the Florida Keys, previous studies have observed increased predation intensity by corallivorous fishes and invertebrates as coral cover declines (Baums, Miller & Szmant, 2003; Burkepile, 2012). In addition, there is established prey preference asymmetries in the Caribbean, where *A. cervicornis* is already the preferred prey of a corallivorous gastropod (Johnston & Miller, 2014). Our results implicate that as heat-stressed corals release more nutrients into their mucus, this may drive stronger predation pressure, at least in bursts associated with the bleaching event itself. This may only maintain a benefit to higher trophic levels at very short time scales, as predation on coral reef dwelling fish is higher when they reside in bleached corals (Coker, Pratchett & Munday, 2009) and corallivorous fish are well known to suffer population declines in response to coral bleaching (Pratchett et al., 2018). Therefore, future studies should investigate how bleaching impacts predator–prey dynamics between corals and corallivorous fish and invertebrates.

### **Stressed corals produce mucus with high antibacterial activity**

This study found that mucus collected from stressed coral fragments eliminated bacteria faster than mucus from healthy fragments from matched genets (Fig. 4). Antibacterial activity of coral mucus is attributed to antimicrobial substances produced by commensal microbes living on the coral surface (Nissimov, Rosenberg & Munn, 2009; Shnit-Orland & Kushmaro, 2009). A previous study of mucus collected from *A. palmata* during a summer bleaching event in the Florida Keys found that mucus from bleached corals lacked antibiotic properties normally found in healthy coral mucus (Ritchie, 2006). The difference in findings between this study and the investigation of antibiotic activity in naturally bleached corals could be attributed to the timing of collections. Mucus in this study was collected as soon as corals became pale, whereas the previous study collected mucus after corals in the Florida Keys had been experiencing high levels of thermal stress and bleaching for about a month according to NOAA reports of the thermal stress event (Eakin et al., 2010). Long-term thermal stress is known to promote coral disease by altering bacterial pathogenicity and host susceptibility (Bruno et al., 2007; Maynard et al., 2015). In our short-term bleaching conditions, the increased protein and lipid content in the mucus (Figs. 3B–3C) may have temporarily improved the antibacterial activity of commensal

microbes that exist in the coral mucus (*Krediet et al., 2013*). Though the mechanism is unclear, Symbiodiniaceae do appear to play a role in a coral's ability to manage immune stress and regulate microbial communities (*Littman, Bourne & Willis, 2010; Wright et al., 2017*).

Other studies that have investigated bacterial growth in coral mucus have found that mucus supports the growth of bacterial species found on corals (*Wild et al., 2004b*), such as *Pseudoalteromonas* and *Vibrio* spp. (*Allers et al., 2008*). Both *Pseudoalteromonas* and *Vibrio* are commonly observed in corals, though they are usually associated with coral stress and disease suggesting that these bacteria employ mechanisms to resist antimicrobial properties of coral mucus (*Ritchie, 2006*) that our lab *E. coli* strain, which was naïve to coral mucus, presumably lacks. This difference in bacterial species could explain the discrepancy in findings. Additionally, the previous experiments measured bacterial growth over 40–50 h. It is possible that we would have found similar results had we measured bacterial growth beyond 5 h.

## CONCLUSIONS

Our results show that thermal stress does not significantly affect the volume of mucus produced by *A. cervicornis* immediately following a bleaching event. Stressed corals produced mucus with higher protein content, higher lipid content, and increased antibacterial activity relative to unstressed controls. Additional lipids and proteins likely come from Symbiodiniaceae and host cells damaged during bleaching rather than from additional investment by the coral host. Short-term nutritional enrichments of mucus released from bleaching corals could promote growth of heterotrophic microbes at the lowest trophic levels of marine ecosystems and thus cause large-scale shifts in a reef's nutrient cycle. Changes in the nutritional composition and antibacterial properties of the mucus should influence coral-associated microbes that contribute to coral disease susceptibility. Future experiments should investigate longer-term effects of thermal stress on mucus production and content to further investigate reef-wide consequences of coral bleaching.

## ACKNOWLEDGEMENTS

We thank the Coral Reef Foundation for providing coral specimen and Sarah Davies for measuring coral surface areas.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was funded by a 2016 PADI grant #21956 awarded to Marie Strader and a University of Texas Co-op Undergraduate Research Fellowship awarded to Heather Genuise. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Grant Disclosures

The following grant information was disclosed by the authors:

2016 PADI: #21956.

University of Texas Co-op Undergraduate Research.

## Competing Interests

The authors declare there are no competing interests.

## Author Contributions

- Rachel M. Wright conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Marie E. Strader conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Heather M. Genuise performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Mikhail Matz conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

## Data Availability

The following information was supplied regarding data availability:

The data and R scripts are available at [https://github.com/rachelwright8/bleached\\_coral\\_mucus](https://github.com/rachelwright8/bleached_coral_mucus).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6849#supplemental-information>.

## REFERENCES

- Allers E, Niesner C, Wild C, Pernthaler J. 2008.** Microbes enriched in seawater after addition of coral mucus. *Applied and Environmental Microbiology* **74**(10):3274–3278 DOI [10.1128/AEM.01870-07](https://doi.org/10.1128/AEM.01870-07).
- Baums IB, Miller MW, Szmant AM. 2003.** Ecology of a corallivorous gastropod, *Coralliophila abbreviata*, on two scleractinian hosts, I: population structure of snails and corals. *Marine Biology* **142**(6):1083–1091 DOI [10.1007/s00227-003-1024-9](https://doi.org/10.1007/s00227-003-1024-9).
- Booth DJ, Beretta GA. 2002.** Changes in a fish assemblage after a coral bleaching event. *Marine Ecology Progress Series* **245**:205–212 DOI [10.3354/meps245205](https://doi.org/10.3354/meps245205).
- Brown B, Bythell J. 2005.** Perspectives on mucus secretion in reef corals. *Marine Ecology Progress Series* **296**:291–309 DOI [10.3354/meps296291](https://doi.org/10.3354/meps296291).
- Bruno JF, Selig ER, Casey KS, Page CA, Willis BL, Harvell CD, Sweatman H, Melendy AM. 2007.** Thermal stress and coral cover as drivers of coral disease outbreaks. *PLOS Biology* **5**(6):1220–1227 DOI [10.1371/journal.pbio.0050124](https://doi.org/10.1371/journal.pbio.0050124).

- Burkepile DE. 2012.** Context-dependent corallivory by parrotfishes in a Caribbean reef ecosystem. *Coral Reefs* **31**(1):111–120 DOI [10.1007/s00338-011-0824-5](https://doi.org/10.1007/s00338-011-0824-5).
- Bythell JC, Wild C. 2011.** Biology and ecology of coral mucus release. *Journal of Experimental Marine Biology and Ecology* **408**(1–2):88–93 DOI [10.1016/j.jembe.2011.07.028](https://doi.org/10.1016/j.jembe.2011.07.028).
- Coker DJ, Pratchett MS, Munday PL. 2009.** Coral bleaching and habitat degradation increase susceptibility to predation for coral-dwelling fishes. *Behavioral Ecology* **20**(6):1204–1210 DOI [10.1093/beheco/arp113](https://doi.org/10.1093/beheco/arp113).
- Crossland CJ. 1987.** *In situ* release of mucus and DOC-lipid from the corals *Acropora variabilis* and *Stylophora pistillata* in different light regimes. *Coral Reefs* **6**(1):35–42 DOI [10.1007/BF00302210](https://doi.org/10.1007/BF00302210).
- Crossland CJ, Barnes DJ, Borowitzka MA. 1980.** Diurnal lipid and mucus production in the staghorn coral *Acropora acuminata*. *Marine Biology* **60**(2–3):81–90 DOI [10.1007/BF00389151](https://doi.org/10.1007/BF00389151).
- Davies PS. 1984.** The role of zooxanthellae in the nutritional energy requirements of *Pocillopora eydouxi*. *Coral Reefs* **2**(4):181–186 DOI [10.1007/BF00263571](https://doi.org/10.1007/BF00263571).
- Eakin CM, Morgan JA, Heron SF, Smith TB, Liu G, Alvarez-Filip L, Baca B, Bartels E, Bastidas C, Bouchon C, Brandt M, Bruckner AW, Bunkley-Williams L, Cameron A, Causey BD, Chiappone M, Christensen TR, Crabbe MJ, Day O, De la Guardia E, Díaz-Pulido G, DiResta D, Gil-Agudelo DL, Gilliam DS, Ginsburg RN, Gore S, Guzmán HM, Hendee JC, Hernández-Delgado EA, Husain E, Jeffrey CF, Jones RJ, Jordán-Dahlgren E, Kaufman LS, Kline DI, Kramer PA, Lang JC, Lirman D, Mallela J, Manfrino C, Maréchal JP, Marks K, Mihaly J, Miller WJ, Mueller EM, Muller EM, Orozco Toro CA, Oxenford HA, Ponce-Taylor D, Quinn N, Ritchie KB, Rodríguez S, Ramírez AR, Romano S, Samhoury JF, Sánchez JA, Schmahl GP, Shank BV, Skirving WJ, Steiner SC, Villamizar E, Walsh SM, Walter C, Weil E, Williams EH, Roberson KW, Yusuf Y. 2010.** Caribbean corals in crisis: record thermal stress, bleaching, and mortality in 2005. *PLOS ONE* **5**(11):e13969 DOI [10.1371/journal.pone.0013969](https://doi.org/10.1371/journal.pone.0013969).
- Greenstein BJ, Curran HA, Pandolfi JM. 1998.** Shifting ecological baselines and the demise of *Acropora cervicornis* in the western North Atlantic and Caribbean Province: a Pleistocene perspective. *Coral Reefs* **17**(3):249–261 DOI [10.1007/s003380050125](https://doi.org/10.1007/s003380050125).
- Hadfield JD. 2010.** MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software* **33**(2):1–22 DOI [10.1002/ana.22635](https://doi.org/10.1002/ana.22635).
- Harriott VJ. 1985.** During a mass bleaching event. *Marine Ecology Progress Series* **21**(1):81–88 DOI [10.3354/meps021081](https://doi.org/10.3354/meps021081).
- Hughes TP, Kerry JT, Baird AH, Connolly SR, Dietzel A, Eakin CM, Heron SF, Hoey AS, Hoogenboom MO, Liu G, McWilliam MJ, Pears RJ, Pratchett MS, Skirving WJ, Stella JS, Torda G. 2018.** Global warming transforms coral reef assemblages. *Nature* **556**:492–496 DOI [10.1038/s41586-018-0041-2](https://doi.org/10.1038/s41586-018-0041-2).
- Johnston L, Miller MW. 2014.** Negative indirect effects of neighbors on imperiled scleractinian corals. *Coral Reefs* **33**(4):1047–1056 DOI [10.1007/s00338-014-1176-8](https://doi.org/10.1007/s00338-014-1176-8).

- Kenkel CD, Meyer E, Matz MV. 2013.** Gene expression under chronic heat stress in populations of the mustard hill coral (*Porites astreoides*) from different thermal environments. *Molecular Ecology* **22**(16):4322–4334 DOI [10.1111/mec.12390](https://doi.org/10.1111/mec.12390).
- Krediet CJ, Ritchie KB, Paul VJ, Teplitski M. 2013.** Coral-associated micro-organisms and their roles in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B: Biological Sciences* **280**(1755):20122328 DOI [10.1098/rspb.2012.2328](https://doi.org/10.1098/rspb.2012.2328).
- Lirman D, Thyberg T, Herlan J, Hill C, Young-Lahiff C, Schopmeyer S, Huntington B, Santos R, Drury C. 2010.** Propagation of the threatened staghorn coral *Acropora cervicornis*: methods to minimize the impacts of fragment collection and maximize production. *Coral Reefs* **29**(3):729–735 DOI [10.1007/s00338-010-0621-6](https://doi.org/10.1007/s00338-010-0621-6).
- Littman RA, Bourne DG, Willis BL. 2010.** Responses of coral-associated bacterial communities to heat stress differ with Symbiodinium type on the same coral host. *Molecular Ecology* **19**(9):1978–1990 DOI [10.1111/j.1365-294X.2010.04620.x](https://doi.org/10.1111/j.1365-294X.2010.04620.x).
- Manzello DP. 2015.** Rapid recent warming of coral reefs in the Florida Keys. *Scientific Reports* **5**:1–10 DOI [10.1038/srep16762](https://doi.org/10.1038/srep16762).
- Matz MV, Wright RM, Scott JG. 2013.** No control genes required: Bayesian analysis of qRT-PCR data. *PLOS ONE* **8**(8):e71448 DOI [10.1371/journal.pone.0071448](https://doi.org/10.1371/journal.pone.0071448).
- Maynard J, Van Hooidek R, Eakin CM, Puotinen M, Garren M, Williams G, Heron SF, Lamb J, Weil E, Willis B, Harvell CD. 2015.** Projections of climate conditions that increase coral disease susceptibility and pathogen abundance and virulence. *Nature Climate Change* **5**:688–695 DOI [10.1038/nclimate2625](https://doi.org/10.1038/nclimate2625).
- Nielsen DA, Petrou K, Gates RD. 2018.** Coral bleaching from a single cell perspective. *ISME Journal* **12**:1558–1567 DOI [10.1038/s41396-018-0080-6](https://doi.org/10.1038/s41396-018-0080-6).
- Niggel W, Glas MS, Laforsch C, Mayr C, Wild C. 2009.** First evidence of coral bleaching stimulating organic matter release by reef corals. In: *Proceedings of the 11th international coral reef symposium, (July)*. 905–910.
- Nissimov J, Rosenberg E, Munn CB. 2009.** Antimicrobial properties of resident coral mucus bacteria of *Oculina patagonica*. *FEMS Microbiology Letters* **292**(2):210–215 DOI [10.1111/j.1574-6968.2009.01490.x](https://doi.org/10.1111/j.1574-6968.2009.01490.x).
- Palmer CV, Bythell JC, Willis BL. 2010.** Levels of immunity parameters underpin bleaching and disease susceptibility of reef corals. *Federation of American Societies for Experimental Biology* **24**(6):1935–1946 DOI [10.1096/fj.09-152447](https://doi.org/10.1096/fj.09-152447).
- Patton JS, Burriss JE. 1983.** Lipid synthesis and extrusion by freshly isolated zooxanthellae (symbiotic algae). *Marine Biology* **75**(2–3):131–136 DOI [10.1007/BF00405995](https://doi.org/10.1007/BF00405995).
- Pfaffl MW. 2001.** A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research* **29**(9):e45 DOI [10.1093/nar/29.9.e45](https://doi.org/10.1093/nar/29.9.e45).
- Pinheiro J, Bates D, DebRoy S, Sarkar D. 2017.** nlme: linear and nonlinear mixed effects models.
- Pinzón JH, Kamel B, Burge CA, Harvell CD, Medina M, Weil E, Mydlarz LD. 2015.** Whole transcriptome analysis reveals changes in expression of immune-related genes during and after bleaching in a reef-building coral. *Royal Society Open Science* **2** DOI [10.1098/rsos.140214](https://doi.org/10.1098/rsos.140214).

- Porter JW, Dustan P, Jaap WC, Patterson KL, Kosmynin V, Meier OW, Patterson ME, Parsons M. 2001. Patterns of spread of coral disease in the Florida Keys. *Hydrobiologia* 460:1–24 DOI 10.1023/A:1013177617800.
- Pratchett MS, Thompson CA, Hoey AS, Cowman PF, Wilson SK. 2018. Effects of coral bleaching and coral loss on the structure and function of reef fish assemblages. In: Van Oppen M, Lough J, eds. *Coral bleaching. Ecological studies (analysis and synthesis)*. Vol 233. Springer, Cham, 265–293.
- R Core Team. 2016. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org/>.
- Richardson LE, Graham NAJ, Pratchett MS, Eurich JG, Hoey AS. 2018. Mass coral bleaching causes biotic homogenization of reef fish assemblages. *Global Change Biology* 24(7):3117–3129 DOI 10.1111/gcb.14119.
- Riegl B, Branch GM. 1995. Effects of sediment on the energy budgets of four scleractinian (Bourne 1900) and five alcyonacean (Lamouroux 1816) corals. *Journal of Experimental Marine Biology and Ecology* 186:259–275 DOI 10.1016/0022-0981(94)00164-9.
- Ritchie K. 2006. Regulation of microbial populations by coral surface mucus and mucus-associated bacteria. *Marine Ecology Progress Series* 322:1–14 DOI 10.3354/meps322001.
- Rodrigues LJ, Grottoli AG. 2007. Energy reserves and metabolism as indicators of coral recovery from bleaching. *Limnology and Oceanography* 52(5):1874–1882 DOI 10.4319/lo.2007.52.5.1874.
- Schneider C, Rasband W, Eliceiri K. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9(7):671–675 DOI 10.1038/nmeth.2089.
- Shnit-Orland M, Kushmaro A. 2009. Coral mucus-associated bacteria: a possible first line of defense. *FEMS Microbiology Ecology* 67(3):371–380 DOI 10.1111/j.1574-6941.2008.00644.x.
- Wild C, Huettel M, Klueter A, Kremb SG, Rasheed MYM, Jørgensen BB. 2004a. Coral mucus functions as an energy carrier and particle trap in the reef ecosystem. *Nature* 428(6978):66–70 DOI 10.1038/nature02344.
- Wild C, Naumann M, Niggel W, Haas A. 2010. Carbohydrate composition of mucus released by scleractinian warm- and cold-water reef corals. *Aquatic Biology* 10(1):41–45 DOI 10.3354/ab00269.
- Wild C, Rasheed M, Werner U, Franke U, Johnstone R, Huettel M. 2004b. Degradation and mineralization of coral mucus in reef environments. *Marine Ecology Progress Series* 267:159–171 DOI 10.3354/meps267159.
- Wild C, Woyt H, Huettel M. 2005. Influence of coral mucus on nutrient fluxes in carbonate sands. *Marine Ecology Progress Series* 287(2014):87–98 DOI 10.3354/meps287087.
- Williams DE, Miller MW. 2011. Attributing mortality among drivers of population decline in *Acropora palmata* in the Florida Keys (USA). *Coral Reefs* 31(2):369–382 DOI 10.1007/s00338-011-0847-y.

- Williams DE, Miller MW, Kramer KL. 2008.** Recruitment failure in Florida Keys *Acropora palmata*, a threatened Caribbean coral. *Coral Reefs* **27**(3):697–705  
DOI [10.1007/s00338-008-0386-3](https://doi.org/10.1007/s00338-008-0386-3).
- Winter RN. 2017.** Environmental controls on the reassembly of symbiodinium communities in reef corals following perturbation: implications for reef futures under climate change. Available at [https://scholarlyrepository.miami.edu/oa\\_dissertations/1870](https://scholarlyrepository.miami.edu/oa_dissertations/1870).
- Winters G, Holzman R, Blehman A, Beer S, Loya Y. 2009.** Photographic assessment of coral chlorophyll contents: implications for ecophysiological studies and coral monitoring. *Journal of Experimental Marine Biology and Ecology* **380**(1–2):25–35  
DOI [10.1016/j.jembe.2009.09.004](https://doi.org/10.1016/j.jembe.2009.09.004).
- Wright RM, Kenkel CD, Dunn CE, Shilling EN, Bay LK, Matz MV. 2017.** Intraspecific differences in molecular stress responses and coral pathobiome contribute to mortality under bacterial challenge in *Acropora millepora*. *Scientific Reports* **7**(1):2609  
DOI [10.1038/s41598-017-02685-1](https://doi.org/10.1038/s41598-017-02685-1).