
8-1-2007

Human Body Temperature (37°C) Increases the Expression of Iron, Carbohydrate, and Amino Acid Utilization Genes in *Escherichia coli* K-12

Christine A. White-Ziegler
Smith College, cwhitezi@smith.edu

Amy J. Malhowski
Smith College

Sarah Young
Smith College

Follow this and additional works at: https://scholarworks.smith.edu/bio_facpubs

 Part of the [Biology Commons](#)

Recommended Citation

White-Ziegler, Christine A.; Malhowski, Amy J.; and Young, Sarah, "Human Body Temperature (37°C) Increases the Expression of Iron, Carbohydrate, and Amino Acid Utilization Genes in *Escherichia coli* K-12" (2007). Biological Sciences: Faculty Publications, Smith College, Northampton, MA.
https://scholarworks.smith.edu/bio_facpubs/242

This Article has been accepted for inclusion in Biological Sciences: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact scholarworks@smith.edu

Human Body Temperature (37°C) Increases the Expression of Iron, Carbohydrate, and Amino Acid Utilization Genes in *Escherichia coli* K-12[∇]

Christine A. White-Ziegler,* Amy J. Malhowski, and Sarah Young

Department of Biological Sciences and Program in Biochemistry, Smith College, Northampton, Massachusetts

Received 21 December 2006/Accepted 15 May 2007

Using DNA microarrays, we identified 126 genes in *Escherichia coli* K-12 whose expression is increased at human body temperature (37°C) compared to growth at 23°C. Genes involved in the uptake and utilization of amino acids, carbohydrates, and iron dominated the list, supporting a model in which temperature serves as a host cue to increase expression of bacterial genes needed for growth. Using quantitative real-time PCR, we investigated the thermoregulatory response for representative genes in each of these three categories (*hisJ*, *cysP*, *srlE*, *garP*, *fes*, and *cirA*), along with the fimbrial gene *papB*. Increased expression at 37°C compared to 23°C was retained in both exponential and stationary phases for all of the genes and in most of the various media tested, supporting the relative importance of this cue in adapting to changing environments. Because iron acquisition is important for both growth and virulence, we analyzed the regulation of the iron utilization genes *cirA* and *fes* and found that growth in iron-depleted medium abrogated the thermoregulatory effect, with high-level expression at both temperatures, contrasting with *papB* thermoregulation, which was not greatly altered by limiting iron levels. A positive role for the environmental regulator H-NS was found for *fes*, *cirA*, *hisJ*, and *srlE* transcription, whereas it had a primarily negative effect on *cysP* and *garP* expression. Together, these studies indicate that temperature is a broadly used cue for regulating gene expression in *E. coli* and that H-NS regulates iron, carbohydrate, and amino acid utilization gene expression.

Bacteria have the ability to keenly sense a multitude of environmental stimuli, such as temperature, pH, osmolarity, oxygen levels, carbon sources, and concentrations of various ions and compounds, including iron (reviewed in references 11, 33, and 36). Elaborate sensory mechanisms have evolved in order for these microorganisms to detect environmental changes and thereby regulate cellular activities to adapt to their changing surroundings. This is particularly evident among bacterial pathogens, in which expression of virulence factors is highly regulated in response to the surrounding environment (reviewed in references 33 and 36). Pathogenic bacteria use environmental cues to distinguish between host and nonhost environments, and subsequent regulation of virulence gene expression can allow a more efficient utilization of resources. Environmental regulation of virulence gene expression is of particular importance in *Escherichia coli*, which can colonize the gut or translocate from that competitive environment to other niches, such as the sterile urinary tract.

Our laboratory has studied the effect of mammalian host temperature (37°C) in controlling pyelonephritis-associated pili, otherwise known as Pap or P fimbriae, that are expressed by uropathogenic strains of *E. coli* and are encoded by the *pap* operon (43, 44). P fimbriae are critical in the colonization of the host upper urinary tract, leading to acute upper urinary tract infections (43, 44). The thermoregulation of *papBA* transcription has been well characterized in the uropathogenic strain and at the molecular level as a transcriptional fusion in

E. coli K-12 (4, 21–23, 70–74). Optimal expression occurs at 37°C, with a 52-fold reduction in *papBA* transcription at 23°C (70). The environmental cue of temperature is important, not just for *pap*, but in the control of a number of virulence genes in *E. coli*, including those encoding fimbriae and toxins (14, 17, 18, 20, 26, 34, 39, 50, 56, 74).

Two proteins, RimJ and H-NS, are known to be important in the regulation of *papBA* transcription in response to temperature and other environmental conditions. RimJ is the N-terminal acetyltransferase of the ribosomal protein S5 (12), and deletion of the *rimJ* gene leads to a loss of thermoregulation so that *papBA* transcription levels are equivalent at both 37°C and 23°C (71). The mechanism by which RimJ represses *papBA* transcription and how the modification of a ribosomal protein might be involved in this process are currently under investigation. H-NS is a histone-like nucleoid-structuring protein that binds to A-T-rich, bent regions of DNA and compacts them (1, 13, 66, 75). H-NS controls the expression of a number of environmentally controlled virulence genes in several gram-negative genera (1). In contrast to many other virulence genes in which H-NS plays only a repressive role, the protein plays both a positive and a negative role in *papBA* transcription (67, 70, 74). At 37°C, introduction of the *hns65I* mutation leads to a decrease in transcription, suggesting that H-NS is required for maximal transcription at this temperature. At 23°C, *papBA* transcription in the *hns65I* mutant strain is higher than in the wild-type strain and approaches the levels observed at 37°C for the mutant strain, indicating a loss of thermoregulation.

In this study, we investigated the genomewide effect of temperature (37°C) on transcription in *E. coli* K-12 to identify other genes that are regulated in a manner similar to that of *pap* and thus might be important in adaptation to and coloni-

* Corresponding author. Mailing address: Department of Biological Sciences, Smith College, Northampton, MA 01063. Phone: (413) 585-3815. Fax: (413) 585-3786. E-mail: cwhitezi@smith.edu.

[∇] Published ahead of print on 25 May 2007.

TABLE 1. Bacterial strains, bacteriophages, primers, and probes used in this study

Strains and bacteriophages	Description	Reference or source
<i>E. coli</i> strains		
MC4100	F ⁻ <i>araD139</i> Δ (<i>lacIPOZYA-argF</i>) <i>U169 rpsL thi-1</i>	9
H1941	MC4100 <i>fur</i>	K. Hantke
NH757	B178 <i>hns651 tyrT</i> β ::Tn10	19
DL1504	MC4100 λ 354 lysogen	5
DL1947	DL1504 <i>hns651</i>	67
CWZ388	DL1504 Δ <i>rimJ</i>	71
CWZ479	H1941 λ 354 lysogen	This study
Bacteriophages		
P1L4	Virulent phage P1	D. A. Low
λ 354	<i>papBA-lacZYA</i> fusion phage	5

zation of the host. Transcriptome analyses based on temperature have been conducted in group A *Streptococcus* (37° and 29°C) (60), *Yersinia pestis* (37°C and 26°C) (24, 38), and *Borrelia burgdorferi* (37°C and 23°C) (7, 52), but to our knowledge, this is the first investigation and analysis of *E. coli*. Here, we present evidence that 126 genes are increased in expression at 37°C, with a significant number of carbohydrate, amino acid, and iron utilization genes represented within this list. We further characterize the thermoregulatory responses for representative genes within these three categories under a variety of growth conditions and demonstrate that, in the majority of cases, a temperature differential is maintained that favors increased expression of these genes at 37°C compared to growth at 23°C. Lastly, we demonstrate that H-NS controls the amino acid, carbohydrate, and iron utilization genes tested in this study, linking this global regulator to acquisition of these important substances required for growth.

MATERIALS AND METHODS

Strains and media. The strains, plasmids, and bacteriophages used in this study are shown in Table 1. Luria-Bertani (LB), and M9 minimal media and antibiotics were prepared as described previously (37, 58).

Construction of CWZ479. Phage λ 354 was retrieved from DL1504 by UV induction as described previously (58). The phage lysate was used to infect H1941 (MC4100 *fur*), and lysogens were selected on LB medium containing kanamycin to create CWZ479 (Table 1) as described previously (72).

Bacterial growth conditions. Bacterial cultures were inoculated and grown as described previously (71, 74). For iron-replete growth conditions at 37°C and 23°C, the bacteria were cultured in M9 glycerol medium (M9 minimal liquid medium containing 2.45 μ M ferric citrate, 30 μ M thiamine, 100 μ M calcium chloride, 1 mM magnesium sulfate, and 0.2% glycerol as a carbon source, pH 7) with aeration. For iron-depleted growth conditions, the bacteria were cultured in M9 glycerol medium containing 200 μ M 2,2'-dipyridyl, an iron-chelating agent. An inoculum from a single colony grown at 37°C was used to initiate parallel, colony-matched cultures grown at 37°C and 23°C. The cells from these cultures were harvested at equivalent optical densities (OD) after approximately 9 to 11 generations of growth. With the exception of stationary-phase cultures, which were harvested at an OD at 600 nm (OD₆₀₀) of 1.4 to 1.6, all cultures were harvested in early to mid-exponential phase (OD₆₀₀ = 0.2 to 0.6). The cell pellets were subsequently frozen in liquid nitrogen and stored at -80°C for RNA isolation.

RNA isolation. For microarray analyses, RNA was isolated by phenol-chloroform extraction as described previously (70) with the modifications that the RNA was subjected to a second DNase digestion in solution and cleaned using an RNeasy column according to the manufacturer's directions (QIAGEN). For quantitative real-time reverse transcription-PCR (qRT-PCR) experiments, RNA isolation was carried out as directed in the QIAGEN RNeasy Mini protocol with

the modifications that cell pellets were resuspended in 100 μ l Tris-EDTA containing lysozyme (400 μ g/ml) and boiled for 90 seconds prior to application to the column, and the RNA samples were digested twice with DNase (first on column and then in solution). RNA concentrations and purity were determined by spectrophotometry. Isolated RNAs were stored at -80°C until they were used.

cDNA synthesis, labeling, and hybridization. Synthesis, labeling of cDNA with Cy3/Cy5, and hybridization were performed using the 3DNA Array 350 RP Expression Array Detection Kit according to the manufacturer's protocol (Genisphere). cDNA for each condition was prepared from 2 μ g total RNA. cDNAs from both temperatures (37°C and 23°C) were cohybridized to slides containing full-length PCR products from all 4,290 annotated open reading frames (ORFs) in *E. coli* MG1655. The slides were produced by the University of Wisconsin-Gene Expression Center (<http://www.biotech.wisc.edu/GEC/>) and obtained at a reduced cost through the Genome Consortium for Active Teaching (<http://www.bio.davidson.edu/projects/gcat/gcat.html>). The hybridization solution (10 μ l of cDNA mixture, 1 μ l of denatured rat Hybloc DNA [1 μ g/ μ l], 20 μ l of 2 \times sodium dodecyl sulfate [SDS] buffer, 2 μ l of dT blocker, and 7 μ l of nuclease-free water) was placed over a prewarmed microarray slide and covered with a Corning 22- by 40-mm coverslip. The slide was then sealed in a hybridization chamber and submerged for 3 days in a 60°C water bath. Microarray slides were washed for 10 min in 2 \times SSC (1 \times SSC is 0.15 M NaCl plus 0.015 M sodium citrate)-0.2% SDS at 37°C, 10 min in 2 \times SSC at room temperature, and 10 min in 0.2 \times SSC at room temperature. The slides were dried by centrifugation at 10,000 rpm for 2 minutes. For the second hybridization to allow the 3DNA capture reagent to bind to the complementary Cy3- or Cy5-specific 3DNA capture sequences ligated to the cDNAs, each microarray received 42 μ l of hybridization solution (2.5 μ l of Cy3 capture reagent, 2.5 μ l of Cy5 capture reagent, 21 μ l of 2 \times SDS buffer, 0.5 μ l antifade reagent, and 15.5 μ l of nuclease-free water) and was hybridized as before via submersion for 3 to 4 h in a 61°C water bath. The slides were washed, dried, and kept in the dark until they were scanned and analyzed.

Microarray data analysis. Five slides were used in the analysis with cDNAs, representing three independent growth experiments and two technical replicates. Hybridized slides were scanned using GenePix Pro 4.1 software (Axon Instruments, Inc.). Photomultiplier tube voltage was altered for the initial scanning to ensure proper channel balance and decrease background. Any gene feature that had a signal-to-background ratio of <2.0 was rejected and not further analyzed. The ratios of median values were used for further analysis. The five slides used for detailed studies were normalized by a Lowess fit using GPROCESSOR, developed by Zhong Guan (<http://bioinformatics.med.yale.edu>). Replicate spots on the arrays were merged using GEPAS (<http://gepas.bioinfo.cipf.es/>). Significance analysis was completed as a one-class response using significance analysis of microarrays (65) with a delta of 0.65 and a median false-discovery rate of 1%. An ORF was considered temperature regulated if it demonstrated a statistically significant change in expression greater than 1.7-fold.

qRT-PCR. Primers and dual-labeled fluorogenic probes for qRT-PCR were created using Primer Express (v.1.5) software from Applied Biosystems. One-step reactions were prepared as directed by the manufacturer (QIAGEN QuantiTect kit or Invitrogen SuperScript III Platinum SYBR Green kit) using 50 ng of total RNA per reaction. An Applied Biosystems 7700 instrument was used for thermocycling with the following conditions: 50°C for 30 min, 95°C for 15 min, 40 cycles of 94°C for 15 seconds, and 60°C for 1 min. All reactions were performed in triplicate, with no reverse transcriptase controls run for each RNA sample to detect DNA contamination.

All reactions were normalized by using the same amount of total RNA (50 ng) in each reaction. Relative levels of gene expression were calculated as previously described (32) using the following equation: $\Delta C_T = C_T$ gene at $x^\circ\text{C}$, X genotype - C_T gene at 37°C M9 glyc, wild type and then transformed to relative changes (n -fold) using $2^{-\Delta\Delta C_T}$. C_T is the cycle number at which the real-time amplification curve crosses the user-defined threshold, $x^\circ\text{C}$ is the temperature at which the RNA was isolated (37°C or 23°C), and X is the genotype of the strain (wild type or mutant). Differences in average C_T values were determined to be statistically significant ($P < 0.05$) by two-way analysis of variance using STATA SE software (StataCorp). The data represent the average change (n -fold) determined from at least three independent experiments.

RESULTS

Microarray design to identify temperature-regulated genes in *E. coli* K-12 MC4100. Because many virulence genes in *E. coli* and other pathogens are regulated by temperature, we wanted to determine on a genomewide scale which genes are

TABLE 2. Genes demonstrating increased expression at 37°C

Gene ^a	Blattner no.	Product ^b	Change (<i>n</i> -fold) at 37°/23°C ^c
Amino acid transport and metabolism			
<i>argD</i>	b3359	Acetylornithine delta-aminotransferase	2.4
<i>aroF</i>	b2601	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase (DAHP synthetase), tyrosine-repressible	2.2
<i>aroG</i>	b0754	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase (DAHP synthetase)	2.0
<i>asnA</i>	b3744	Asparagine synthetase A	2.4
<i>carB</i>	b0033	Carbamoyl phosphate synthase, large subunit	2.8
<i>cysD</i>	b2752	ATP-sulfurylase, subunit 2 (ATP-sulfate adenyltransferase)	2.2
<i>cysH</i>	b2762	3'-Phosphoadenosine 5'-phosphosulfate (PAPS) reductase	2.6
<i>cysM</i>	b2421	Cysteine synthase B (<i>O</i> -acetylserine sulfhydrylase B)	2.4
<i>gcvP</i>	b2903	Glycine cleavage complex protein P, glycine decarboxylase, PLP dependent	1.9
<i>hisC</i>	b2021	Histidinol phosphate aminotransferase	1.9
<i>hisJ</i>	b2309	Histidine transport protein (ABC superfamily, <i>peri_bind</i>)	2.6
<i>hisQ</i>	b2308	Histidine and lysine/arginine/ornithine transport system (ABC superfamily, membrane)	2.2
<i>idnD</i>	b4267	L-Idonate 5-dehydrogenase, NAD binding	1.8
<i>livG</i>	b3455	ATP-binding component of high-affinity branched-chain amino acid transport system	1.9
<i>livK</i>	b3458	High-affinity branched-chain amino acid transport protein (ABC superfamily, <i>peri_bind</i>)	2.4
<i>lysC</i>	b4024	Aspartokinase III, lysine sensitive	2.6
<i>lysP</i>	b2156	Lysine-specific permease	2.5
<i>nanA</i>	b3225	<i>N</i> -Acetylneuraminate lyase (aldolase)	2.1
<i>tesA</i>	b0494	Acyl-coenzyme A thioesterase I; also functions as protease I	2.0
<i>trpD</i>	b1263	Anthranilate synthase component II	2.1
<i>ybiK</i>	b0828	L-asparaginase	1.9
<i>yecC</i>	b1917	Putative ATP-binding component of a transport system	2.5
<i>yecS</i>	b1918	Putative amino acid transport protein (ABC superfamily, membrane)	2.1
Carbohydrate transport and metabolism			
<i>araA</i>	b0062	L-arabinose isomerase	1.7
<i>galK</i>	b0757	Galactokinase	3.3
<i>garD</i>	b3128	(D)-galactarate dehydrogenase	3.0
<i>garL</i>	b3126	Alpha-dehydro-beta-deoxy-D-glucarate aldolase	5.4
<i>garP</i>	b3127	Putative transport protein	6.3
<i>lamB</i>	b4036	Maltoporin, high-affinity receptor for maltose and maltoseoligosaccharides; phage lambda receptor	2.3
<i>mglA</i>	b2149	ATP-binding component of methyl-galactoside transport and galactose taxis	3.3
<i>mglB</i>	b2150	Galactose transport protein (ABC superfamily, <i>peri_bind</i>)	3.1
<i>mglC</i>	b2148	Methyl-galactoside transport and galactose taxis	4.4
<i>nagE</i>	b0679	PTS family enzyme IIC (N terminal); enzyme IIB (center); enzyme IIC (C terminal)	2.4
<i>sgbH</i>	b3581	3-Keto-L-gulonate 6-phosphate decarboxylase	1.8
<i>srlB</i>	b2704	PTS family enzyme IIA, glucitol/sorbitol specific	2.2
<i>uidA</i>	b1617	Beta-D-glucuronidase	2.3
<i>uxuB</i>	b4323	D-mannonate oxidoreductase	2.1
<i>yeiC</i>	b2166	Putative kinase	2.4
Cell motility and secretion			
<i>fliQ</i>	b1949	Flagellar biosynthesis	1.9
Cell wall/membrane biogenesis			
<i>nmpC</i>	b0553	DLP12 prophage; outer membrane porin, at locus of <i>qsr</i> prophage	10.0
<i>ompT</i>	b0565	DLP12 prophage; protease VII, outer membrane protein 3b (a), putative porin	3.3
<i>yciD</i>	b1256	Outer membrane protein W; colicin S4 receptor; putative transport protein	3.0
Coenzyme transport and metabolism			
<i>btuB</i>	b3966	Outer membrane porin: vitamin B ₁₂ /cobalamin transport, receptor for E colicins,	2.1
<i>entC</i>	b0593	Isochorismate hydroxymutase 2, enterochelin biosynthesis	2.0
<i>folE</i>	b2153	GTP cyclohydrolase I	2.7
<i>ubiA</i>	b4040	4-Hydroxybenzoate-octaprenyltransferase	2.1
Defense mechanisms			
<i>hsdR</i>	b4350	Endonuclease R, host restriction	2.8
Energy production and conversion			
<i>fumB</i>	b4122	Fumarase B (fumarate hydratase class I), anaerobic isozyme	1.9
<i>galT</i>	b0758	Galactose-1-phosphate uridylyltransferase	2.6
<i>hybA</i>	b2996	Hydrogenase-2 small subunit	2.1
<i>napF</i>	b2208	Fe-S ferredoxin-type protein, electron transfer	1.7
<i>nirB</i>	b3365	Nitrite reductase [NAD(P)H] subunit	3.3
<i>sucD</i>	b0729	Succinyl-coenzyme A synthetase, alpha subunit	1.9

Continued on following page

TABLE 2—Continued

Gene ^a	Blattner no.	Product ^b	Change (<i>n</i> -fold) at 37°/23°C ^c
<i>ybiC</i>	b0801	Putative dehydrogenase	3.1
<i>yqhD</i>	b3011	Putative alcohol dehydrogenase	3.3
Inorganic-ion transport and metabolism			
<i>cirA</i>	b2155	Outer membrane pore protein, receptor for colicin I, requires TonB	2.8
<i>cysA</i>	b2422	ATP-binding component of sulfate permease A protein; chromate resistance	2.3
<i>cysC</i>	b2750	Adenosine 5'-phosphosulfate kinase	2.7
<i>cysI</i>	b2763	Sulfite reductase, alpha subunit	2.2
<i>cysJ</i>	b2764	Sulfite reductase (NADPH), flavoprotein beta subunit	2.7
<i>cysN</i>	b2751	ATP-sulfurylase (ATP:sulfate adenyltransferase), subunit 1, probably a GTPase	2.1
<i>cysP</i>	b2425	Thiosulfate binding protein	3.1
<i>fecA</i>	b4291	KpLE2 phage-like element; outer membrane porin, receptor for ferric citrate	2.5
<i>fecR</i>	b4292	KpLE2 phage-like element; regulator in multicomponent regulatory system with FecI/FecA	3.3
<i>fepA</i>	b0584	Outer membrane porin, receptor for ferric enterobactin and colicins B and D	2.0
<i>fepB</i>	b0592	Ferric enterobactin (enterochelin) binding protein; periplasmic component	1.9
<i>fepC</i>	b0588	ATP-binding component of ferric enterobactin transport	2.0
<i>fepG</i>	b0589	Ferric enterobactin transport protein	2.0
<i>fes</i>	b0585	Enterochelin esterase	2.7
<i>yedO</i>	b1018	Hypothetical protein	1.8
Lipid transport and metabolism			
<i>acs</i>	b4069	Acetyl-coenzyme A synthetase	2.4
<i>fadE</i>	b0221	Medium-long-chain fatty acyl-coenzyme A dehydrogenase	1.8
<i>garR</i>	b3125	Tartronate semialdehyde reductase (TSAR)	1.8
Nucleotide transport and metabolism			
<i>allB</i>	b0512	Allantoinase	1.9
<i>rihA</i>	b0651	Pyrimidine-specific nucleoside hydrolase	2.0
<i>rihC</i>	b0030	Nucleoside hydrolase	3.3
Posttranslational modification, protein turnover, chaperones			
<i>ccmC</i>	b2199	Heme export protein (ABC superfamily, membrane)	2.3
<i>ccmG</i>	b2195	Heme lyase/disulfide oxidoreductase (thiol-disulfide interchange protein)	1.8
<i>cysU</i>	b2424	Sulfate, thiosulfate transport system permease T protein	2.7
<i>yeeD</i>	b2012	Hypothetical protein	3.0
Replication, recombination, and repair			
<i>nohB</i>	b0560	DLP12 prophage; bacteriophage DNA- packaging protein	2.0
<i>sbcB</i>	b2011	Exonuclease I, 3'→'-specific; deoxyribophosphodiesterase	2.0
<i>trs5_1</i>	b0259	CP4-6 prophage; IS5 protein 1	2.0
<i>trs5_10</i>	b3218	IS5 protein 10	2.3
<i>trs5_11</i>	b3505	IS5 protein 11	2.2
<i>trs5_3</i>	b0656	IS5 protein	2.2
<i>trs5_4</i>	b1331	IS5 protein	2.1
<i>trs5_5</i>	b1370	IS5Y transposase	2.4
<i>trs5_6</i>	b1994	IS5 protein	2.4
<i>trs5_7</i>	b2030	IS5 protein	2.6
<i>trs5_8</i>	b2192	IS5 protein	2.5
<i>trs5_9</i>	b2982	IS5 protein	3.2
Secondary-metabolite biosynthesis, transport and catabolism			
<i>entB</i>	b0595	2,3-Dihydro-2,3-dihydroxybenzoate synthetase, isochroismatase	2.1
<i>entE</i>	b0594	2,3-Dihydroxybenzoate-AMP ligase	2.2
<i>idnO</i>	b4266	5-Keto-D-gluconate 5-reductase	2.0
<i>srlD</i>	b2705	Glucitol (sorbitol)-6-phosphate dehydrogenase	1.9
<i>yeiN</i>	b2165	Hypothetical protein	3.1
Signal transduction mechanisms			
<i>yedT</i>	b1025	Putative transmembrane protein	1.7
Transcription			
<i>fecI</i>	b4293	KpLE2 phage-like element; sigma 19 factor of RNA polymerase	2.2
<i>lrhA</i>	b2289	NADH dehydrogenase transcriptional regulator, LysR family; modulates SprE (RssB) activity	4.4
<i>yijC</i>	b3963	Putative regulator (TetR/AcrR family)	2.4
General function prediction or function unknown			
<i>yadS</i>	b0157	Putative membrane protein	2.2

Continued on following page

TABLE 2—Continued

Gene ^a	Blattner no.	Product ^b	Change (<i>n</i> -fold) at 37°/23°C ^c
<i>yddA</i>	b1496	Putative ABC transport system ATP-binding protein	4.8
<i>yeeE</i>	b2013	Putative membrane component of transport system	2.2
<i>yfiD</i>	b2579	Putative formate acetyltransferase	4.4
Not in COGS database			
<i>b1172</i>	b1172	Hypothetical protein	2.0
<i>b2191</i>	b2191	Unknown CDS	2.3
<i>cdaR</i>	b0162	Regulator of D-galactarate, D-glucarate, and D-glycerate metabolism	1.9
<i>cysW</i>	b2423	Sulfate transport system permease W protein	2.6
<i>malM</i>	b4037	Periplasmic protein of <i>mal</i> regulon	2.4
<i>pppA</i>	b2972	Bifunctional prepilin peptidase	2.7
<i>srlA</i>	b2702	PTS family enzyme IIC, glucitol/sorbitol specific	2.0
<i>srlE</i>	b2703	PTS family enzyme IIBC, glucitol/sorbitol specific	2.5
<i>ybcT</i>	b0556	DLP12 prophage; bacteriophage lambda endopeptidase homolog	3.5
<i>ybcU</i>	b0557	DLP12 prophage; bacteriophage lambda Bor lipoprotein homolog, involved in serum resistance	3.5
<i>yddB</i>	b1495	Hypothetical protein	8.0
<i>yedO</i>	b1919	D-cysteine desulfhydrase, PLP-dependent enzyme	2.4
<i>ygaW</i>	b2670	Putative membrane protein	4.9
<i>yhcE</i>	b3217	Unknown CDS	2.1
<i>yijD</i>	b3964	Putative membrane protein	3.4
<i>yjiD</i>	b4326	Unknown CDS	2.4
<i>ylaC</i>	b0458	Putative membrane protein	2.1
<i>ymfL</i>	b1147	e14 prophage; putative negative regulator	2.3

^a Genes involved in iron utilization and/or known to be Fe²⁺-Fur regulated are shown in boldface.

^b Product descriptions and functional categories are based upon the *E. coli* K-12 COGS categorization (63). Descriptions were shortened in some instances.

^c Change is indicated as the average ratio of medians (37°C/23°C). All genes included showed a statistically significant increase at 37°C as described in Materials and Methods and based on the results from five microarrays.

modulated in adapting to both human host temperature (37°C) and ambient room temperature (23°C). Growth and medium conditions were chosen (M9 glycerol medium with aeration) to maximize *papBA* transcription at 37°C so we could use this operon as a temperature-regulated positive control in our microarrays. Parallel cultures grown at 37°C and 23°C were harvested in exponential phase at 9 to 11 generations of growth after inoculation. Thus, the results presented here reflect the adapted state and signify genes whose expression is differentially maintained over long-term growth at a given temperature.

Overall, 126 genes were found to be more highly expressed at 37°C (Table 2), whereas 297 genes were more highly expressed at 23°C (C. A. White-Ziegler, S. Um, N. M. Pérez, A. L. Berns, A. J. Malhowski, and S. Young, unpublished data). The *lacZ*, *lacY*, and *lacA* genes, contained on a chromosomally located *papBA* transcriptional fusion and driven by the pBA promoter, showed the greatest increase at 37°C (13.7-fold, 7.5-fold, and 5.4-fold, respectively), supporting the validity of the experimental design.

COGS categorization of genes. The 126 genes with increased expression at 37°C were categorized using the Clusters of Orthologous Groups of Proteins (COGS) database (63). At least one gene was found in each category, with the exception of cell cycle control (Table 2 and Fig. 1). Twelve genes within the category of replication, recombination, and repair were increased at 37°C; with the exception of *sbcB*, all encode proteins with phage/transposon-related functions (Table 2 and Fig. 1). Approximately 12% of the genes with increased transcription at 37°C have unknown functions (Table 2 and Fig. 1). The other three categories of genes that dominated the response

were those associated with amino acid (19%), carbohydrate (12%), and inorganic-ion (12%) transport and metabolism.

Human body temperature increases carbohydrate and amino acid uptake and utilization gene expression. Expression levels of 15 genes involved in carbon transport and utilization and 23 genes involved in synthesis and transport of amino acids were significantly increased at 37°C in comparison to 23°C (Table 2). These genes showed increased expression in the absence of the cognate sugar and, in the case of amino acid transporters, in the absence of the specific amino acid. The carbohydrate utilization genes that were increased at 37°C are involved in the catabolism of a diversity of sugars, including arabinose, maltose, galactose, D-galactarate, D-glucarate, D-gluconuride, glucitol, and sorbitol. A variety of amino acid transporters demonstrated increased expression at 37°C, whereas biosynthesis was focused on histidine and cysteine (Table 2). Genes required for the uptake and utilization of other important compounds for growth, including sulfur/thio-sulfate (*cysDHM*) and cobalamin (*btuB*), were similarly increased at 37°C.

Human body temperature increases expression of genes for iron uptake and utilization. Interestingly, 12 genes involved in iron utilization were identified, representing 10% of all genes with increased expression at 37°C (Table 2). They represent ferric enterobactin uptake genes (*fepA*, *fepB*, *fepC*, and *fepG*), enterobactin synthesis genes (*entB*, *entC*, and *entE*), and the enterobactin esterase gene *fes*. In addition, the genes *fecA*, *fecI*, *fecR*, and *cirA*, involved in utilization of other iron forms (ferric-dicitrate, ferrous iron, ferri-copragen/rhodotorulic acid, ferrioxamine B, and ferric-dihydrobenzoate), were increased in expression at 37°C compared to 23°C.




FIG. 1. Percentages of ORFs demonstrating increased expression at 37°C compared to 23°C in each of 20 functional categories. ORFs were placed in functional categories according to the *E. coli* K-12 COGS categorization (63).

Based on other studies, it is known that transcription of all of these genes is responsive to the iron concentration, showing high expression under iron-depleted conditions but repressed by the transcriptional regulator Fur (ferric uptake regulator) when iron is present (reviewed in references 16 and 25). In our microarrays, the bacteria were grown in iron-replete medium, where Fur binds iron and act as a repressor. Thus, these results indicate that the increased expression of these genes is due to the effect of temperature and not to limiting iron conditions.

Several additional genes that are regulated by Fe²⁺-Fur demonstrated higher expression at 37°C than at 23°C (35). Due to their similarity to ABC transporters (YddA) and TonB-dependent outer membrane receptors (YddB), it has been hypothesized that these two proteins may form part of a new iron uptake system (35). Interestingly, the genes were also identified in a transposon mutagenesis screen and found to be required for optimal growth at 37°C in rich medium (57). They are two of the most highly expressed genes at 37°C compared to 23°C (4.8- and 8.0-fold, respectively). *yciD* (*ompW*) encodes a S4 colicin receptor and putative transport protein (48), similar to the iron acquisition genes *cirA* and *fepA*, which also serve as colicin receptors. *garPLR*, *napF*, *nirB*, and *ycdO* are also included in Fe²⁺-Fur-controlled operons (35) and demonstrated temperature-regulated expression in this study.

Thermoregulation of amino acid, carbohydrate, and iron utilization genes occurs in both exponential and stationary phases. We analyzed the expression of six genes, two from each

of the amino acid, carbohydrate, and iron utilization gene categories, at 37°C and 23°C under a variety of growth conditions. Both *cysP* and *hisJ* encode periplasmic binding proteins that are involved in thiosulfate (59) and histidine (31) transport, respectively. For carbohydrate utilization genes, we examined *srlE*, a subunit of the glucitol/sorbitol PTS permease (49), and *garP*, a putative glucarate transporter (46). For iron utilization genes, we studied *fes* and *cirA*. *Fes* is a cytoplasm protein that functions in releasing iron bound to the bacterial iron chelator enterobactin (6). *CirA* is an outer membrane receptor with broad specificity that has been postulated to transport iron complexes of enterobactin and is known to transport colicins I/V and catechol-substituted cephalosporins (42). We also analyzed the expression of *papB*, a known temperature-regulated virulence gene. *papB* is the first gene in the multicistronic *papBA* operon that encodes pyelonephritis-associated pili (Pap) (reviewed in reference 28).

In concordance with our microarray results, all six genes demonstrated statistically significant thermoregulatory responses in exponentially growing cells in M9 glycerol medium (Fig. 2). The amino acid utilization genes *hisJ* and *cysP* demonstrated the smallest differential in expression based on temperature, with 1.9- and 3.1-fold-higher expression, respectively, at 37°C compared to 23°C. *garP* and *srlE* demonstrated higher differentials in expression due to temperature, with 7.2- and 3.4-fold increases, respectively, in their transcription at 37°C compared to 23°C. The relative expression levels of *fes* and *cirA* were




FIG. 2. Effects of growth phase and medium on the thermoregulation of gene expression for amino acid, carbohydrate, and iron utilization genes. The bars indicate relative levels of expression measured

3.4-fold and 3.5-fold greater, respectively, at 37°C than at 23°C. *papB* demonstrated a 96.7-fold-greater level of expression at 37°C than at 23°C, similar to previous results using a β-galactosidase assay, where transcription from the *papBA* promoter was 52-fold higher at 37°C than at 23°C (70). Together, our data support a model in which temperature serves as an important environmental cue that modulates gene expression.

All cultures grown to stationary phase in M9 glycerol medium still retained a thermoregulatory response, indicating that temperature serves to maintain higher levels of expression at 37°C independent of the growth phase (Fig. 2). For the iron utilization genes (*fes* and *cirA*) and *papB*, transcription at 37°C was increased significantly in stationary phase, leading to a higher differential in expression between 37°C and 23°C (13.4-, 9.3-, and 994-fold, respectively). For the carbohydrate utilization genes (*garP* and *srlE*) and *hisJ*, expression was decreased at both 37°C and 23°C in stationary phase, but a statistically significant thermoregulatory response was retained (3.7-, 4.9-, and 7.7-fold higher at 37°C, respectively). Growth in stationary phase decreased *cysP* expression at 37°C but did not alter transcription significantly at 23°C, leading to a smaller differential in gene expression based on temperature during this growth phase.

Glucose as a carbon source abrogates thermoregulation for the carbohydrate genes, but not the amino acid or iron utilization genes. Growth in M9 medium utilizing glucose as a carbon source abrogated the thermoregulatory response for the carbohydrate utilization genes (*garP* and *srlE*) so that they had equivalent expression at both 37°C and 23°C (Fig. 2). All of the other genes retained a statistically significant differential in expression based on temperature when grown in M9 glucose medium (Fig. 2). Similar to stationary phase, growth on glucose increased expression of the iron utilization gene *cirA* at 37°C, resulting in a 6.5-fold differential in expression between the two temperatures, whereas glucose increased *fes* expression at both 37°C and 23°C so that the differential in their expression was only 1.9-fold. Growth in M9 glucose decreased the expression of *hisJ* and *cysP* at both temperatures in comparison to glycerol as a carbon source and resulted in a greater differential in expression based on temperature (4.1-fold and 6.0-fold, respectively). *papB* demonstrated greatly reduced levels of transcription at both temperatures so that the differential in expression between 37°C and 23°C was only 12.4-fold compared to the 97-fold observed in M9 glycerol.

Growth in LB medium had varied effects on the thermoregulatory response. Growth in LB medium greatly increased expression of *cysP* so that equivalent expression levels of these genes were observed at both 37°C and 23°C, abrogating the thermoregulatory response (Fig. 2). For *papB* and *hisJ*, LB

in the wild-type strain DL1504 in M9 glycerol medium in exponential phase (M9 glyc), M9 glycerol medium in stationary phase (M9 glyc stat), M9 glucose medium in exponential phase (M9 gluc), and LB medium in exponential phase (LB). Relative levels of expression are expressed in comparison to the wild-type strain grown in M9 glycerol medium at 37°C. A bracket indicates a statistically significant difference in expression levels based on temperature under the given condition and is accompanied by the ratio of expression (37°C/23°C). Error is expressed as ±1 standard deviation from the mean.




FIG. 3. Effects of temperature and iron concentration on *cirA*, *fes*, and *papBA* transcription. The bars indicate relative levels of expression measured in the wild-type strain DL1504 under iron-replete (+ Fe) and iron-depleted (– Fe) conditions. Relative levels of expression are expressed in comparison to the wild-type strain grown in iron-replete (M9 glycerol) medium at 37°C. A bracket indicates a statistically significant difference in expression levels based on temperature under the given condition and is accompanied by the ratio of expression (37°C/23°C). Error is expressed as ± 1 standard deviation from the mean.

decreased expression at both temperatures, but statistically significant temperature differentials (4.8 and 2.6, respectively) were retained (Fig. 2). Interestingly, for the iron utilization genes *cirA* and *fes*, LB significantly decreased the expression of both genes at 37°C but had little effect on expression at 23°C, so that *fes* and *cirA* were more highly expressed at 23°C than at 37°C under these growth conditions (Fig. 2). For *srlE*, LB increased expression at 37°C, leading to a 13.3-fold differential between its expression at 37°C and at 23°C. *garP* expression increased at both 37°C and 23°C, but there was significant colony variability, so that an accurate relative expression could not be determined (data not shown).

Low-iron conditions abrogate the thermoregulatory effect on *cirA* and *fes*, but not *pap*, transcription. Because iron acquisition is essential to bacterial growth and impacts uropathogenic *E. coli* virulence (30, 54, 64), we further assessed the relative importance of iron availability and temperature as cues for regulating transcription of the iron utilization genes *fes* and *cirA* by measuring expression under iron-depleted conditions at both 37°C and 23°C in the wild-type strain (Fig. 3). Iron-depleted conditions were obtained by adding 200 μ M 2,2'-

dipyridyl, an iron-chelating agent, to M9 glycerol medium for growth. *cirA* and *fes* transcription levels were higher under iron-depleted conditions than under iron-replete conditions at both temperatures. *fes* demonstrated 3.3- and 3.0-fold increases at 37°C and 23°C, respectively, under iron-depleted conditions compared to iron-replete conditions at 37°C (Fig. 3). Similarly, *cirA* transcription demonstrated 1.6- and 2.5-fold increases at 37°C and 23°C, respectively, in iron-depleted medium compared to iron-replete medium at 37°C (Fig. 3). These results demonstrate that a lack of iron has a stimulatory effect on the transcription of these genes at both temperatures and that the thermoregulatory effect on *cirA* and *fes* transcription is abrogated when iron is limiting.

Because research has indicated that the urinary tract is an iron-limited environment (61) and that iron affects the expression of certain fimbrial operons (27), we investigated whether iron availability, in addition to temperature, might serve as a cue for controlling *papBA* gene expression. Our data demonstrate that expression levels for *papBA* are similar under iron-depleted conditions to those seen in iron-replete media at both temperatures and that thermoregulation of *papBA* transcription still occurs when iron is limiting (Fig. 3). Thus, while *fes*, *cirA*, and *papBA* exhibited thermoregulated expression under iron-replete conditions, they showed much different expression patterns under iron-limiting conditions, where *fes* and *cirA* are no longer affected by temperature but *papBA* remains thermoregulated.

The thermoregulator H-NS, but not RimJ, alters *cirA* and *fes* transcription. We wanted to assess whether temperature regulation of *cirA* and *fes* under iron-replete conditions was mediated by thermoregulators known to control *papBA* transcription, thereby suggesting a coordinate thermoregulation of fimbrial expression with iron acquisition. Thermoregulation of *cirA* and *fes* was investigated by analyzing the relative levels of gene expression at 37°C and 23°C in the *hns651* strain. The *hns651* mutation contains an insertion sequence that leads to the loss of H-NS expression (70). At 37°C, both *fes* and *cirA* expression levels were decreased an average of 3.3- and 4.9-fold, respectively, in the *hns651* strain compared to the wild type at 37°C (Fig. 4). This decrease in expression in the mutant, in the absence of a temperature change, indicates that H-NS has a positive effect on *cirA* and *fes* transcription at 37°C. At 23°C, relative levels of *cirA* and *fes* in the *hns651* mutant were similar to that measured in the wild-type strain at 23°C (Fig. 4). In the *hns651* mutant, there was no longer a significant difference in the levels of *cirA* and *fes* transcription based on temperature, primarily due to decreased expression at 37°C rather than an alleviation of a repressive effect of H-NS at 23°C.

RimJ is an N-terminal acetyltransferase that represses *papBA* transcription (71–73), and thermoregulatory repression was relieved in the $\Delta rimJ$ mutant so that *papBA* transcription levels were equivalent at 37°C and 23°C (71). In a $\Delta rimJ$ strain, there was an average 3.2- and 3.3-fold difference in the levels of *fes* and *cirA* expression, respectively, at 37°C compared to 23°C (Fig. 4). These results are similar to those observed in the wild-type strain, indicating that *cirA* and *fes* are not under the thermoregulatory control of RimJ.

Thermoregulation of *cirA* and *fes* transcription occurs independently of Fur. Although there was no previous research suggesting a thermoregulatory role for Fur, we investigated




FIG. 4. Effects of the $\Delta rimJ$, *hns651*, and *fur* mutations on *cirA* and *fes* transcription. The bars indicate relative levels of expression measured in the wild-type strain DL1504 (wt), in the $\Delta rimJ$ mutant strain CWZ388 ($\Delta rimJ$), in the *hns651* mutant strain DL1947 (*hns651*), and in the *fur* mutant strain CWZ479 (Δfur). Relative levels of expression are expressed in comparison to the wild-type strain grown in M9 glycerol medium at 37°C. A bracket indicates a statistically significant difference in expression levels based on temperature under the given condition and is accompanied by the ratio of expression (37°C/23°C). Error is expressed as ± 1 standard deviation from the mean.

whether Fur might have a dual regulatory role in sensing both iron concentration and temperature to control *fes* and *cirA* transcription. As expected, introduction of the *fur* mutation caused an increase in transcription of *fes* and *cirA* at both 37°C and 23°C in the absence of this transcriptional repressor (Fig. 4). However, unlike the experiment in which iron was limiting and *cirA* and *fes* transcription was no longer thermoregulated (Fig. 3), analysis of *fes* and *cirA* transcription in the *fur* mutant strain showed a thermoregulatory trend similar to that seen in the wild-type strain (Fig. 4). *fes* and *cirA* showed 2.3- and 1.9-fold differences, respectively, in expression levels at 37°C compared to 23°C in the *fur* mutant (Fig. 4). Thus, our results indicate that thermoregulation of *fes* and *cirA* occurs in the absence of Fur.

H-NS controls the regulation of carbohydrate and amino acid utilization genes. Because of the role of H-NS in controlling iron utilization gene transcription, we wanted to explore if H-NS played a similar regulatory role for the amino acid and carbohydrate utilization genes in this study. Similar to the iron utilization genes, a *hns651* mutation led to a 3.4-fold decrease in *srhE* expression at 37°C compared to the wild-type strain at 37°C, thereby abrogating any differential in expression due to temperature and indicating a positive role for H-NS in the regulation of *srhE* (Fig. 5). For *hisJ*, there were statistically significant decreases in transcription at both 37°C and 23°C by the *hns651* mutation (2.3- and 5.1-fold, respectively), indicating that H-NS has a positive role in controlling transcription of this gene at both temperatures (Fig. 5). H-NS plays a predominantly repressive role for transcription of *cysP* at low temperature, with the *hns651* mutation leading to statistically significant increased transcription at 23°C and a subsequent loss of a thermoregulatory differential in gene expression. *garP* expression demonstrated a purely negative effect of H-NS on tran-




FIG. 5. Effects of the *hns651* mutation on *hisJ*, *cysP*, *srhE*, and *garP* transcription. The bars indicate relative levels of expression measured in the wild-type strain DL1504 (wt) and in the *hns651* mutant strain DL1947 (*hns651*). Relative levels of expression are expressed in comparison to the wild-type strain grown in M9 glycerol medium at 37°C. A bracket indicates a statistically significant difference in expression levels based on temperature under the given condition and is accompanied by the ratio of expression (37°C/23°C). Error is expressed as ± 1 standard deviation from the mean.

scription, as an *hns651* mutation led to increased expression of *garP* at both temperatures (Fig. 5). Together, our results indicate that H-NS may have a broad regulatory role in controlling gene expression that is responsive to temperature but that there are a variety of modes by which H-NS exerts this influence.

DISCUSSION

Our microarray data indicate that mammalian host temperature (37°C) serves to increase and maintain iron, carbohydrate, and amino acid utilization genes at a higher steady-state level of expression than at 23°C. While temperature has been widely studied as a factor controlling virulence gene expression, our study and others indicate that temperature is used genomewide as a cue for adapting gene expression that may facilitate colonization of a mammalian host (7, 8, 24, 38, 45, 60). In particular, our study shows commonality with *Y. pestis*, in which the expression levels of a significant proportion of genes for amino acid biosynthesis and carbohydrate utilization were increased at 37°C compared to growth at 26°C (24, 38), and with group A *Streptococcus*, in which temperature was identified as an important cue in controlling the transcription of iron utilization genes (60).

We propose that temperature may serve as a sentinel cue to quickly allow adaptation of the organism to its environment. Upon entry into a human host, temperature would increase and maintain higher levels of expression of the genes needed for growth. This would allow the bacterium to preemptively prepare for and utilize a diverse set of carbohydrate, amino acid, and iron sources that it might encounter as it transits through different niches in the host, allowing it to more efficiently draw upon the available resources. Interestingly, a recent study demonstrated that a number of iron, carbohydrate, and amino acid utilization genes were increased in expression in bacteria obtained from patients colonized with the asymp-

omatic-bacteriuria *E. coli* strain 83972 (53). Together with preliminary experiments from our laboratory demonstrating that a shift from 23°C to 37°C causes a quick increase in iron and fimbrial gene transcription (data not shown), these data support the model that temperature may be an important *in vivo* cue for modulating bacterial gene expression.

The role of temperature in regulating iron utilization genes is particularly interesting because of evidence that these genes are increased in expression within a mammalian host and that they are required for virulence *in vivo*. As mentioned previously, bacteria collected from patients colonized by the asymptomatic-bacteriuria *E. coli* strain 83972 demonstrated increased expression of genes within the *fep*, *ent*, *fhu*, and *iut* iron systems (53). Similarly, bladder infection in mice from uropathogenic *E. coli* led to increased expression of a number of iron acquisition genes (61), while in a mouse model of peritonitis, transcriptional fusions to iron acquisition genes were isolated (51). Torres et al. assessed the effect of iron acquisition on virulence, demonstrating that uropathogenic *E. coli* compromised in the ability to acquire iron through multiple pathways (*tonB* mutants) were attenuated and those deficient in a single pathway were less successful at colonization in mixed competition in mice (64). Other studies demonstrated urovirulence for individual iron acquisition systems and the presence of additional iron utilization systems in the genomes of pathogenic *E. coli* (10, 15, 30, 47, 55, 62, 69). Together, these studies provide *in vivo* evidence for the importance of iron acquisition for *E. coli* virulence.

Our study delineates the increased expression of several genes in *E. coli* K-12 in the enterobactin system (*ent*, *fep*, and *fes*) and the ferric citrate (*fec*) uptake systems at human body temperature, supporting the broad use of this cue in regulating multiple iron uptake pathways. This is further supported by our results indicating that the Fhu system (*fhuA*), involved in the uptake of ferrichrome and other iron forms, is also temperature regulated (data not shown). Uropathogenic and enterohemorrhagic *E. coli* strains are known to contain additional systems for iron uptake beyond those found in *E. coli* K-12 (47, 69), and it would be interesting to investigate if temperature regulation is broadly applicable to these iron utilization systems as well.

Our assessment of temperature regulation demonstrated that the majority of genes retained a higher level of expression at 37°C than at 23°C under a variety of growth conditions, including medium, growth phase, and carbon source, supporting the importance of temperature in modulating gene expression. At the same time, it should be noted that these conditions often influenced gene expression so that the level of expression at 37°C and/or 23°C was altered, leading to an expansion or contraction of the temperature differential, depending on the input of additional environmental cues and reflecting the ability of *E. coli* to integrate multiple environmental cues and adapt accordingly. Some clear exceptions to this generality were observed in which a given environmental cue led to an abrogation of the thermoregulatory response.

Regulation of *fes* and *cirA* by iron and temperature provide an example in which multiple cues are used for adapting gene expression. One could imagine that it could be advantageous to the bacterium to use both cues to adapt to various conditions within the host. In our *in vitro* experiments, limiting iron is the

overriding cue and has the ability to abrogate the thermoregulatory response, significantly increasing expression of *cirA* and *fes* regardless of temperature. This response would be what one would expect, given the requirement for iron for bacterial growth and the regulation of these genes by Fur. When might it be advantageous to have temperature regulation of iron acquisition genes? Either upon initial entry into the host or in niches where iron is available within the host, using the cue of host temperature to increase iron acquisition utilization gene expression would benefit the bacterium by allowing it to more efficiently capture and store iron that could subsequently be used when iron-limited niches are encountered.

Similar to the iron utilization genes in our study, others have noted the regulation of virulence gene expression by both temperature and iron availability. In *Pseudomonas aeruginosa*, the *tolRQ* and *tolA* genes, involved in pyocin transport, are optimally expressed at 37°C under iron-limiting conditions; low temperature and high iron concentrations decrease the expression of these genes (29). Similarly, Shiga toxin synthesis in *Shigella dysenteriae* is increased by 37°C temperature and low-iron conditions (68). In group A *Streptococcus*, temperature is an important cue in controlling the transcription of iron utilization genes and hemolysins (60). This coordinate regulation suggests a model in which a temperature of 37°C and iron-limiting conditions cause high expression of a number of genes required for virulence, conditions that most closely replicate the host environment.

Based on these observations and the evidence suggesting that the mammalian host is a 37°C, iron-limited environment (53, 61), we hypothesized that the *pap* fimbrial genes might be more highly expressed under iron-limiting conditions. This, however, was not the case, as our results showed that *papB* was transcribed at similar levels at 37°C when grown under iron-depleted conditions and iron-replete conditions (Fig. 3). Previous experiments demonstrated that high iron concentrations do not affect *papBA* transcription (74), and our results presented here do not support a role for iron or the Fur protein in controlling *papBA* gene expression (data not shown). Experiments growing uropathogenic *E. coli* *in vitro* in human urine at 37°C induced *pap* transcription (61), and previous studies identified the roles of several individual environmental stimuli in controlling *papBA* transcription (74), indicating that while iron may not play a role in *pap* transcription, there are likely multiple overlapping cues, including temperature, that control *pap* expression *in vivo*. It should be noted that our experiments were conducted in an *E. coli* K-12 strain, and future studies will address whether *papBA* transcription is modulated in the same manner in a uropathogenic *E. coli* strain.

Our results indicate that H-NS regulates iron, carbohydrate, and amino acid utilization gene expression in *E. coli*, supporting a broad role for this regulator in the adaptation to temperature. For *cirA* and *fes*, the *hns651* mutation causes decreased expression of these genes at 37°C, suggesting an activating role for H-NS in the transcription of the genes at host temperature. We also found this to be the case for *hisJ* and *srlE*, where H-NS has an activating role at one or both temperatures. This regulation is different from that of many H-NS controlled genes, where H-NS serves as a transcriptional silencer in response to a variety of environmental cues and deletion of H-NS leads to high-level expression under the

normally repressive cue (reviewed in references 1 and 13). Since there is no evidence that H-NS can act as a transcriptional activator, it has been postulated that H-NS may play an indirect role (reviewed in reference 13). For the *Erwinia chrysanthemi* pectate lyase synthesis genes (40, 41) and *E. coli* flagellar synthesis genes (2, 3), it has been found that H-NS exerts its control, not on these promoters directly, but at the promoters of transcriptional repressors that act upon these target genes. Similarly, it may be that H-NS disrupts the regulation of *cirA*, *fes*, *srlE*, and *hisJ* through indirect effects. It is interesting that the *hns651* mutation has a similar effect on the *papBA* operon, causing a 7.6-fold decrease in transcription at 37°C (67, 70, 74). Recent microarray results in our laboratory indicate that H-NS controls a majority of the temperature-regulated genes identified in this study (unpublished data), supporting a broad role of H-NS in adapting to host and ambient temperature conditions.

While temperature has been known to be a control for the expression of virulence genes, such as fimbriae, toxins, and adhesins (33), this study and others demonstrate that temperature may have a broader effect to fine tune and regulate a number of genes to presumably allow more efficient colonization of the host. Future studies to investigate and understand the pathways of this thermal regulation could yield valuable anti-infective targets for chemotherapeutic drugs that would decrease the ability of bacteria to compete and survive within the host.

ACKNOWLEDGMENTS

We thank the Genome Consortium for Active Teaching and the University of Wisconsin Gene Expression Center for making microarrays available at a reduced cost for use by the two undergraduates, S.Y. and A.M., who completed the microarray experiments in this study. We are grateful to present and former Smith College students, staff, and faculty for their technical assistance and advice, including Michelle Ploutz, Scott Edmands, Eva Ladow, Adam Hall, Nick Horton, and Lori Sanders.

This work was supported by National Institutes of Health grant GM62792 to C.W.-Z., by the Albert F. Blakeslee Trust, and by Smith College. A.M. and S.Y. were supported by student grants through the Albert F. Blakeslee Trust and by the Howard Hughes Medical Institute grant awarded to Smith College.

REFERENCES

1. **Atlung, T., and H. Ingmer.** 1997. H-NS: a modulator of environmentally regulated gene expression. *Mol. Microbiol.* **24**:7–17.
2. **Bertin, P., F. Hommais, E. Krin, O. Soutourina, C. Tendeng, S. Derzelle, and A. Danchin.** 2001. H-NS and H-NS-like proteins in Gram-negative bacteria and their multiple role in the regulation of bacterial metabolism. *Biochimie* **83**:235–241.
3. **Bertin, P., E. Terao, E. H. Lee, P. Lejeune, C. Colson, A. Danchin, and E. Collatz.** 1994. The H-NS protein is involved in the biogenesis of flagella in *Escherichia coli*. *J. Bacteriol.* **176**:5537–5540.
4. **Blyn, L. B., B. A. Braaten, C. A. White-Ziegler, D. H. Rolfson, and D. A. Low.** 1989. Phase-variation of pyelonephritis-associated pili in *Escherichia coli*: evidence for transcriptional regulation. *EMBO J.* **8**:613–620.
5. **Braaten, B. A., X. Nou, L. S. Kaltenbach, and D. A. Low.** 1994. Methylation patterns in *pap* regulatory DNA control pyelonephritis-associated pili phase variation in *E. coli*. *Cell* **76**:577–588.
6. **Brickman, T. J., and M. A. McIntosh.** 1992. Overexpression and purification of ferric enterobactin esterase from *Escherichia coli*. Demonstration of enzymatic hydrolysis of enterobactin and its iron complex. *J. Biol. Chem.* **267**:12350–12355.
7. **Brooks, C. S., P. S. Hefty, S. E. Jolliff, and D. R. Akins.** 2003. Global analysis of *Borrelia burgdorferi* genes regulated by mammalian host-specific signals. *Infect. Immun.* **71**:3371–3383.
8. **Caimano, M. J., C. H. Eggers, K. R. Hazlett, and J. D. Radolf.** 2004. RpoS is not central to the general stress response in *Borrelia burgdorferi* but does control expression of one or more essential virulence determinants. *Infect. Immun.* **72**:6433–6445.
9. **Casadaban, M.** 1976. Transposition and fusion of the *lac* genes to selected promoters in *E. coli* using bacteriophage lambda and Mu. *J. Mol. Biol.* **104**:541–555.
10. **Chapman, T. A., X. Y. Wu, I. Barchia, K. A. Bettelheim, S. Driesen, D. Trott, M. Wilson, and J. J. Chin.** 2006. Comparison of virulence gene profiles of *Escherichia coli* strains isolated from healthy and diarrheic swine. *Appl. Environ. Microbiol.* **72**:4782–4795.
11. **Cotter, P. A., and J. F. Miller.** 1998. *In vivo* and *ex vivo* regulation of bacterial virulence gene expression. *Curr. Opin. Microbiol.* **1**:17–26.
12. **Cumberlidge, A. G., and K. Isono.** 1979. Ribosomal protein modification in *Escherichia coli*. I. A mutant lacking the N-terminal acetylation of protein S5 exhibits thermosensitivity. *J. Mol. Biol.* **131**:169–189.
13. **Dorman, C. J.** 2004. H-NS: a universal regulator for a dynamic genome. *Nat. Rev. Microbiol.* **2**:391–400.
14. **Dorman, C. J., and N. N. Bhriain.** 1992. Thermal regulation of *fimA*, the *Escherichia coli* gene coding for the type 1 fimbrial subunit protein. *FEMS Microbiol. Lett.* **78**:125–130.
15. **Durant, L., A. Metais, C. Soulama-Mouze, J. M. Genevard, X. Nassif, and S. Esaich.** 2006. Identification of candidates for a subunit vaccine against extraintestinal pathogenic *Escherichia coli*. *Infect. Immun.* **75**:1916–1925.
16. **Earhart, C. F.** 1996. Uptake and metabolism of iron and molybdenum, p. 1075–1090. *In* F. C. Neidhardt, R. Curtis III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. ASM Press, Washington, DC.
17. **Edwards, R. A., L. H. Keller, and D. M. Schifferli.** 1998. Improved allelic exchange vectors and their use to analyze 987P fimbria gene expression. *Gene* **207**:149–157.
18. **Edwards, R. A., and D. M. Schifferli.** 1997. Differential regulation of *fasA* and *fasH* expression of *Escherichia coli* 987P fimbriae by environmental cues. *Mol. Microbiol.* **25**:797–809.
19. **Falconi, M., V. McGovern, C. Gualerzi, D. Hillyard, and N. P. Higgins.** 1991. Mutations altering chromosomal protein H-NS induce mini-Mu transposition. *New Biol.* **3**:615–625.
20. **Gally, D. L., J. A. Bogan, B. I. Eisenstein, and I. C. Blomfield.** 1993. Environmental regulation of the *fim* switch controlling type 1 fimbrial phase variation in *Escherichia coli* K-12: effects of temperature and media. *J. Bacteriol.* **175**:6186–6193.
21. **Göransson, M., K. Forsman, and B. E. Uhlin.** 1989. Regulatory genes in the thermoregulation of *Escherichia coli* pili gene transcription. *Genes Dev.* **3**:123–130.
22. **Göransson, M., B. Sonden, P. Nilsson, B. Dagberg, K. Forsman, K. Emanuelsson, and B. E. Uhlin.** 1990. Transcriptional silencing and thermoregulation of gene expression in *Escherichia coli*. *Nature* **344**:682–685.
23. **Göransson, M., and B. E. Uhlin.** 1984. Environmental temperature regulates transcription of a virulence pili operon in *E. coli*. *EMBO J.* **3**:2885–2888.
24. **Han, Y., D. Zhou, X. Pang, Y. Song, L. Zhang, J. Bao, Z. Tong, J. Wang, Z. Guo, J. Zhai, Z. Du, X. Wang, X. Zhang, J. Wang, P. Huang, and R. Yang.** 2004. Microarray analysis of temperature-induced transcriptome of *Yersinia pestis*. *Microbiol. Immunol.* **48**:791–805.
25. **Hantke, K., and V. Braun.** 2000. The art of keeping low and high iron concentrations in balance, p. 275–288. *In* G. Storz and R. Hengge-Aronis (ed.), *Bacterial stress responses*. ASM Press, Washington, DC.
26. **Jordi, B. J., B. Dagberg, L. A. de Haan, A. M. Hamers, B. A. van der Zeijst, W. Gastra, and B. E. Uhlin.** 1992. The positive regulator CfaD overcomes the repression mediated by histone-like protein H-NS (H1) in the CFA/I fimbrial operon of *Escherichia coli*. *EMBO J.* **11**:2627–2632.
27. **Karjalainen, T. K., D. G. Evans, D. J. Evans, Jr., D. Y. Graham, and C. H. Lee.** 1991. Iron represses the expression of CFA/I fimbriae of enterotoxigenic *E. coli*. *Microb. Pathog.* **11**:317–323.
28. **Krabbe, M., N. Weyand, and D. Low.** 2000. Environmental control of pilus gene expression, p. 305–322. *In* G. Storz and R. Hengge-Aronis (ed.), *Bacterial stress responses*. ASM Press, Washington, DC.
29. **Lafontaine, E. R., and P. A. Sokol.** 1998. Effects of iron and temperature on expression of the *Pseudomonas aeruginosa* *tolQRA* genes: role of the ferric uptake regulator. *J. Bacteriol.* **180**:2836–2841.
30. **Leveille, S., M. Caza, J. R. Johnson, C. Clabots, M. Sabri, and C. M. Dozois.** 2006. Iha from an *Escherichia coli* urinary tract infection outbreak clonal group A strain is expressed *in vivo* in the mouse urinary tract and functions as a catechololate siderophore receptor. *Infect. Immun.* **74**:3427–3436.
31. **Liu, P. Q., and G. F. Ames.** 1998. *In vitro* disassembly and reassembly of an ABC transporter, the histidine permease. *Proc. Natl. Acad. Sci. USA* **95**:3495–3500.
32. **Livak, K. J., and T. D. Schmittgen.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* **25**:402–408.
33. **Mahan, M. J., J. M. Schlauch, and J. J. Mekalanos.** 1996. Environmental regulation of virulence gene expression in *Escherichia*, *Salmonella*, and *Shigella* spp., p. 2803–2816. *In* F. C. Neidhardt, R. Curtis III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M.

- Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. ASM Press, Washington, DC.
34. Martínez-Laguna, Y., E. Calva, and J. L. Puente. 1999. Autoactivation and environmental regulation of *bfpT* expression, the gene coding for the transcriptional activator of *bfpA* in enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **33**:153–166.
 35. McHugh, J. P., F. Rodríguez-Quinones, H. Abdul-Tehrani, D. A. Svis-tunenko, R. K. Poole, C. E. Cooper, and S. C. Andrews. 2003. Global iron-dependent gene regulation in *Escherichia coli*. A new mechanism for iron homeostasis. *J. Biol. Chem.* **278**:29478–29486.
 36. Mekalanos, J. J. 1992. Environmental signals controlling expression of virulence determinants in bacteria. *J. Bacteriol.* **174**:1–7.
 37. Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratories, Cold Spring Harbor, NY.
 38. Motin, V. L., A. M. Georgescu, J. P. Fitch, P. P. Gu, D. O. Nelson, S. L. Mabery, J. B. Garnham, B. A. Sokhansanj, L. L. Ott, M. A. Coleman, J. M. Elliott, L. M. Kegelmeyer, A. J. Wyrobek, T. R. Slezak, R. R. Brubaker, and E. Garcia. 2004. Temporal global changes in gene expression during temperature transition in *Yersinia pestis*. *J. Bacteriol.* **186**:6298–6305.
 39. Nagy, L. K., T. Mackenzie, D. J. Pickard, and G. Dougan. 1986. Effects of immune colostrum on the expression of a K88 plasmid encoded determinant: Role of plasmid stability and influence of phenotypic expression of K88 fimbriae. *J. Gen. Microbiol.* **132**:2497–2503.
 40. Nasser, W., M. Faelen, N. Hugouvieux-Cotte-Pattat, and S. Reverchon. 2001. Role of the nucleoid-associated protein H-NS in the synthesis of virulence factors in the phytopathogenic bacterium *Erwinia chrysanthemi*. *Mol. Plant-Microbe Interact.* **14**:10–20.
 41. Nasser, W., and S. Reverchon. 2002. H-NS-dependent activation of pectate lyases synthesis in the phytopathogenic bacterium *Erwinia chrysanthemi* is mediated by the PecT repressor. *Mol. Microbiol.* **43**:733–748.
 42. Nikaido, H., and E. Y. Rosenberg. 1990. Cir and Fiu proteins in the outer membrane of *Escherichia coli* catalyze transport of monomeric catechols: study with beta-lactam antibiotics containing catechol and analogous groups. *J. Bacteriol.* **172**:1361–1367.
 43. Normark, S., D. Lark, R. Hull, M. Norgren, M. Bøga, P. O'Hanley, G. Schoolnik, and S. Falkow. 1983. Genetics of digalactoside-binding adhesion from a uropathogenic *Escherichia coli*. *Infect. Immun.* **41**:942–949.
 44. O'Hanley, P., D. A. Low, I. Romero, D. Lark, K. Vosti, S. Falkow, and G. Schoolnik. 1985. Gal-Gal binding and hemolysin phenotypes and genotypes associated with uropathogenic *Escherichia coli*. *N. Engl. J. Med.* **313**:414–420.
 45. Ojaimi, C., C. Brooks, S. Casjens, P. Rosa, A. Elias, A. Barbour, A. Jasinskas, J. Benach, L. Katona, J. Radolf, M. Caimano, J. Skare, K. Swingle, D. Akins, and I. Schwartz. 2003. Profiling of temperature-induced changes in *Borrelia burgdorferi* gene expression by using whole genome arrays. *Infect. Immun.* **71**:1689–1705.
 46. Pao, S. S., I. T. Paulsen, and M. H. Saier, Jr. 1998. Major facilitator superfamily. *Microbiol. Mol. Biol. Rev.* **62**:1–34.
 47. Perna, N. T., G. Plunkett III, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grotbeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamouis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner. 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**:529–533.
 48. Pilsil, H., D. Smajs, and V. Braun. 1999. Characterization of colicin S4 and its receptor, OmpW, a minor protein of the *Escherichia coli* outer membrane. *J. Bacteriol.* **181**:3578–3581.
 49. Postma, P. W., J. W. Lengeler, and G. R. Jacobson. 1993. Phosphoenolpyruvate:carbohydrate phosphotransferase systems of bacteria. *Microbiol. Rev.* **57**:543–594.
 50. Puente, J. L., D. Bieber, S. W. Ramer, W. Murray, and G. K. Schoolnik. 1996. The bundle-forming pili of enteropathogenic *Escherichia coli*: transcriptional regulation by environmental signals. *Mol. Microbiol.* **20**:87–100.
 51. Redford, P., P. L. Roesch, and R. A. Welch. 2003. DegS is necessary for virulence and is among extraintestinal *Escherichia coli* genes induced in murine peritonitis. *Infect. Immun.* **71**:3088–3096.
 52. Revel, A. T., A. M. Talaat, and M. V. Norgard. 2002. DNA microarray analysis of differential gene expression in *Borrelia burgdorferi*, the Lyme disease spirochete. *Proc. Natl. Acad. Sci. USA* **99**:1562–1567.
 53. Roos, V., and P. Klemm. 2006. Global gene expression profiling of the asymptomatic bacteriuria *Escherichia coli* strain 83972 in the human urinary tract. *Infect. Immun.* **74**:3565–3575.
 54. Russo, T. A., U. B. Carlino, A. Mong, and S. T. Jodush. 1999. Identification of genes in an extraintestinal isolate of *Escherichia coli* with increased expression after exposure to human urine. *Infect. Immun.* **67**:5306–5314.
 55. Russo, T. A., C. D. McFadden, U. B. Carlino-MacDonald, J. M. Beanan, T. J. Barnard, and J. R. Johnson. 2002. IronN functions as a siderophore receptor and is a urovirulence factor in an extraintestinal pathogenic isolate of *Escherichia coli*. *Infect. Immun.* **70**:7156–7160.
 56. Schmoll, T., M. Ott, B. Oudega, and J. Hacker. 1990. Use of a wild-type gene fusion to determine the influence of environmental conditions on expression of the S fimbrial adhesin in an *Escherichia coli* pathogen. *J. Bacteriol.* **172**:5103–5111.
 57. Serina, S., F. Nozza, G. Nicastro, F. Faggioni, H. Mottl, G. Deho, and A. Polissi. 2004. Scanning the *Escherichia coli* chromosome by random transposon mutagenesis and multiple phenotypic screening. *Res. Microbiol.* **155**:692–701.
 58. Silhavy, T. J., M. L. Berman, and L. W. Enquist. 1984. Experiments with gene fusions. Cold Spring Harbor Laboratories, Cold Spring Harbor, NY.
 59. Sirko, A., M. Zatyka, E. Sadowy, and D. Hulanicka. 1995. Sulfate and thiosulfate transport in *Escherichia coli* K-12: evidence for a functional overlapping of sulfate- and thiosulfate-binding proteins. *J. Bacteriol.* **177**:4134–4136.
 60. Smoot, L. M., J. C. Smoot, M. R. Graham, G. A. Somerville, D. E. Sturdevant, C. A. Migliaccio, G. L. Sylva, and J. M. Musser. 2001. Global differential gene expression in response to growth temperature alteration in group A *Streptococcus*. *Proc. Natl. Acad. Sci. USA* **98**:10416–10421.
 61. Snyder, J. A., B. J. Haugen, E. L. Buckles, C. V. Lockett, D. E. Johnson, M. S. Donnenberg, R. A. Welch, and H. L. Mobley. 2004. Transcriptome of uropathogenic *Escherichia coli* during urinary tract infection. *Infect. Immun.* **72**:6373–6381.
 62. Steele, M., K. Ziebell, Y. Zhang, A. Benson, P. Konczyk, R. Johnson, and V. Gannon. 2006. Identification of *Escherichia coli* O157:H7 genomic regions conserved in genotypes associated with human infection. *Appl. Environ. Microbiol.*
 63. Tatusov, R. L., E. V. Koonin, and D. J. Lipman. 1997. A genomic perspective on protein families. *Science* **278**:631–637.
 64. Torres, A. G., P. Redford, R. A. Welch, and S. M. Payne. 2001. TonB-dependent systems of uropathogenic *Escherichia coli*: aerobactin and heme transport and TonB are required for virulence in the mouse. *Infect. Immun.* **69**:6179–6185.
 65. Tusher, V. G., R. Tibshirani, and G. Chu. 2001. Significance analysis of microarrays applied to the ionizing radiation response. *Proc. Natl. Acad. Sci. USA* **98**:5116–5121.
 66. Ussery, D. W., J. C. D. Hinton, B. J. A. M. Jordi, P. E. Granum, A. Seirafi, R. J. Stephen, A. E. Tupper, G. Berridge, J. M. Sidebotham, and C. F. Higgins. 1994. The chromatin-associated protein H-NS. *Biochimie* **76**:968–980.
 67. van der Woude, M. W., L. S. Kaltenbach, and D. A. Low. 1995. Leucine-responsive regulatory protein plays dual roles as both an activator and a repressor of the *Escherichia coli* *pap* fimbrial operon. *Mol. Microbiol.* **17**:303–312.
 68. Weinstein, D. L., R. K. Holmes, and A. D. O'Brien. 1988. Effects of iron and temperature on Shiga-like toxin I production by *Escherichia coli*. *Infect. Immun.* **56**:106–111.
 69. Welch, R. A., V. Burland, G. Plunkett III, P. Redford, P. Roesch, D. Rasko, E. L. Buckles, S. R. Liou, A. Boutin, J. Hackett, D. Stroud, G. F. Mayhew, D. J. Rose, S. Zhou, D. C. Schwartz, N. T. Perna, H. L. Mobley, M. S. Donnenberg, and F. R. Blattner. 2002. Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **99**:17020–17024.
 70. White-Ziegler, C. A., M. L. Angus Hill, B. A. Braaten, M. W. van der Woude, and D. A. Low. 1998. Thermoregulation of *Escherichia coli* *pap* transcription: H-NS is a temperature-dependent DNA methylation blocking factor. *Mol. Microbiol.* **28**:1121–1137.
 71. White-Ziegler, C. A., A. M. Black, S. H. Eliades, S. Young, and K. Porter. 2002. The *N*-acetyltransferase RimJ responds to environmental stimuli to repress *pap* fimbrial transcription in *Escherichia coli*. *J. Bacteriol.* **184**:4334–4342.
 72. White-Ziegler, C. A., L. B. Blyn, B. A. Braaten, and D. A. Low. 1990. Identification of a genetic locus in *Escherichia coli* involved in the thermoregulation of the *pap* operon. *J. Bacteriol.* **172**:1775–1782.
 73. White-Ziegler, C. A., and D. A. Low. 1992. Thermoregulation of the *pap* operon: evidence for the involvement of RimJ, the *N*-terminal acetylase of ribosomal protein S5. *J. Bacteriol.* **174**:7003–7012.
 74. White-Ziegler, C. A., A. Villapakkam, K. Ronaszeki, and S. D. Young. 2000. H-NS controls *pap* and *daa* fimbrial transcription *Escherichia coli* in response to multiple environmental cues. *J. Bacteriol.* **182**:6391–6400.
 75. Williams, R. M., and S. Rimsky. 1997. Molecular aspects of the *E. coli* nucleoid protein, H-NS: a central controller of gene regulatory networks. *FEMS Microbiol. Lett.* **156**:175–185.